## **List of Typically Avoided Situations for Trauma Survivors**

Some typical examples of upsetting situations for trauma survivors that usually lead to avoidance include:

- In cases of assault, being around men you don't know, especially those who somehow remind you of the assailant (e.g., a man of the same race)
- 2. Someone standing close or coming up suddenly
- 3. Being touched by someone (especially someone unfamiliar)
- 4. Doing something similar to the trauma situation (e.g., for motor vehicle accident survivors, driving or riding in a car)
- 5. Walking down a street or being out in the open
- 6. Being alone at home (day or night)
- 7. Going somewhere alone at night
- 8. Being in a crowded mall or store
- 9. Talking to strangers
- 10. Driving a car or being stopped at a stoplight
- 11. Being in a parking lot
- 12. Riding in elevators or being in small, confined spaces
- 13. Reading about a similar event in the newspaper or hearing about it on television
- 14. Talking with someone about the trauma
- 15. Watching movies that remind you of the trauma (e.g., combat films, assault scenes)
- 16. Going to the area of town where the traumatic event happened
- 17. Riding public transportation
- 18. Hugging and kissing significant others
- 19. Sexual or physical contact

- 20. Listening to a song that you heard during the traumatic event or that was from that same time
- 21. Watching the news on TV
- 22. Wearing makeup or looking attractive
- 23. Going to a movie that has some violence
- 24. Taking an exercise class
- 25. Driving a car with unknown people and piles of unidentifiable material next to the road that you fear could be explosives
- 26. Coming upon a stopped car while driving and not being able to get around it
- 27. Smelling food or spices similar to those present during the traumatic experience

## Safety Considerations When Constructing the In Vivo Exposure Hierarchy

It is important that the situations you plan to confront during in vivo exposure are safe or low-risk. We do not want you to put situations that are dangerous or high-risk on your list. For example, we would not want you to walk alone in areas where drugs are sold or in a park where crimes are regularly committed. Instead, we want you to pick situations that trigger your anxiety but are realistically safe. For example, you could walk in a public park with another person, or plan to walk alone in a safe area of the city.