

The Endocrine Society's
CLINICAL | GUIDELINES

Evaluation and Treatment of
Adult Growth Hormone Deficiency:
An Endocrine Society Clinical Practice Guideline



Growth Hormone Guideline Task Force: Mark E. Molitch, David R. Clemmons, Saul Malozowski, George R. Merriam, Stephen M. Shalet, and Mary Lee Vance, for The Endocrine Society's Clinical Guidelines Subcommittee

Northwestern University Feinberg School of Medicine (M.E.M.), Chicago, Illinois; University of North Carolina School of Medicine (D.R.C.), Chapel Hill, North Carolina; National Institute of Diabetes, Digestive and Kidney Disease (S.M.), National Institutes of Health, Bethesda, Maryland; University of Washington, Veterans Affairs Puget Sound Health Care System (G.R.M.), Seattle and Tacoma, Washington; Christie Hospital (S.M.S.), Manchester, United Kingdom; and University of Virginia Health Science Center (M.L.V.), Charlottesville, Virginia

Disclaimer: Clinical Practice Guidelines are developed to be of assistance to endocrinologists by providing guidance and recommendations for particular areas of practice. The Guidelines should not be considered inclusive of all proper approaches or methods, or exclusive of others. The Guidelines cannot guarantee any specific outcome, nor do they establish a standard of care. The Guidelines are not intended to dictate the treatment of a particular patient. Treatment decisions must be made based on the independent judgment of healthcare providers and each patient's individual circumstances.

The Endocrine Society makes no warranty, express or implied, regarding the Guidelines and specifically excludes any warranties of merchantability and fitness for a particular use or purpose. The Society shall not be liable for direct, indirect, special, incidental, or consequential damages related to the use of the information contained herein.

First published in *The Journal of Clinical Endocrinology & Metabolism*, Vol. 91, pages 1621–1634.

© The Endocrine Society, 2006



The Endocrine Society's
CLINICAL | GUIDELINES

Evaluation and Treatment of
Adult Growth Hormone Deficiency:
An Endocrine Society Clinical Practice Guideline



Table of Contents

Introduction	4
GHD in Adults	5
AGHD in patients with prior childhood GHD	
Acquired GHD secondary to structural lesions or trauma	
Adult-onset idiopathic GHD	
Recommendations	
Diagnosis of GHD	8
Recommendations	
Suggestions	
Consequences of GHD and Benefits of Treatment with GH	10
Body composition	
Bone health	
Cardiovascular health	
Mortality in hypopituitarism	
Quality of life	
Recommendations	
Side Effects and Risks Associated with GH Therapy	14
Recommendation	
Suggestion	
Treatment Regimens	16
Recommendations	
Conclusion, Summary of Recommendations & Suggestions	18
Appendix	20
References	21
Order Form	31
Reprint Information, Questions & Correspondences	Inside Back Cover

Abstract

Objective: The objective is to provide guidelines for the evaluation and treatment of adults with GH deficiency (GHD).

Participants: The chair of the Task Force was selected by the Clinical Guidelines Subcommittee of The Endocrine Society (TES). The chair selected five other endocrinologists and a medical writer, who were approved by TES Council. One closed meeting of the group was held. There was no corporate funding, and members of the group received no remuneration.

Evidence: Only fully published, peer-reviewed literature was reviewed. The grades of evidence used are outlined in the *Appendix*.

Consensus Process: Consensus was achieved through one group meeting and e-mailing of drafts that were written by the group with grammatical/style help from

the medical writer. Drafts were reviewed successively by the Clinical Guidelines Subcommittee, the Clinical Affairs Committee, and TES Council, and a version was placed on the TES Web site for comments. At each level, the task force incorporated needed changes.

Conclusions: GHD can persist from childhood or be newly acquired. Confirmation through stimulation testing is usually required unless there is a proven genetic/structural lesion persistent from childhood. GH therapy offers benefits in body composition, exercise capacity, skeletal integrity, and quality of life measures and is most likely to benefit those patients who have more severe GHD. The risks of GH treatment are low. GH dosing regimens should be individualized. The final decision to treat adults with GHD requires thoughtful clinical judgment with a careful evaluation of the benefits and risks specific to the individual. (*J Clin Endocrinol Metab* 91: 0000–0000, 2006)

Introduction

When GH Therapy was first introduced in the 1950s, the preparation was extracted from human cadaver pituitaries and therefore was restricted in its supply (1). Extracted preparations of GH were used for the treatment of children with GH deficiency (GHD) until 1985, when its use was halted worldwide because of its association with cases of Creutzfeldt-Jakob disease. Since 1985, all GH in clinical use has been recombinant DNA-derived biosynthetic human GH that is free of the Creutzfeldt-Jakob prions and is available from several pharmaceutical firms. The current preparations have a biopotency of 3 IU/mg, using the World Health Organization (WHO) reference preparation 88/624 (2).

Before biosynthetic GH became available, treatment was limited to children with short stature and proven GHD. Now GH is approved by the U.S. Food and Drug Administration (FDA) and other regulatory agencies for treatment of short stature from causes other than GHD, such as Turner's syndrome, renal failure, small size for gestational age, Prader-Willi syndrome, and, most recently, idiopathic short stature. Because GH was only the second (after insulin) recombinant DNA preparation to be approved, the FDA required all manufacturers to carry out postmarketing safety surveillance studies. In most circumstances, pediatric GH therapy has been discontinued once final height has been achieved after epiphyseal closure. Issues regarding continuation of GH treatment in GH-deficient children will be discussed below.

GH treatment of GH-deficient adults was approved by the FDA in 1996. Subsequently, much clinical experience has been evaluated regarding the treatment of adults with GHD. Although treatment appears to be

safe overall, certain areas require long-term surveillance, such as risks of glucose intolerance, pituitary/hypothalamic tumor recurrence, and cancer (3). Benefits of GH treatment of GH-deficient adults have been found in body composition, bone health, cardiovascular risk factors, and quality of life indicators. However, reductions in cardiovascular events and mortality have yet to be demonstrated, and treatment costs remain high.

It is the purpose of these guidelines to summarize information regarding adult GHD (AGHD) and to provide recommendations related to several specific questions:

- Who are appropriate candidates for GH therapy?
- What tests should be used to diagnose GHD, and what criteria are necessary to make this diagnosis with the various tests?
- What are the benefits of treatment with GH in GH-deficient adults?
- What are the risks of treatment with GH in GH-deficient adults?
- What treatment regimens should be used, and how should these be monitored?

The evidence used for these recommendations has been graded according to uniform criteria (see *Appendix*). Greater weight was given to randomized, placebo-controlled trials than to open-label, uncontrolled studies. It should be emphasized at the outset that the decision to treat adults with GHD requires thoughtful clinical judgment. For each potential patient, a careful evaluation of the benefits and risks is required before a decision to treat can be reached. Furthermore, if there is a decision to treat, then periodic reevaluation of treatment is warranted.

Abbreviations: AGHD, Adult GHD; BMD, bone mineral density; DXA, dual energy x-ray absorptiometry; GHD, GH deficiency; IGHD, isolated GHD; IMT, intimal-medial thickness; ITT, insulin tolerance test; LDL, low-density lipoprotein; LV, left ventricular; MPH, multiple pituitary hormone deficiency; SMR, standardized mortality ratio.

GHD in Adults

Adults with GHD can be grouped into three categories: those who had prior childhood GHD, those who acquire GHD secondary to structural lesions or trauma, and those with idiopathic GHD. Childhood GHD is generally further divided into those with organic causes and those with idiopathic causes.

AGHD in patients with prior childhood GHD

Mutations in early-appearing transcription factors tend to cause multiple pituitary hormone deficiencies (MPHD; e.g. in mutations of HESXI, PROPI, PIT-I, and LHX3/4), whereas others can cause isolated deficiencies (e.g. GHD in Rieger syndrome due to PITX2 mutation) (4,5) (Table 1).

Four types of Mendelian disorders of the GH gene have been described (6). Isolated GHD (IGHD) IA and IB are both inherited in an autosomal recessive manner resulting in undetectable or very low GH levels. Patients with undetectable GH (IGHD IA) levels often develop anti-GH antibodies when treated with GH. IGHD II is inherited in an autosomal dominant manner with variable clinical severity. IGHD III is an X-linked disorder often associated with hypogammaglobulinemia.

Mutations of the gene encoding the GHRH receptor have been identified in a number of kindreds with GHD (7). Mutations in the GSA gene can lead to GHRH resistance and thereby GHD (8).

GHD is occasionally associated with congenital anatomical changes in the pituitary region or other structures of the brain, as well as with malformations outside the neurocranium, usually in association

with other pituitary hormone deficiencies (9–12), as listed in Table 1.

The term “empty sella syndrome” is descriptive, signifying primarily that the sella turcica does not contain normal pituitary structures; however, the radiological demonstration of an empty sella does not provide insight into pituitary function. In many cases, the anatomical abnormalities are associated with endocrine dysfunction, including both IGHD and MPHD (13). An empty sella was observed in 8.8% of patients with IGHD and 34.9% of those with MPHD (13).

Tumors in the pituitary and hypothalamic area may cause hypopituitarism primarily or after treatment with surgery and/or irradiation. The most common tumors are pituitary adenomas and craniopharyngiomas; others are listed in Table 1.

Infiltrative diseases (Langerhans cell histiocytosis, sarcoidosis, tuberculosis) of the hypothalamus and stalk commonly cause hypopituitarism and diabetes insipidus. Lymphocytic hypophysitis may involve the pituitary and stalk.

GH status evolves with time after cranial radiotherapy and depends on dose. The younger the patient, the longer the interval after radiotherapy, and the higher the dose, the greater the chance of developing GHD after irradiation. There is a greater than 50% likelihood of GHD if the biological effective dose is greater than 40 Gy (14).

In nearly all series, idiopathic is the category that accounts for most individuals with childhood GHD (15–21). In these studies, all patients were documented biochemically to be GH deficient in childhood, but at

reassessment in adulthood most had normal GH responses when tested. This finding raises interesting questions about the nature of the defect in GH secretion during childhood in this group. The diagnostic threshold for GHD is arbitrarily defined, and there is poor reproducibility of the GH response to provocative testing within individuals. On these grounds alone, it would be anticipated that a considerable number of those considered GH deficient at one time might be normal at reevaluation. Furthermore, it is likely that in a proportion of these patients, the childhood diagnosis was constitutional delay in growth and puberty and not isolated idiopathic GHD, but the initial GH provocative tests carried out without estrogen “priming” failed to make this distinction (21). Finally, it remains possible that transient GHD in childhood is a real entity, although longitudinally obtained proof is lacking. Because of the greater GH requirements for normal growth in children, it is possible that in some patients the GHD was partial but severe enough to prevent normal growth as a child and not severe enough to cause symptoms or meet criteria for GHD as an adult.

In contrast to the population with isolated idiopathic GHD, young adults diagnosed as having organic GHD in childhood, as a consequence of a mass lesion, pituitary surgery, high-dose irradiation damage to the hypothalamic-pituitary axis, or a combination of these, much less commonly revert to normal GH status (20). Those with genetic and embryopathic defects do not revert to normal GH status.

Acquired GHD secondary to structural lesions or trauma

The most common cause of GHD in adults is a pituitary adenoma or treatment of the adenoma with pituitary surgery or radiotherapy. Pituitary microadenomas are very rarely, if ever, associated with hypopituitarism. Macroadenomas are more frequently associated with pituitary hormone deficiencies, with 30-60% having one or more anterior pituitary hormone deficiencies. The likely mechanism of hypopituitarism in most patients is compression of the portal vessels in the pituitary stalk, secondary to either the expanding tumor

mass directly or raised intrasellar pressure (22). Derangement of central endocrine regulation also occurs with parapituitary space-occupying lesions such as craniopharyngiomas, Rathke's cleft cysts, arachnoid cysts, meningiomas, dysgerminomas, metastatic tumors, and astrocytomas/gliomas.

Hypopituitarism can be a consequence of pituitary surgery. The incidence and degree of hypopituitarism depend on several factors, including the size of the original tumor, the degree of infiltration, and the experience of the surgeon. Surgery for pituitary adenomas may also be followed by significant recovery of pituitary function; up to 50% of patients recover at least one pituitary hormone that had been deficient after transsphenoidal surgery (23–25). Postoperative improvement is more likely if there is no tumor on postoperative imaging and no neurosurgical or pathological evidence that the tumor is invasive (23). GH is less likely to recover than gonadotropins, ACTH, and TSH (25). There is evidence that among those patients in whom recovery of pituitary function occurs, the process begins immediately after surgery (24).

Irradiation commonly causes hypopituitarism and may be progressive over time. By 10 yr after conventional, fractionated irradiation, varying degrees of hypopituitarism are present in over 50% of patients (26, 27). Single dose, stereotactic radiotherapy also leads to hypopituitarism, and preliminary data suggest a similar rate (28).

Traumatic brain injury has been reported to cause GHD and varying degrees of hypopituitarism in more than 25% of patients (29–32). The best time to assess pituitary function in such patients has not been determined.

Adult-onset idiopathic GHD

As defined by strict hormonal criteria, adult-onset idiopathic GHD is very rare. There is no single biological marker in an adult who is suspected of being GH deficient that offers the same diagnostic usefulness as the growth rate of a child. GH is usually the first of the anterior pituitary hormones to be affected by

pathological insults. Consequently, in a patient with MPHD, the probability of GHD is extremely high. This conclusion could be applied to those patients with idiopathic MPHD in just the same manner as it has been for those with an organic cause for their MPHD. However, no studies documenting such a transition from isolated to multiple hormone losses have been reported.

The more difficult issue concerns the patient in whom a diagnosis of isolated idiopathic GHD of adult onset is being considered. Truncal obesity will be present, and it is now established in clinically nonobese healthy adults that relative adiposity, in the abdominal region in particular, is associated with a blunted GH response to stimulation (33, 34); hence, GH status will often appear to be subnormal. Obesity *per se* is almost always associated with a normal IGF-I level. Therefore, to

make the diagnosis of idiopathic GHD, an IGF-I level below the age-corrected lower limit of normal should also be present.

Recommendations

- Patients with childhood-onset GHD who are candidates for GH therapy should be retested for GHD as adults unless they have known mutations, embryopathic lesions, or irreversible structural lesions/ damage (level of evidence, high).
- Adult patients with structural hypothalamic/pituitary disease, surgery or irradiation in these areas, or other pituitary hormone deficiencies should be considered for evaluation for acquired GHD (level of evidence, high).

TABLE 1. Causes of GHD

Congenital

Genetic
Transcription factor defects (PIT-1, PROP-1, LHX3/4, HESX-1, PITX-2)
GHRH receptor gene defects
GH secretagogue receptor gene defects
GH gene defects
GH receptor/post receptor defects
Prader-Willi syndrome
Associated with brain structural defects
Agenesis of corpus callosum
Septo-optic dysplasia
Empty sella syndrome
Holoprosencephaly
Encephalocele
Hydrocephalus
Arachnoid cyst
Associated with midline facial defects
Single central incisor
Cleft lip/palate

Acquired

Trauma
Perinatal
Postnatal
Central nervous system infection
Tumors of hypothalamus or pituitary
Pituitary adenoma
Craniopharyngioma
Rathke's cleft cyst
Glioma/astrocytoma
Germinoma
Metastatic
Other
Infiltrative/granulomatous disease
Langerhans cell histiocytosis
Sarcoidosis
Hypophysitis
Other
Cranial irradiation
Surgery
Idiopathic

Diagnosis of GHD

Clinically, adults with GHD tend to have a relative increase in fat mass and a relative decrease in muscle mass and, in many instances, decreased energy and quality of life. These characteristics are obviously nonspecific but, in the appropriate clinical setting (see above), may point to GHD. The next step in such an evaluation is hormonal testing.

GH is secreted in an episodic manner that is modified by age and sex. Because of this pulsatile secretion, multiple sampling of GH levels would be ideal, but it is not a practical procedure in clinical practice. For this reason, current diagnostic testing involves provocative tests of GH secretion. However, these tests have intrinsic false-positive error rates. Additionally, the insulin tolerance test (ITT), which has been considered the “gold standard,” may carry increased risk in patients with seizure disorders or cardiovascular disease and requires constant monitoring even in healthy adults, although it is quite safe in experienced hands. Ghigo and colleagues (35) showed that the combined administration of arginine, which presumably reduces hypothalamic somatostatin secretion, and GHRH is safe and provides a strong stimulus to GH secretion that is less affected by aging or obesity and thus could be used as an alternative to the ITT as a test of pituitary GHD.

A recent study evaluated the relative performance of GHRH-arginine, the ITT, arginine alone, clonidine, levodopa, and the combination of arginine plus levodopa (36). The five tests were administered in random order to 39 patients with MPPHD; to 21 patients with one or no pituitary deficiency other than GH; and to 34 sex-, age-, and body mass index-matched controls. The overall performance of the GHRH-arginine test, with 95% sensitivity and 91% specificity at a GH cutoff of 4.1 $\mu\text{g/liter}$ at the central laboratory used, compared well to the ITT, which had an optimal GH cutoff of 5.1

$\mu\text{g/liter}$ (96% sensitivity and 92% specificity). The performance of the other tests was much poorer. As expected, the discriminating power of all tests was reduced in patients with fewer pituitary hormone deficits, *i.e.* the patients posing the greatest diagnostic challenge, but again the GHRH-arginine test performed almost as well as the ITT. Because the GHRH-arginine test is generally well tolerated and free of the potentially serious side effect of hypoglycemia, it is beginning to gain wider use for patients with suspected GHD of pituitary origin, the majority of AGHD patients. However, because GHRH directly stimulates the pituitary, it can give a falsely normal GH response in patients with GHD of hypothalamic origin, *e.g.* those having received irradiation of the hypothalamic-pituitary region (37). In such a circumstance, if hypoglycemic stimulation testing is contraindicated, arginine alone may be used without concomitant GHRH, using a lower cutoff of 1.4 $\mu\text{g/liter}$ (36).

Having normal levels of IGF-I and IGFBP-3 does not exclude a diagnosis of GHD in adults (36, 38–40). However, IGF-I can be of some diagnostic assistance if levels are below the age-adjusted normal range.

Several studies involving panhypopituitary patients have shown that under certain circumstances GH stimulation tests may be unnecessary to diagnose AGHD (39, 41, 42). In one study, the proportion of patients with GH responses to provocative testing that were below 2.5 $\mu\text{g/ml}$ increased with the number of other pituitary hormone deficiencies. The presence of three or more other deficits, together with a low serum IGF-I level (<84 ng/ml in the Esoterix assay), was as specific a predictor as any of the GH provocative tests employed (39). Thus, one might conclude that GH testing could be omitted in these patients. Not all insurers' requirements, however, have been modified to

reflect this information, and many still require the results of a GH stimulation test.

Biochemical criteria for the diagnosis of AGHD are complicated by the lack of normative data that are age- and sex-adjusted, by assay variability, and by the stimulus used. With polyclonal RIAs, the cutoff values for stimulated GH levels for diagnosing AGHD were established at levels between 3 and 5 $\mu\text{g/liter}$ (43). Whether lower cutoffs should be used with the newer, more sensitive, two-site assays has not been definitively determined. Still, according to the multicenter study cited above (36), which used an immunochemiluminescent two-site assay, the values of 5.1 $\mu\text{g/liter}$ for the ITT and 4.1 $\mu\text{g/liter}$ for GHRH-arginine test had sufficient specificity and sensitivity for the diagnosis of AGHD.

The transition from pediatric to adult care is an appropriate time for reassessment of GH status. Patients with a high likelihood of having permanent GHD are those who have MPHD and a serum IGF-I concentration below the normal range if associated with one or more of the following: 1) a radiologically confirmed congenital anomaly in the sellar or suprasellar region; 2) known acquired hypothalamic-pituitary disease, *e.g.* craniopharyngioma; 3) previous surgery for lesions directly affecting the hypothalamic-pituitary region or radiotherapy for malignant disease that included a high dose of irradiation to the hypothalamic-pituitary region; or 4) a proven genetic/molecular defect involving the capacity to secrete GH. If children in these categories have a low IGF-I level on no GH treatment, this generally suffices to document continuing GHD. Those children with idiopathic GHD, either isolated or with one additional hormone deficit, are less likely to have permanent GHD and should be retested in early adulthood using the stimulation tests and criteria outlined above.

Testing should be conducted after discontinuation of GH treatment to avoid possible suppression of endogenous responses. The interval between the reevaluation and the discontinuation of GH should not be less than 1 month. (It should be noted that no formal studies have addressed this interval and this recommendation is based on personal practice.)

Recommendations

- The insulin tolerance and the GHRH-arginine tests have sufficient sensitivity and specificity to establish the diagnosis of GHD. However, in those with clearly established recent (within 10 yr) hypothalamic causes of suspected GHD, *e.g.* irradiation, testing with GHRH-arginine may be misleading (level of evidence, high).
- Because of the irreversible nature of the cause of the GHD in children with structural lesions with multiple hormone deficiencies and those with proven genetic causes, a low IGF-I level at least 1 month off GH therapy is sufficient documentation of persistent GHD without additional provocative testing (level of evidence, moderate).

Suggestions

- A normal IGF-I level does not exclude the diagnosis of GHD but makes provocative testing mandatory to make the diagnosis of GHD (level of evidence, high).
- A low IGF-I level, in the absence of catabolic conditions and liver disease, indicates severe GHD and may be useful in identifying patients who may benefit from treatment (level of evidence, moderate).
- The presence of deficiencies in three or more pituitary axes strongly suggests the presence of GHD, and in this context provocative testing is optional (level of evidence, moderate).

Consequences of GHD and Benefits of Treatment with GH

The benefits of treatment with GH among patients with GHD occur in several domains: body composition, bone health, cardiovascular risk factors, and quality of life. Mortality is increased in patients with hypopituitarism, and the role of GHD in this mortality will be discussed.

Body composition

One of the most consistent responses to GH administration is increased lipolysis. Before treatment, AGHD patients often have increased fat mass, and some studies have shown a preferential increase in visceral fat (44–46). Several studies have found significant decreases in total body fat content in response to GH therapy. This effect occurs in both sc and visceral fat. Using computerized tomography scanning, some investigators have reported a preferential effect of GH on visceral fat (47–50). This change occurs within 6 months after the initiation of therapy, and it is maintained if treatment is continued.

Untreated adults with GHD have also been shown to have decreased lean body mass compared with age- and sex-matched controls (44, 46, 51). There is usually an increase in muscle mass in response to GH; however, the degree of change is less than the reduction in fat mass (48, 52–54). Several studies have attempted to determine whether this change in muscle mass results in increased strength and/or exercise capacity. Some but not all studies have shown increases in isometric or isokinetic strength (48, 49, 55–59). Even in those studies showing an increase in strength, the absolute values attained do not equal those of control subjects without GHD. In some but not all short- and long-term studies, exercise capacity and physical performance were improved by GH therapy, with parameters such as

VO₂ max and maximum work capacity being significantly increased (59–62).

Bone health

Multiple studies using either bone densitometry [dual energy x-ray absorptiometry (DXA)] or quantitative computerized tomography have shown that, on average, bone mineral density (BMD) in adults severely deficient in GH is approximately 1 sd score below the mean (63–65), even when the possible effects of hypogonadism or glucocorticoid overreplacement are factored out (63, 64, 66). Approximately 20% of adult-onset and 35% of childhood-onset adult patients with GHD have BMD T-scores below –2.5 (the threshold for the diagnosis of osteoporosis). The age of onset of GHD appears to determine the severity of osteopenia. Whether their GHD is adult onset or childhood onset, patients younger than 30 yr have the most severe osteopenia, whereas subjects older than 60 yr do not differ from controls without GHD. Subjects between 30 and 45 yr of age have intermediate severity osteopenia (67). The severity of GHD correlates with the severity of osteopenia (68). GH-deficient children who do not receive replacement therapy during puberty and after reaching final height have reduced peak bone mass, which is not reached until a decade after linear growth ceases (69).

On histological examination of iliac crest bone biopsies, GHD patients show an increase in the volume of trabecular bone, increased resorption, and increased osteoid thickness, suggesting delayed mineralization (70). Fracture rates that are increased 2- to 5-fold compared with rates in non-GHD control populations have been reported (71–73). Levels of circulating and urinary markers of bone resorption and formation are

variable, however, and therefore not routinely used in clinical practice.

GH replacement has an eventual overall anabolic effect on bone, but the effects are complex and the results biphasic. GH stimulates both bone formation and resorption (74, 75). With less than 12 months of treatment, BMD by DXA scanning may not increase and may even decrease (75, 76). After 18-24 months of treatment, however, most studies have shown increases of 4–10% in BMD (usually by DXA), generally with greater effects at vertebral than at femoral sites (74, 77, 78). When subjects were stratified on the basis of severity of bone mineral loss, those subjects with Z scores worse than -2 had the greatest improvement in response to treatment (79). One characteristic that is apparent in almost all studies is that the BMD of men responds better to GH therapy than the BMD of women (80, 81). The effects of GH replacement on BMD may plateau after several years of treatment, and one study suggests that in patients who remain osteopenic adding a bisphosphonate may result in further improvement (82). This study suggests a beneficial effect of adding bisphosphonate therapy to GH on fracture risk; to date, however, there are no reports of controlled studies of the effects of long-term GH replacement on the fracture rate in AGHD patients.

Special considerations pertain to childhood-onset GHD and the transition to adulthood. Any evaluation of BMD in these patients has to take into account volumetric assessment. Some of these patients may not have reached their true potential maximum bone volume and may continue to increase bone volume with GH therapy. DXA scanning does not directly measure bone volume, although correction formulas can be applied (83).

An important question is whether these transition patients require GH replacement beyond the time of their reaching final height to achieve normal peak bone mass. In normal adults, 95% peak bone mass is achieved by the mid-twenties, occurring later in men than women (84). However, subjects with hypopituitarism due to delayed onset of puberty or lack of normal gonadotropin secretion may lag behind in terms of the age at which they reach peak bone mass (85). After

discontinuation of GH therapy, there is a reduced acquisition of bone mineral content (86–88). This discontinuation of therapy usually occurs at age 15–17 yr, an age when normal subjects are still increasing bone mass. An important issue is whether therapy should be maintained or reinstated at least until these subjects reach peak bone mass. Four studies have demonstrated that continuing/reinstating GH therapy for periods of up to 2 yr in patients who had completed growth resulted in significantly greater BMD than that in patients who had equally severe GHD but received no treatment (78, 81, 85, 89), but one study did not (90). Overall, these studies suggest, therefore, that patients with childhood-onset GHD who have low age-adjusted bone mineral content would benefit from continued treatment.

These findings suggest that GHD patients should have a DXA measurement of BMD before treatment and, if it is abnormal, at least every 2 yr thereafter. They particularly highlight the possible detrimental effects of stopping GH treatment for more than 18 months during the transition from pediatric to adult GH replacement, when linear growth has ceased but bone mass continues to accrue, and suggest that if GH treatment is interrupted at this time, retesting and reinstatement of transitional and then adult GH doses be completed as expeditiously as possible.

Cardiovascular health

GH has both direct effects on vascular function and effects mediated through IGF-I that may have opposing actions. In general, most of the cardiovascular risk that has been defined in patients with GHD appears to be related to hypertension, inflammation, dyslipidemia, and insulin resistance. In severe GHD, patients tend to be substantially more hypertensive, and this results in impaired vasodilatation responses to stress and/or exercise (91). Importantly, GH replacement therapy has been shown to increase flow-mediated dilatation and to reduce arterial stiffness (92). In several large trials, GH replacement has resulted in a slight decrease in blood pressure (93).

Inflammatory markers are elevated in patients with GHD (94–96). Administration of GH reduces C-reactive protein concentrations, and this effect persists for as long as 18 months (94, 96). GH also affects lipoprotein metabolism. Increased total and low-density lipoprotein (LDL) cholesterol, decreased high-density lipoprotein cholesterol, and elevated apolipoprotein B-100 have been reported in 26–45% of GH-deficient adults (97). Most but not all studies have shown increases in high-density lipoprotein and decreases in LDL and total cholesterol after institution of GH replacement therapy (50, 53, 90, 93, 98–100). However, no studies have determined whether GH has an additive effect over and above optimum therapy with statins; therefore this remains an open question.

A fourth cardiovascular parameter is insulin resistance. The net effect of GH replacement on insulin resistance is difficult to predict. GH replacement lowers fat mass, and increasing IGF-I improves insulin sensitivity (101), but GH also has direct insulin antagonistic effects in the liver and other tissues. Insulin clamp studies have shown that if high doses of GH are given, then insulin sensitivity deteriorates acutely as a result of increased free fatty acid release (102–106). However, low doses given for 6–12 months cause no change in insulin sensitivity (102,103,104,105,106). Because individual patients have differential sensitivity in these parameters, it is not surprising that some show a worsening of insulin sensitivity after administration of GH, whereas others show little change. A recent meta-analysis of placebo-controlled studies showed that GH therapy was associated with a slight rise in both fasting glucose and fasting insulin levels (93). However, there is limited evidence as to whether this effect persists after long-term GH treatment has altered body composition.

Increased intimal-medial thickness (IMT) and abnormal arterial wall dynamics have been documented in GHD (92, 107–109). One study showed that subjects with a low IGF-I had the greatest increase in IMT (110). Three small studies have shown that administration of GH to GH-deficient adults and/or children resulted in decreased IMT (53, 111, 112). In epidemiological studies, increases in IMT have predicted the development of symptomatic coronary disease occurring approximately 8 yr after the initial

measurements (113). This finding suggests that patients with an IMT response may have a significant improvement in cardiovascular outcome, but as yet, this question has not been specifically addressed in patients with GHD.

Cardiac function may also be significantly impaired in GHD. Patients with childhood-onset GHD had reduced left ventricular (LV), posterior wall, and interventricular septal thickness, and LV diameter and mass as evaluated by echocardiography (114, 115). In GHD patients younger than 40 yr, whether their GHD was of adult or childhood onset, there was LV systolic dysfunction at rest and after peak physical exercise as compared with control subjects (116). Analysis of several studies has shown that the most consistent increases after GH administration were in LV mass, LV end diastolic volume, and stroke volumes (117). It is possible that changes in these parameters correlate with the reported subjective benefits in increased exercise tolerance and energy that have been reported by GH-deficient patients after replacement therapy.

Mortality in hypopituitarism

Epidemiological studies have shown that adults with hypopituitarism have increased mortality compared with nonhypopituitary populations adjusted for age and sex. The causes of premature mortality were cardiovascular and cerebrovascular disease. Because of cranial radiation, surgery, and other hormone replacement therapy issues, it cannot be concluded that this premature mortality can be attributed solely to GHD.

Several retrospective epidemiological studies that excluded patients with Cushing's disease and acromegaly have demonstrated premature mortality in patients with pituitary lesions treated with surgery and cranial radiation. All-cause standardized mortality ratios (SMRs) were 1.73–1.87 (118–120). Cardiovascular SMRs ranged between 1.41 and 1.9 (118, 121–123). Cerebrovascular SMRs were higher, ranging between 2.44 and 3.39 (119, 121). The higher SMRs for cerebrovascular mortality were associated with the diagnosis of craniopharyngioma and high rates

of cranial irradiation (119, 121). Untreated gonadal steroid deficiency was also associated with increased risk for premature death in one study (119).

Another contributor to premature mortality is a progression of pituitary disease that requires additional surgery. In a study of 281 patients who underwent surgery and cranial radiation (1946–1988), 35 had a regrowth of the pituitary adenoma that required another operation. Twenty-five of these 35 patients died (cardiovascular SMR, 3.74; cerebrovascular SMR, 3.77, both compared with the general population). In the 246 patients who did not have tumor regrowth, the overall SMR was 1.71 (cardiovascular SMR, 1.56; cerebrovascular SMR, 3.54, both compared with the general population) (123). There is no evidence that GH treatment affects the rate of pituitary tumor regrowth.

Quality of life

Quality of life is usually assessed via self-administered questionnaires that reflect a variety of health-related, economic, and social factors. Quality of life measures may be broadly correlated with, but are different from, assessments of affect or cognition. Disease-specific quality of life assessment questionnaires have been validated and are now widely used (124, 125).

Quality of life evaluations of GHD patients have shown a high degree of variability. For example, in the untreated state, some patients reported severe

impairment in quality of life and some said their quality of life was normal (126). In particular, significant impairment in quality of life was less frequently observed in adults with childhood-onset GHD (98). The area of quality of life most likely to be affected by GHD was energy and vitality (127). Some studies showed definite benefit after patients received GH replacement therapy, but in others improvements were more limited or no improvement was seen (48, 90, 125–130). The degree of improvement in quality of life is generally proportional to the deviation from normality at the outset (128, 129) but shows no correlation with the degree of improvement in IGF-1 levels (125). In practice, this means that if the quality of life of the patients is normal at baseline, no improvement will be seen with GH replacement (130). Some studies have shown that much of the improvement in quality of life occurs within the first 3 months of GH replacement (129, 131). Some long-term studies have shown sustained benefit in some aspects of quality of life among treated patients as compared with untreated patients (132).

Recommendations

- GH therapy of GH-deficient adults offers significant clinical benefits in body composition, exercise capacity, skeletal integrity, and quality of life measures (level of evidence, moderate).
- GH treatment is most likely to benefit those patients who have more severe clinical and biochemical GHD (level of evidence, moderate).

Side Effects and Risks Associated with GH Therapy

Initial studies of GH replacement used high doses and were associated with numerous side effects (47, 133–136). Consequently, dosages were reduced, resulting in fewer adverse effects and a much improved safety profile (137–148).

Most adverse effects are related to the metabolic effects of GH and are dose related. The most common side effects are related to fluid retention, occurring in 5–18% of patients, and include paresthesias, joint stiffness, peripheral edema, arthralgia, and myalgia. Carpal tunnel syndrome occurs in approximately 2% of treated AGHD patients. Adult patients who are older, heavier, or female are more prone to develop these complications (148). Most of these adverse reactions improve with dose reduction. Increased blood pressure is seen when fluid retention occurs, but this problem can be avoided with appropriate dosing (149).

Insulin resistance and type 2 diabetes have been reported in a few patients in the early large clinical trials (104). As noted above, there is considerable variability in changes in insulin sensitivity due to differences in body composition, age, and genetic predisposition. In the placebo-controlled study by Hoffman *et al.* (48), GH therapy was associated with a worsening of glucose tolerance to impaired glucose tolerance in 13% and to diabetes in 4% of patients, the total number with worsening being greater than what was seen with placebo. Thus, with current dosing regimens, there may be a slight excess risk of diabetes mellitus; monitoring of diabetic patients for changes in medication needs is appropriate.

Retinopathy is an extremely rare complication of GH therapy. Two patients without diabetes, one an adult and the other a 9-yr-old patient with Turner's syndrome, developed retinopathy while receiving GH but

improved after its withdrawal (150–152). By contrast, none of 85 children with IGHD who received GH for 6.4 ± 2.9 yr developed retinopathy (153).

Benign intracranial hypertension has been linked to GH treatment in children (154), but only one case has been reported in an adult (155). Gynecomastia has been reported in normal elderly individuals receiving GH in high doses (156, 157). Galactorrhea has not been reported.

There has been a concern that GH therapy and its attendant increase in IGF-I could lead to tumor recurrence or the development of malignancies. However, an increase in the recurrence rates of either intracranial or extracranial tumors has not been demonstrated in AGHD (158,159). Fradkin *et al.* (160) did report an increase in leukemia in children treated with GH, but the excess risk could be attributed to the presence of other tumors and/or radiotherapy. There was no increase in leukemia among pediatric patients with idiopathic GHD who received GH (160). In a second series (161), mortality from colorectal cancer and Hodgkin's disease was increased in a cohort of 1848 GHD patients who received GH during childhood; however, the number of cases was small (only two cases of each), and treatment parameters differed from modern day dosing regimens. No increased rates of leukemia were reported in this cohort. There are no published reports of long-term observational studies in patients with AGHD treated with GH with respect to the development of malignancies.

Although not an adverse effect, a recent study showed that GH replacement caused a lowering of serum free T₄ levels, perhaps due to increased deiodination of T₄ (162). In another study from this group, GH replacement was found to cause a lowering of serum

cortisol levels, bringing out central hypoadrenalism that had been masked, likely due to enhanced conversion of cortisone to cortisol during the GH-deficient state (163). Thus, free T₄ levels should be monitored during GH treatment, and doses of T₄ should be adjusted as necessary (162). Similarly, the hypothalamic-pituitary-adrenal axis should be reassessed in GHD patients during GH therapy, if they had not been previously found to be deficient in this axis, and glucocorticoid replacement should be instituted if necessary.

Recommendation

- GH treatment is contraindicated in the presence of an active malignancy (level of evidence, low).

Suggestion

- GH treatment in patients with diabetes mellitus may require adjustments in antidiabetic medications (level of evidence, moderate).

Treatment Regimens

GH dosing in adults was initially adopted from pediatric practice and was subsequently found to be supraphysiological and to cause high rates of adverse effects. Adults generally are much more susceptible to the side effects of GH than are children, even at doses achieving similar IGF-I responses (13, 128, 131, 133).

Dosing plans have evolved from weight-based dosing to individualized dose-titration strategies. Two studies that formally compared these dosing approaches found that adverse effects were less than half as frequent in the dose-titration group as in the weight-based dosing group, and the final maintenance doses were somewhat lower in the dose-titration group than in the weight-based dosing group (133, 142).

In general, women require higher doses of GH to achieve the same IGF-I response (131, 164). However, when men and women with similar IGF-I responses were matched, the effects of GH on clinical endpoints such as body fat, LDL cholesterol, and circulating markers of bone turnover were still blunted in women (164). Cook et al. (165) reported similar contrasting results for men and women, and they found that much higher GH doses were needed to achieve the same IGF-I levels in women receiving oral estrogen replacement. As women come off estrogen therapy or are switched from oral to transdermal estrogen, GH doses may need to be lowered.

GH secretion normally decreases with age, and older patients have an increased susceptibility to GH-related side effects. Therefore, GH dose requirements should be lower in older patients. By the same token, higher doses may be appropriate in some transition and young adult patients (89). For patients aged 30–60 yr, a starting dose of 300 $\mu\text{g}/\text{d}$ usually will not be associated with side effects. Daily dosing should be increased by 100–200 μg every 1 to 2 months, the goals being an appropriate

clinical response, an avoidance of side effects, and an IGF-I level in the age-adjusted reference range. A commonly used target is the upper half of that range, although no published studies offer specific guidance in this regard. Clinical benefits may not become apparent for 6 or more months of treatment. Older (>60 yr) patients should be started on lower doses (100–200 $\mu\text{g}/\text{d}$) and increased more slowly. Younger (<30 yr) patients may benefit from higher initial doses (400–500 $\mu\text{g}/\text{d}$); for patients transitioning from pediatric treatment, even higher doses may be appropriate. Women who are on oral estrogen replacement usually need substantially higher doses of GH, but those on transdermal estrogen preparations may not (165).

After maintenance doses have been achieved, monitoring usually occurs at 6-month intervals. Such monitoring should include a clinical evaluation, an assessment of side effects, and measurement of IGF-I levels. The lipid profile and a fasting glucose should be assessed annually. If the initial bone DXA scan is abnormal, then repeat evaluations at 1- to 2-yr intervals may be useful in assessing the need for additional treatment modalities. Assessment of quality of life provides another modality for monitoring the response to therapy. Hypopituitary patients on thyroid hormone replacement may need dose adjustments after starting GH replacement, and the hypothalamic-pituitary-adrenal axis should also be reevaluated, as noted above. These recommendations for monitoring are based on clinical experience rather than being validated by large, controlled studies.

The transition from pediatric to adult care raises specific physiological and logistical issues. As discussed above, peak bone mass has generally not been achieved at the time final height is attained. Stopping GH at that point has also been associated with a worsening of lipoprotein

profiles, body composition, and quality of life scores (89,99). Consideration should therefore be given to minimizing lengthy interruptions in GH therapy. However, it is important to retest such individuals, especially those who had IGHD.

How long to administer GH therapy is unclear. If benefits are being achieved, there is no particular reason to stop treatment. On the other hand, if there are no apparent or objective benefits of treatment after at least 1 yr of treatment, discontinuing GH therapy may be appropriate.

Recommendations

- GH dosing regimens should be individualized rather than weight-based (level of evidence, high).
- GH treatment should start with low doses and be titrated according to clinical response, side effects, and IGF-I levels (level of evidence, high).
- GH dosing should take age, sex, and estrogen status into consideration (level of evidence, high).
- During GH treatment, patients should be monitored at 1- to 2-month intervals during dose titration and semiannually thereafter with a clinical assessment and an evaluation for adverse effects, IGF-I levels, and other parameters of GH response (level of evidence, moderate).

Conclusion, Summary of Recommendations & Suggestions

Conclusion

GH therapy has been shown to benefit many adults with GHD. It is critical to identify appropriate candidates in whom the clinical context suggests that GHD may be present. Confirmation of GHD before beginning therapy is crucial and usually involves biochemical testing. The demonstrated benefits of GH therapy include improvements in body composition, exercise capacity, skeletal integrity, lipids, and quality of life. Although it has been suggested that GH treatment may reduce the increased vascular mortality associated with hypopituitarism, this has not yet been proved. It should be emphasized that long-term clinical outcome studies on hard endpoints such as fractures, clinical heart disease, cancer, and mortality are still lacking at present. Dosing should be individualized with attention to avoidance of side effects. Periodic monitoring will be necessary for adverse effects and physiological benefit.

Recommendations

- Patients with childhood-onset GHD who are appropriate candidates for GH therapy should be retested for GHD as adults unless they have known mutations, embryopathic lesions, or irreversible structural lesions/damage (level of evidence, high).
- Adult patients with evidence of structural hypothalamic/pituitary disease, surgery or irradiation to these areas, or other pituitary hormone deficiencies should be considered for evaluation for acquired GHD (level of evidence, high).
- The ITT or the GHRH-arginine test is the preferred test for establishing the diagnosis of GHD. However, in those with clearly established recent hypothalamic causes of suspected GHD, *e.g.* irradiation, testing with GHRH-arginine may be

misleading (level of evidence, high).

- Because of the irreversible nature of the cause of the GHD in children with structural lesions with multiple hormone deficiencies and those with proven genetic causes, a low IGF-I level at least 1 month off GH therapy is sufficient documentation of persistent GHD without additional provocative testing (level of evidence, moderate).
- GH therapy of GH-deficient adults offers significant clinical benefits in body composition, exercise capacity, skeletal integrity, and quality of life measures (level of evidence, moderate).
- GH treatment is most likely to benefit those patients who have more severe clinical and biochemical abnormalities and should be encouraged in such patients (level of evidence, moderate).
- GH treatment is contraindicated in the presence of an active malignancy (level of evidence, low).
- GH dosing regimens should be individualized rather than weight-based (level of evidence, high).
- GH treatment should start with low doses and be titrated according to clinical response, side effects, and IGF-I levels (level of evidence, high).
- GH dosing should take age, sex, and estrogen status into consideration (level of evidence, high).
- During GH treatment, patients should be monitored at 1- to 2-month intervals during dose titration and semiannually thereafter with a clinical assessment and an evaluation for adverse effects, IGF-I levels, and other parameters of GH response (level of evidence, moderate).

Suggestions

- A normal IGF-I level does not exclude the diagnosis of GHD and, in the context of other pituitary disease, makes provocative testing mandatory to make the diagnosis of GHD (level of evidence, high).
- A low IGF-I level, in the absence of catabolic conditions and liver disease, indicates severe GHD and may be useful in identifying patients who will

benefit from treatment (level of evidence, moderate).

- The presence of deficiencies in three or more pituitary axes strongly suggests the presence of GHD, and in this context provocative testing is optional (level of evidence, moderate).
- GH treatment in patients with diabetes mellitus may require adjustments in antidiabetic medications (level of evidence, moderate).

Appendix

Overall grades of evidence

High	Further research is very unlikely to change our confidence in the estimate of effect.
Moderate	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
Low	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.
Very Low	Any estimate of effect is very uncertain.

Above based on criteria for assigning grade of evidence

Types of evidence

High	Randomized trial
Low	Observational study (cohort studies, case-control studies, interrupted time series analyses, controlled before and after studies)
Very low	Any other evidence

Decrease grade if:

- Serious (–1) or very serious (–2) limitation to study quality
- Important inconsistency (–1)
- Some (–1) or major (–2) uncertainty about directness
- Imprecise or sparse data (–1)
- High probability of reporting bias (–1)

Increase grade if:

- Strong evidence of association-significant relative risk of >2 based on consistent evidence from two or more observational studies with no plausible confounders (+1)
- Very strong evidence of association-significant relative risk of >5 based on direct evidence with no major threats to validity (+2)
- Evidence of a dose-response gradient
- All plausible confounders would have reduced the effect (+1)

Recommendation: ‘Do it’ or ‘Don’t do it’—indicating a judgment that most well-informed people would make

Suggestion: ‘Probably do it’ or ‘Probably don’t do it’—indicating a judgment that a majority of well-informed people would make but a substantial minority would not

References

1. **Raben MS** 1958 Treatment of a pituitary dwarf with human growth hormone. *J Clin Endocrinol* 18:901-903
2. **MacGillivray MH, Blizzard RM** 1994 Rationale for dosing recombinant human growth hormone by weight rather than units. *Growth Genet Horm* 10:7-9
3. **Growth Hormone Research Society** 2001 Critical evaluation of the safety of recombinant human growth hormone administration: statement from the Growth Hormone Research Society. *J Clin Endocrinol Metab* 86:1868-1870
4. **Cohen LE, Radovick S** 2002 Molecular basis of combined pituitary hormone deficiencies. *Endocr Rev* 23:431-442
5. **Dattani MT** 2005 Growth hormone deficiency and combined pituitary hormone deficiency: does the genotype matter? *Clin Endocrinol (Oxf)* 63:121-130
6. **Moseley CT, Phillips 3rd JA** 2000 Pituitary gene mutations and the growth hormone pathway. *Semin Reprod Med* 18:21-29
7. **Wajnrajch MP, Gertner JM, Harbison MD, Chua Jr SC, Leibel RL** 1996 Nonsense mutation in the human growth hormone-releasing hormone receptor causes growth failure analogous to the little (lit) mouse. *Nat Genet* 12:88-90
8. **Mantovani G, Maghnie M, Weber G, DeMenis E, Brunelli V, Cappa M, Loli P, Beck-Peccoz P, Spada A** 2003 Growth hormone-releasing hormone resistance in pseudohypoparathyroidism type ia: new evidence for imprinting of the Gs a gene. *J Clin Endocrinol Metab* 88:4070-4074
9. **Hintz RL, Menking M, Sotos JF** 1968 Familial holoprosencephaly with endocrine dysgenesis. *J Pediatr* 72:81-87
10. **Zuppinger KA, Sutter M, Zurbrugg RP, Joss EE, Oetliker O** 1971 Cleft lip and choroidal coloboma associated with multiple hypothalamo-pituitary dysfunctions. *J Clin Endocrinol Metab* 33:934-939
11. **Stelling M, Goldstein D, Johanson A, Blizzard R** 1980 Hypopituitarism and dysmorphism: problems in paediatric endocrinology. In: La Caupa C, Root A, eds. *Proceedings of Sero Symposium*. New York: Academic Press; 101-112
12. **Bierich JR, Christie M, Heinrich JJ, Martinez AS** 1991 New observations on midline defects: co-incidence of anophthalmos, microphthalmos and cryptophthalmos with hypothalamic disorders. *Eur J Pediatr* 150:246-249
13. **Cacciari E, Zucchini S, Ambrosetto P, Tani G, Carla G, Cicognani A, Pirazzoli P, Sganga T, Balsamo A, Cassio A** 1994 Empty sella in children and adolescents with possible hypothalamic-pituitary disorders. *J Clin Endocrinol Metab* 78:767-771
14. **Gleeson HK, Gattamaneni HR, Smethurst L, Brennan BM, Shalet SM** 2004 Reassessment of growth hormone status is required at final height in children treated with growth hormone replacement after radiation therapy. *J Clin Endocrinol Metab* 89:662-666
15. **de Boer H, Bolk GJ, Van der Veen EA** 1995 Clinical aspects of growth hormone deficiency in adults. *Endocr Rev* 16:63-86
16. **Clayton PE, Price DA, Shalet SM** 1987 Growth hormone state after completion of treatment with growth hormone. *Arch Dis Child* 62:222-226
17. **Cacciari E, Tassoni P, Parisi G, Pirazoli P, Zucchini S, Mandini M, Cicognani A, Balsamo A** 1992 Pitfalls in diagnosing impaired growth hormone (GH) secretion: retesting after replacement therapy of 63 patients defined as GH deficient. *J Clin Endocrinol Metab* 74:1284-1289
18. **Wacharasindhu S, Cotterill AM, Camacho-Hubner C, Besser GM, Savage MO** 1996 Normal growth hormone secretion in growth hormone insufficient children re-tested after completion of linear growth. *Clin Endocrinol (Oxf)* 45:553-556
19. **Longobardi S, Merola B, Pivonello R, Di Rella F, Di Somma C, Colao A, Ghigo E, Camanni F, Lombardi G** 1996 Re-evaluation of growth hormone (GH) secretion in 69 adults diagnosed as GH-deficient patients during childhood. *J Clin Endocrinol Metab* 81:1244-1247
20. **Nicolson A, Toogood AA, Rahim A, Shalet SM** 1996 The prevalence of severe growth hormone deficiency in adults who received growth hormone replacement in childhood. *Clin Endocrinol (Oxf)* 44:311-316
21. **Tauber M, Moulin P, Pienkowski C, Jouret B, Rochiccioli P** 1997 Growth hormone (GH) retesting and auxological data in 131 GH-deficient patients after completion of treatment. *J Clin Endocrinol Metab* 82:352-356
22. **Arafah BM, Prunty D, Ybarra J, Hlavin ML, Selman WR** 2000 The dominant role of increased intrasellar pressure in the pathogenesis of hypopituitarism, hyperprolactinemia, and headaches in patients with pituitary adenomas. *J Clin Endocrinol Metab* 85:1789-1793
23. **Webb SM, Rigla M, Wagner A, Oliver B, Bartumeus F** 1999 Recovery of hypopituitarism after neurosurgical treatment of pituitary adenomas. *J Clin Endocrinol Metab* 84:3696-3700

24. **Arafah BM, Kailani SH, Nekl KE, Gold RS, Selman WR** 1994 Immediate recovery of pituitary function after transsphenoidal resection of pituitary macroadenomas. *J Clin Endocrinol Metab* 79:348-354
25. **Arafah BM** 1986 Reversible hypopituitarism in patients with large non-functioning pituitary adenomas. *J Clin Endocrinol Metab* 62:1173-1179
26. **Littley MD, Shalet SM, Beardwell CG, Ahmed SR, Applegate G, Sutton ML** 1989 Hypopituitarism following external radiotherapy for pituitary tumours in adults. *Q J Med* 70:145-160
27. **Snyder PJ, Fowble BF, Schatz NJ, Savino PJ, Gennarelli TA** 1986 Hypopituitarism following radiation therapy of pituitary adenomas. *Am J Med* 81:457-462
28. **Feigl GC, Bonelli CM, Berghold A, Mokry M** 2002 Effects of gamma knife radiosurgery of pituitary adenomas on pituitary function. *J Neurosurg* 97(5 Suppl):415-421
29. **Agha A, Rogers B, Sherlock M, O'Kelly P, Tormey W, Phillips J, Thompson CJ** 2004 Anterior pituitary dysfunction in survivors of traumatic brain injury. *J Clin Endocrinol Metab* 89:4929-4936
30. **Leal-Cerro A, Flores JM, Rincon M, Murillo F, Pujol M, Garcia-Pesquera F, Dieguez C, Casanueva FF** 2005 Prevalence of hypopituitarism and growth hormone deficiency in adults long-term after severe traumatic brain injury. *Clin Endocrinol (Oxf)* 62:525-532
31. **Aimaretti G, Ambrosio MR, Di Somma C, Fusco A, Cannavo S, Gasperi M, Scaroni C, DeMarinis L, Benvenega S, degli Uberti EC, Lombardi G, Mantero F, Martino E, Giordano G, Ghigo E** 2004 Traumatic brain injury and subarachnoid haemorrhage are conditions at high risk for hypopituitarism: screening study at 3 months after the brain injury. *Clin Endocrinol (Oxf)* 61:320-326
32. **Aimaretti G, Ambrosio MR, Di Somma C, Gasperi M, Cannavo S, Scaroni C, Fusco A, Del Monte P, De Menis E, Faustini-Fustini M, Grimaldi F, Logoluso F, Razzore P, Rovere S, Benvenega S, degli Uberti EC, De Marinis L, Lombardi G, Mantero F, Martino E, Giordano G, Ghigo E** 2005 Residual pituitary function after brain injury-induced hypopituitarism: a prospective 12-month study. *J Clin Endocrinol Metab* 90:6085-6092
33. **Vahl N, Jorgensen JO, Jurik AG, Christiansen JS** 1996 Abdominal adiposity and physical fitness are major determinants of the age associated decline in stimulated GH secretion in healthy adults. *J Clin Endocrinol Metab* 81:2209-2215
34. **Bonert VS, Elashoff JD, Barnett P, Melmed S** 2004 Body mass index determines evoked growth hormone (GH) responsiveness in normal healthy male subjects: diagnostic caveat for adult GH deficiency. *J Clin Endocrinol Metab* 89:3397-3401
35. **Aimaretti G, Corneli G, Razzore P, Bellone S, Baffoni C, Arvat E, Camanni F, Ghigo E** 1998 Comparison between insulin-induced hypoglycemia and growth (GH)-releasing hormone + arginine as provocative tests for the diagnosis of GH deficiency in adults. *J Clin Endocrinol Metab* 83:1615-1618
36. **Biller BM, Samuels MH, Zagar A, Cook DM, Arafah BM, Bonert V, Stavrou S, Kleinberg DL, Chipman JJ, Hartman ML** 2002 Sensitivity and specificity of six tests for the diagnosis of adult growth hormone deficiency. *J Clin Endocrinol Metab* 87:2067-2079
37. **Darzy KH, Aimaretti G, Wieringa G, Gattamaneni HR, Ghigo E, Shalet SM** 2003 The usefulness of the combined growth hormone (GH)-releasing hormone and arginine stimulation test in the diagnosis of radiation-induced GH deficiency is dependent on the post-irradiation time interval. *J Clin Endocrinol Metab* 88:95-102
38. **Hoffman DM, O'Sullivan AJ, Baxter RC, Ho KK** 1994 Diagnosis of growth-hormone deficiency in adults. *Lancet* 343:1064-1068
39. **Hartman ML, Crowe BJ, Biller BM, Ho KK, Clemmons DR, Chipman JJ; HyposCCS Advisory Board; U.S. HypoCCS Study Group** 2002 Which patients do not require a GH stimulation test for the diagnosis of adult GH deficiency? *J Clin Endocrinol Metab* 87:477-485
40. **Hilding A, Hall K, Wivall-Helleryd IL, Saaf M, Melin AL, Thoren M** 1999 Serum levels of insulin-like growth factor I in 152 patients with growth hormone deficiency, aged 19-82 years, in relation to those in healthy subjects. *J Clin Endocrinol Metab* 84:2013-2019
41. **Juul A, Kastrup KW, Pedersen SA, Skakkebaek NE** 1997 Growth hormone (GH) provocative retesting of 108 young adults with childhood-onset GH deficiency and the diagnostic value of insulin-like growth factor I (IGF-I) and IGF-binding protein-3. *J Clin Endocrinol Metab* 82:1195-1201
42. **Toogood AA, Beardwell CG, Shalet SM** 1994 The severity of growth hormone deficiency in adults with pituitary disease is related to the degree of hypopituitarism. *Clin Endocrinol (Oxf)* 41:511-516
43. **Growth Hormone Research Society** 1998 Consensus guidelines for the diagnosis and treatment of adults with growth hormone deficiency: summary statement of the Growth Hormone Research Society Workshop on Adult Growth Hormone Deficiency. *J Clin Endocrinol Metab* 83:379-381
44. **Hoffman DM, O'Sullivan AJ, Freund J, Ho KK** 1995 Adults with growth hormone deficiency have abnormal

- body composition but normal energy metabolism. *J Clin Endocrinol Metab* 80:72-77
45. **Beshyah SA, Freemantle C, Thomas E, Rutherford O, Page B, Murphy M, Johnston DG** 1995 Abnormal body composition and reduced bone mass in growth hormone deficient hypopituitary adults. *Clin Endocrinol (Oxf)* 42:179-189
 46. **Johannsson G, Marin P, Lonn L, Ottosson M, Stenlof K, Bjorntorp P, Sjostrom L, Bengtsson BA** 1997 Growth hormone treatment of abdominally obese men reduces abdominal fat mass, improves glucose and lipoprotein metabolism, and reduces diastolic blood pressure. *J Clin Endocrinol Metab* 82:727-734
 47. **Bengtsson BA, Eden S, Lonn L, Kvist H, Stokland A, Lindstedt G, Bosaeus I, Tolli J, Sjostrom L, Isaksson OG** 1993 Treatment of adults with growth hormone (GH) deficiency with recombinant human GH. *J Clin Endocrinol Metab* 76:309-317
 48. **Hoffman AR, Kuntze JE, Baptista J, Baum HB, Baumann GP, Biller BM, Clark RV, Cook D, Inzucchi SE, Kleinberg D, Klibanski A, Phillips LS, Ridgway EC, Robbins RJ, Schlechte J, Sharma M, Thorner MO, Vance ML** 2004 Growth hormone (GH) replacement therapy in adult-onset GH deficiency: effects on body composition in men and women in a double-blind, randomized, placebo-controlled trial. *J Clin Endocrinol Metab* 89:2048-2056
 49. **Chrisoulidou A, Beshyah SA, Rutherford O, Spinks TJ, Mayet J, Kyd P, Anyaoku V, Haida A, Ariff B, Murphy M, Thomas E, Robinson S, Foale R, Johnston DG** 2000 Effects of 7 years of growth hormone replacement therapy in hypopituitary adults. *J Clin Endocrinol Metab* 85:3762-3769
 50. **Attanasio AF, Bates PC, Ho KK, Webb SM, Ross RJ, Strasburger CJ, Bouillon R, Crowe B, Selander K, Valle D, Lamberts SW** 2002 Hypopituitary Control and Complications Study International Advisory Board. Human growth hormone replacement in adult hypopituitary patients: long-term effects on body composition and lipid status--3-year results from the HypoCCS Database. *J Clin Endocrinol Metab* 87:1600-1606
 51. **Salomon F, Cuneo RC, Hesp R, Sonksen P** 1989 The effects of treatment with recombinant human growth hormone on body composition and metabolism in adults with growth hormone deficiency. *N Engl J Med* 321:1797-1803
 52. **Al-Shoumer K, Page B, Thomas E, Murphy M, Beshyah SA, Johnston DG** 1996 Effects of four years' treatment with biosynthetic human growth hormone (GH) on body composition in GH-deficient hypopituitary adults. *Eur J Endocrinol* 135:559-567
 53. **Gibney J, Wallace JD, Spinks T, Schnorr L, Ranicar A, Cuneo RC, Lockhart S, Burnand KG, Salomon F, Sonksen PH, Russell-Jones D** 1999 The effects of 10 years of recombinant human growth hormone in adult GH deficient patients. *J Clin Endocrinol Metab* 84:2596-2602
 54. **Attanasio A, Howell S, Bates P, Frewer P, Chipman J, Blum WF, Shalet SM** 2002 Body composition, IGF-I and IGFBP-3 concentrations as outcome measures in severely GH-deficient GHD patients. *J Clin Endocrinol Metab* 87:3368-3372
 55. **Svensson J, Stibrant Sunnerhagen K, Johannsson G** 2003 Five years of growth hormone replacement therapy in adults: age and gender related changes in isometric and isokinetic muscle strength. *J Clin Endocrinol Metab* 88:2061-2069
 56. **Johannsson G, Grimby G, Sunnerhagen K, Bengtsson B** 1997 Two years of growth hormone (GH) treatment increase isometric and isokinetic muscle strength in GH-deficient adults. *J Clin Endocrinol Metab* 82:2877-2884
 57. **Cuneo RC, Salomon F, Wiles CM, Hesp R, Sonksen PH** 1991 Growth hormone treatment in growth hormone-deficient adults. I. Effects on muscle mass and strength. *J Appl Physiol* 70:688-694
 58. **Janssen YJ, Doornbos J, Roelfsema F** 1999 Changes in muscle volume, strength, and bioenergetics during recombinant human growth hormone (GH) therapy in adults with GH deficiency. *J Clin Endocrinol Metab* 84:279-284
 59. **Woodhouse LJ, Asa SL, Thomas SG, Ezzat S** 1999 Measures of submaximal aerobic performance evaluate and predict functional response to growth hormone (GH) treatment in GH-deficient adults. *J Clin Endocrinol Metab* 84:4570-4577
 60. **Cuneo RC, Salomon F, Wiles CM, Hesp R, Sonksen PH** 1991 Growth hormone treatment in growth hormone deficient adults. II. Effects on exercise performance. *J Appl Physiol* 70:695-700
 61. **Elgzyri T, Castenfors J, Hagg E, Backman C, Thoren M, Brammert M** 2004 The effects of GH replacement therapy on cardiac morphology and function, exercise capacity and serum lipids in elderly patients with GH deficiency. *Clin Endocrinol (Oxf)* 61:113-122
 62. **Jorgensen J, Thuesen L, Muller J, Ovesen P, Skakkebaek N, Christiansen J** 1994 Three years of growth hormone treatment in growth hormone deficient adults: near normalization of body composition and physical performance. *Eur J Endocrinol* 130:224-228
 63. **Holmes SJ, Economou G, Whitehouse RW, Adams JE, Shalet SM** 1994 Reduced bone mineral density in patients with adult onset growth hormone deficiency. *J Clin Endocrinol Metab* 78:669-674

64. **Murray RD, Columb B, Adams JE, Shalet SM** 2004 Low bone mass is an infrequent feature of the adult growth hormone deficiency syndrome in middle-age adults and the elderly. *J Clin Endocrinol Metab* 89:1124-1130
65. **Rosen T, Hansson T, Granhed H, Szucs J, Bengtsson BA** 1993 Reduced bone mineral content in adult patients with growth hormone deficiency. *Acta Endocrinol (Copenh)* 129:201-206
66. **Kaufman J, Taelman P, Vermeulen A, Vandeweghe M** 1992 Bone mineral status in growth hormone-deficient males with isolated and multiple pituitary insufficiencies of childhood onset. *J Clin Endocrinol Metab* 74:118-123
67. **Lissett CA, Murray RD, Shalet SM** 2002 Timing of onset of growth hormone deficiency is a major influence of insulin-like growth factor I status in adult life. *Clin Endocrinol (Oxf)* 57:35-40
68. **Colao A, Di Somma C, Pivonello R, Loche S, Aimaretti G, Cerbone G, Faggiano A, Corneli G, Ghigo E, Lombardi G** 1999 Bone loss is correlated to the severity of growth hormone deficiency in adult patients with hypopituitarism. *J Clin Endocrinol Metab* 84:1919-1924
69. **Hyer S, Rodin D, Tobias J, Leiper A, Nussey S** 1992 Growth hormone deficiency during puberty reduces adult bone mineral density. *Arch Dis Childhood* 67:1472-1474
70. **Bravenboer N, Holzmann P, de Boer H, Blok GJ, Lips P** 1996 Histomorphometric analysis of bone mass and bone metabolism in growth hormone deficient adult men. *Bone* 18:551-557
71. **Rosen T, Wilhelmsen L, Landin-Wilhelmsen K, Lappas G, Bengtsson BA** 1997 Increased fracture frequency in adult patients with hypopituitarism and GH deficiency. *Eur J Endocrinol* 137:240-245
72. **Wuster C, Abs R, Bengtsson BA, Bennmarker H, Feldt-Rasmussen U, Hernberg-Stahl E, Monson JP, Westberg B, Wilton P; KIMS Study Group and the KIMS International Board** 2001 The influence of growth hormone deficiency, growth hormone replacement therapy, and other aspects of hypopituitarism on fracture rate and bone mineral density. *J Bone Miner Res* 16:398-405
73. **Vestergaard P, Jorgensen JO, Hagen C, Hoeck HC, Laurberg P, Rejnmark L, Brixen K, Weeke J, Andersen M, Conceicao FL, Nielsen TL, Mosekilde L** 2002 Fracture risk is increased in patients with GH deficiency or untreated prolactinomas—a case-control study. *Clin Endocrinol (Oxf)* 56:159-167
74. **Baum HB, Biller BM, Finkelstein JS, Cannistraro KB, Oppenheim DS, Schoenfeld DA, Michel TH, Wittink H, Klibanski A** 1996 Effects of physiologic growth hormone therapy on bone density and body composition in patients with adult-onset growth hormone deficiency. A randomized, placebo-controlled trial. *Ann Intern Med* 125:883-890
75. **Hansen T, Brixen K, Vahl N, Jorgensen JO, Christiansen JS, Mosekilde L, Hagen C** 1996 Effects of 12 months of growth hormone (GH) treatment on calcitropic hormones, calcium homeostasis, and bone metabolism in adults with acquired GH deficiency: a double-blind, randomized, placebo-controlled study. *J Clin Endocrinol Metab* 81:3352-3359
76. **Holmes SJ, Whitehouse RW, Swindell R, Economou G, Adams JE, Shalet SM** 1995 Effect of growth hormone replacement on bone mass in adults with adult-onset growth hormone deficiency. *Clin Endocrinol (Oxf)* 42:627-633
77. **Biller BM, Sesimo G, Baum HB, Hayden D, Schoenfeld D, Klibanski A** 2000 Withdrawal of long-term physiological growth hormone (GH) administration: differential effects on bone density and body composition in men with adult-onset GH deficiency. *J Clin Endocrinol Metab* 85:970-976
78. **Shalet SM, Shavrikova E, Cromer M, Child CJ, Keller E, Zapletalova J, Moshang T, Blum WF, Chipman JJ, Quigley CA, Attanasio AF** 2003 Effect of growth hormone (GH) treatment on bone in postpubertal GH-deficient patients: a 2-year randomized, controlled dose-ranging study. *J Clin Endocrinol Metab* 88:4124-4129
79. **Johannsson G, Rosen T, Bosaeus I, Sjostrom L, Bengtsson BA** 1996 Two years of growth hormone (GH) treatment increases bone mineral content and density in hypopituitary patients with adult onset GH deficiency. *J Clin Endocrinol Metab* 81:2865-2873
80. **Drake WM, Rodriguez-Arnao J, Weaver JU, James IT, Coyte D, Spector TD, Besser GM, Monson JP** 2001 The influence of gender on the short- and long-term effects of growth hormone replacement on bone metabolism and bone mineral density in hypopituitary adults: a 5-year study. *Clin Endocrinol (Oxf)* 54:525-532
81. **Bex M, Abs R, Maiter D, Beckers A, Lamberigts G, Bouillon R** 2002 The effects of growth hormone replacement therapy on bone metabolism in adult-onset growth hormone deficiency: a 2-year open randomized controlled multicenter trial. *J Bone Miner Res* 17:1081-1094
82. **Biermasz NR, Hamdy NA, Pereira AM, Romijn JA, Roelfsema F** 2004 Long-term skeletal effects of recombinant human growth hormone (rhGH) alone and rhGH combined with alendronate in GH-deficient adults: a seven-year follow-up study. *Clin Endocrinol (Oxf)* 60:568-575
83. **Olney RC** 2003 Regulation of bone mass by growth hormone. *Med Pediatr Oncol* 41:228-234
84. **Teegarden D, Proulx WR, Martin BR, Zhao J, McCabe GP, Lyle RM, Peacock M, Slemenda C, Johnston CC, Weaver CM** 1995 Peak bone mass in young women. *J Bone Miner Res* 10:711-715

85. **Baroncelli GI, Bertelloni S, Sodini F, Saggese G** 2003 Acquisition of bone mass in normal individuals and in patients with growth hormone deficiency. *J Pediatr Endocrinol Metab* 16(Suppl 2):327-335
86. **Baroncelli GI, Bertelloni S, Sodini F, Saggese G** 2004 Longitudinal changes in lumbar bone mineral density (BMD) in patients with GH deficiency after discontinuation of treatment at final height; timing and peak values for lumbar BMD. *Clin Endocrinol (Oxf)* 60:175-184
87. **Mukherjee A, Attanasio A, Shalet SM** 2003 Skeletal requirements for optimal growth hormone replacement in the transitional years. *Growth Horm IGF Res* 13(Suppl A):S130-135
88. **Drake WM, Carroll PV, Maher KT, Metcalfe K, Shaw NJ, Dunger DB, Cheetham TD, Savage MO, Monson JP** 2003 The effect of cessation of growth hormone (GH) therapy on bone mineral accretion in GH-deficient adolescents at the completion of linear growth. *J Clin Endocrinol Metab* 88:1658-1663
89. **Underwood LE, Attie KM, Baptista J; Genentech Collaborative Study Group** 2003 Growth hormone (GH) dose-response in young adults with childhood-onset GH deficiency: a two-year, multicenter, multiple-dose, placebo-controlled study. *J Clin Endocrinol Metab* 88:5273-5280
90. **Mauras N, Pescovitz OH, Allada V, Messig M, Wajnrajch MP, Lippe B; Transition Study Group** 2005 Limited efficacy of growth hormone (GH) during transition of GH-deficient patients from adolescence to adulthood: a phase III multicenter, double-blind, randomized two-year trial. *J Clin Endocrinol Metab* 90:3946-3955
91. **Boger RH, Skamira C, Bode-Boger SM, Brabant G, von zur Muhlen A, Frolich JC** 1996 Nitric oxide may mediate the hemodynamic effects of recombinant growth hormone in patients with acquired growth hormone deficiency. A double-blind, placebo-controlled study. *J Clin Invest* 98:2706-2713
92. **Smith JC, Evans LM, Wilkinson I, Goodfellow J, Cockcroft JR, Scanlon MF, Davies JS** 2002 Effects of GH replacement on endothelial function and large-artery stiffness in GH-deficient adults: a randomized, double-blind, placebo-controlled study. *Clin Endocrinol (Oxf)* 56:493-501
93. **Maison P, Griffin S, Nicoue-Beglah M, Haddad N, Balkau B, Chanson P** 2004 Impact of growth hormone (GH) treatment on cardiovascular risk factors in GH-deficient adults: a meta-analysis of blinded, randomized, placebo-controlled trials. *J Clin Endocrinol Metab* 89:2192-2199
94. **Sesnilo G, Biller BM, Llevadot J, Hayden D, Hanson G, Rifai N, Klibanski A** 2000 Effects of growth hormone administration on inflammatory and other cardiovascular risk markers in men with growth hormone deficiency. A randomized, controlled clinical trial. *Ann Intern Med* 133:111-122
95. **Sesnilo G, Miller K, Hayden D, Klibanski A** 2001 Inflammatory cardiovascular risk markers in women with hypopituitarism. *J Clin Endocrinol Metab* 86:5774-5781
96. **Sesnilo G, Biller BM, Llevadot J, Hayden D, Hanson G, Rifai N, Klibanski A** 2001 Effects of growth hormone (GH) administration on homocyst(e)ine levels in men with GH deficiency: a randomized controlled trial. *J Clin Endocrinol Metab* 86:1518-1524
97. **Bengtsson BA, Abs R, Bennmarker H, Monson JP, Feldt-Rasmussen U, Hernberg-Stahl E, Westberg B, Wilton P, Wuster C** 1999 The effects of treatment and the individual responsiveness to growth hormone (GH) replacement therapy in 665 GH-deficient adults. KIMS Study Group and the KIMS International Board. *J Clin Endocrinol Metab* 84:3929-3935
98. **Attanasio AF, Lamberts SW, Matranga AM, Birkett MA, Bates PC, Valk NK, Hilsted J, Bengtsson BA, Strasburger CJ** 1997 Adult growth hormone (GH)-deficient patients demonstrate heterogeneity between childhood onset and adult onset before and during human GH treatment. Adult Growth Hormone Deficiency Study Group. *J Clin Endocrinol Metab* 82:82-88
99. **Colao A, Di Somma C, Salerno M, Spinelli L, Orio F, Lombardi G** 2002 The cardiovascular risk of GH-deficient adolescents. *J Clin Endocrinol Metab* 87:3650-3655
100. **Kearney T, de Gallegos CN, Proudler A, Parker K, Anyaoku V, Bannister P, Venkatesan S, Johnston DG** 2003 Effects of short- and long-term growth hormone replacement on lipoprotein composition and on very-low-density lipoprotein and low-density lipoprotein apolipoprotein B100 kinetics in growth hormone-deficient hypopituitary subjects. *Metabolism* 52:50-59
101. **Clemmons DR** 2004 The relative roles of growth hormone and IGF-1 in controlling insulin sensitivity. *J Clin Invest* 113:25-27
102. **al-Shoumer KA, Gray R, Anyaoku V, Hughes C, Beshyah S, Richmond W, Johnston DG** 1998 Effects of four year's treatment with biosynthetic human growth hormone (GH) on glucose homeostasis, insulin secretion and lipid metabolism in GH-deficient adults. *Clin Endocrinol (Oxf)* 48:795-802
103. **Weaver JU, Monson JP, Noonan K, John WG, Edwards A, Evans KA, Cunningham J** 1995 The effect of low dose recombinant human growth hormone replacement on regional fat distribution, insulin sensitivity, and cardiovascular risk factors in hypopituitary adults. *J Clin Endocrinol Metab* 80:153-159

104. Svensson J, Fowelin J, Landin K, Bengtsson BA, Johansson JO 2002 Effects of seven years of GH-replacement therapy on insulin sensitivity in GH-deficient adults. *J Clin Endocrinol Metab* 87:2121-2127
105. Yuen K, Cook D, Ong K, Chatelain P, Fryklund L, Gluckman P, Ranke MB, Rosenfeld R, Dunger D 2002 The metabolic effects of short-term administration of physiological versus high doses of GH therapy in GH deficient adults. *Clin Endocrinol (Oxf)* 57:333-341
106. Segerlantz M, Brammert M, Manhem P, Laurila E, Groop LC 2003 Inhibition of lipolysis during acute GH exposure increase insulin sensitivity in previously untreated GH-deficient adults. *Eur J Endocrinol* 149:511-519
107. Leonsson M, Hulthe J, Oscarsson J, Johannsson G, Wendelhag I, Wikstrand J, Bengtsson BA 2002 Intima-media thickness in cardiovascularly asymptomatic hypopituitary adults with growth hormone deficiency: relation to body mass index, gender, and other cardiovascular risk factors. *Clin Endocrinol (Oxf)* 57:751-759
108. Murata M, Kaji H, Mizuno I, Sakurai T, Iida K, Okimura Y, Chihara K 2003 A study of carotid intima-media thickness in GH-deficient adults during onset among adults and children. *Eur J Endocrinol* 148:333-338
109. Markussis V, Beshyah SA, Fisher C, Parker KH, Nicolaides AN, Johnston DG 1997 Abnormal carotid arterial wall dynamics in symptom-free hypopituitary adults. *Eur J Endocrinol* 136:157-164
110. Colao A, Di Somma C, Filippella M, Rota F, Pivonello R, Orio F, Vitale G, Lombardi G 2004 Insulin-like growth factor-1 deficiency determines increased intima-media thickness at common carotid arteries in adult patients with growth hormone deficiency. *Clin Endocrinol (Oxf)* 61:360-366
111. Pfeifer M, Verhovec R, Zizek B, Prezelj J, Poredos P, Clayton RN 1999 Growth hormone (GH) treatment reverses early atherosclerotic changes in GH-deficient adults. *J Clin Endocrinol Metab* 84:453-457
112. Borson-Chazot F, Serusclat A, Kalfallah Y, Ducottet X, Sassolas G, Bernard S, Labrousse F, Pastene J, Sassolas A, Roux Y, Berthezene F 1999 Decrease in carotid intima-media thickness after one year growth hormone (GH) treatment in adults with GH deficiency. *J Clin Endocrinol Metab* 84:1329-1333
113. Hodis HN, Mack WJ, LaBree L, Selzer RH, Liu CR, Liu CH, Azen SP 1998 The role of carotid arterial intima-media thickness in predicting clinical coronary events. *Ann Intern Med* 128:262-269
114. Jallad RS, Liberman B, Vianna CB, Vieira ML, Ramires JA, Knoepfelmacher M 2003 Effects of growth hormone replacement therapy on metabolic and cardiac parameters, in adult patients with childhood-onset growth hormone deficiency. *Growth Horm IGF Res* 13:81-88
115. Colao A, di Somma C, Cuocolo A, Spinelli L, Tedesco N, Pivonello R, Bonaduce D, Salvatore M, Lombardi G 2001 Improved cardiovascular risk factors and cardiac performance after 12 months of growth hormone (GH) replacement in young adult patients with GH deficiency. *J Clin Endocrinol Metab* 86:1874-1881
116. Colao A, Vitale G, Pivonello R, Ciccarelli A, Di Somma C, Lombardi G 2004 The heart: an end-organ of GH action. *Eur J Endocrinol* 151(Suppl 1):S93-S101
117. Maison P, Chanson P 2003 Cardiac effects of growth hormone in adults with growth hormone deficiency: a meta-analysis. *Circulation* 108:2648-2652
118. Bates AS, Van't Hoff W, Jones PJ, Clayton RN 1996 The effect of hypopituitarism on life expectancy. *J Clin Endocrinol Metab* 81:1169-1172
119. Tomlinson JW, Holden N, Hills RK, Wheatley K, Clayton RN, Bates AS, Sheppard MC, Stewart PM 2001 Association between premature mortality and hypopituitarism. West Midlands Prospective Hypopituitary Study Group. *Lancet* 357:425-431
120. Rosen T, Bengtsson BA 1990 Premature mortality due to cardiovascular disease in hypopituitarism. *Lancet* 336:285-288
121. Bulow B, Hagmar L, Mikoczy Z, Nordstrom CH, Erfurth EM 1997 Increased cerebrovascular mortality in patients with hypopituitarism. *Clin Endocrinol (Oxf)* 46:75-81
122. Nilsson B, Gustavasson-Kadaka E, Bengtsson BA, Jonsson B 2000 Pituitary adenomas in Sweden between 1958 and 1991: incidence, survival and mortality. *J Clin Endocrinol Metab* 85:1420-1425
123. Erfurth EM, Bulow B, Nordstrom CH, Mikoczy Z, Hagmar L, Stromberg U 2004 Doubled mortality rate in irradiated patients reoperated for regrowth of a macroadenoma of the pituitary gland. *Eur J Endocrinol* 150:497-502
124. Holmes SJ, McKenna SP, Doward LC, Hunt SM, Shalet SM 1995 Development of a questionnaire to assess the quality of life of adults with growth hormone deficiency. *Endocrinol Metab* 2:63-69
125. Rosilio M, Blum WF, Edwards DJ, Shavrikova EP, Valle D, Lamberts SW, Erfurth EM, Webb SM, Ross RJ, Chihara K, Henrich G, Herschbach P, Attanasio AF 2004 Long-term improvement of quality of life during growth hormone (GH) replacement therapy in adults with GH deficiency, as measured by questions on life satisfaction-hypopituitarism (QLS-H). *J Clin Endocrinol Metab* 89:1684-1693

126. **Holmes SJ, Shalet SM** 1995 Factors influencing the desire for long-term growth hormone replacement in adults. *Clin Endocrinol (Oxf)* 43:151-157
127. **McGauley GA** 1989 Quality of life assessment before and after growth hormone treatment in adults with growth hormone deficiency. *Acta Paediatr Scand Suppl* 356:70-72; discussion, 73-74
128. **Murray RD, Shalet SM** 2002 Adult growth hormone replacement: lessons learned and future direction. *J Clin Endocrinol Metab* 87:4427-4428
129. **Murray RD, Skillicorn CJ, Howell SJ, Lissett CA, Rahim A, Smethurst LE, Shalet SM** 1999 Influences on quality of life in GH deficient adults and their effect on response to treatment. *Clin Endocrinol (Oxf)* 51:565-573
130. **Baum HB, Katznelson L, Sherman JC, Biller BM, Hayden DL, Schoenfeld DA, Cannistraro KE, Klibanski A** 1998 Effects of physiological growth hormone (GH) therapy on cognition and quality of life in patients with adult-onset GH deficiency. *J Clin Endocrinol Metab* 83:3184-3189
131. **Drake WM, Coyte D, Camacho-Hubner C, Jivanji NM, Kaltsas G, Wood DF, Trainer PJ, Grossman AB, Besser GM, Monson JP** 1998 Optimizing growth hormone replacement therapy by dose titration in hypopituitary adults. *J Clin Endocrinol Metab* 83:3913-3919
132. **Gilchrist FJ, Murray RD, Shalet SM** 2002 The effect of long-term untreated growth hormone deficiency (GHD) and 9 years of GH replacement on the quality of life (QoL) of GH-deficient adults. *Clin Endocrinol (Oxf)* 57:363-370
133. **Johannsson G, Rosen T, Bengtsson BA** 1997 Individualized dose titration of growth hormone (GH) during GH replacement in hypopituitary adults. *Clin Endocrinol (Oxf)* 47:571-581
134. **Jorgensen JO, Pedersen SA, Thuesen L, Jorgensen J, Ingemann-Hansen T, Skakkebaek NE, Christiansen JS** 1989 Beneficial effects of growth hormone treatment in GH-deficient adults. *Lancet* 1:1221-1225
135. **Whitehead HM, Boreham C, McIlrath EM, Sheridan B, Kennedy L, Atkinson AB, Hadden DR** 1992 Growth hormone treatment of adults with growth hormone deficiency: results of a 13-month placebo controlled cross-over study. *Clin Endocrinol (Oxf)* 36:45-52
136. **Binnerts A, Swart GR, Wilson JH, Hoogerbrugge N, Pols HA, Birkenhager JC, Lamberts SW** 1992 The effect of growth hormone administration in growth hormone deficient adults on bone, protein, carbohydrate and lipid homeostasis, as well as on body composition. *Clin Endocrinol (Oxf)* 37:79-87
137. **Chong PK, Jung RT, Scrimgeour CM, Rennie MJ, Paterson CR** 1994 Energy expenditure and body composition in growth hormone deficient adults on exogenous growth hormone. *Clin Endocrinol (Oxf)* 40:103-110
138. **Amato G, Carella C, Fazio S, La Montagna G, Cittadini A, Ssabatini D, Marciano-Mone C, Sacca L, Bellastella A** 1993 Body composition, bone metabolism, and heart structure and function in growth hormone (GH)-deficient adults before and after GH replacement therapy at low doses. *J Clin Endocrinol Metab* 77:1671-1676
139. **Orme SM, Sebastian JP, Oldroyd B, Stewart SP, Grant PJ, Stickland MH, Smith MA, Belchetz PE** 1992 Comparison of measures of body composition in a trial of low dose growth hormone replacement therapy. *Clin Endocrinol (Oxf)* 37:453-459
140. **O'Halloran DJ, Tsatsoulis A, Whitehouse RW, Holmes SJ, Adams JE, Shalet SM** 1993 Increased bone density after recombinant human growth hormone (GH) therapy in adults with isolated GH deficiency. *J Clin Endocrinol Metab* 76:1344-1348
141. **Vandeweghe M, Taelman P, Kaufman JM** 1993 Short- and long-term effects of growth hormone treatment on bone turnover and bone mineral content in adult growth hormone-deficient males. *Clin Endocrinol (Oxf)* 39:409-415
142. **Hoffman AR, Strasburger CJ, Zagar A, Blum WF, Kehely A, Hartman ML; T002 Study Group** 2004 Efficacy and tolerability of an individualized dosing regimen for adult growth hormone replacement therapy in comparison with fixed body weight-based dosing. *J Clin Endocrinol Metab* 89:3224-3233.
143. **Kehely A, Bates PC, Frewer P, Birkett M, Blum WF, Mamessier P, Ezzat S, Ho KK, Lombardi G, Luger A, Marek J, Russell-Jones D, Sonksen P, Attanasio AF** 2002 Short-term safety and efficacy of human GH replacement therapy in 595 adults with GH deficiency: a comparison of two dosage algorithms. *J Clin Endocrinol Metab* 87:1974-1979
144. 2004 Physicians' desk reference. Montvale, NJ: Medical Economics Co.
145. **Bryant J, Loveman E, Cave C, Chase D, Milne R** 2002 Endocrinology trial design: adverse event reporting in randomised controlled trials of recombinant human GH in GH-deficient adults. *J Endocrinol* 175:545-552
146. **Cummings DE, Merriam GR** 2003 Growth hormone therapy in adults. *Annu Rev Med* 54:513-533
147. **Carroll PV, Christ ER, Bengtsson BA, Carlsson L, Christiansen JS, Clemmons D, Hintz R, Ho K, Laron Z, Sizonenko P, Sonksen PH, Tanaka T, Thorne M** 1998 Growth hormone deficiency in adulthood and the effects of growth hormone replacement: a review. *Growth Hormone*

Research Society Scientific Committee. *J Clin Endocrinol Metab* 83:382-395

148. **Holmes SJ, Shalet SM** 1995 Which adults develop side-effects of growth hormone replacement? *Clin Endocrinol (Oxf)* 43:143-149
149. **Thuesen L, Jorgensen JO, Muller JR, Kristensen BO, Skakkebaek NE, Vahl N, Christiansen JS** 1994 Short- and long-term cardiovascular effect of growth hormone therapy in growth hormone deficient adults. *Clin Endocrinol (Oxf)* 41:615-620
150. **Koller EA, Green L, Gertner JM, Bost M, Malozowski S** 1998 Retinal changes mimicking diabetic retinopathy in two nondiabetic, growth hormone-treated patients. *J Clin Endocrinol Metab* 83:2380-2383
151. **Koller EA, Green L, Malozowski S** 2000 Comment on growth hormone therapy and retinal changes mimicking diabetic retinopathy. *J Clin Endocrinol Metab* 85:923 (Letter)
152. **Hansen R, Koller EA, Malozowski S** 2000 Full remission of growth hormone (GH)-induced retinopathy after GH treatment discontinuation: long-term follow-up. *J Clin Endocrinol Metab* 85:2627
153. **Radetti G, Gentili L, Lepidi M** 2000 Comment on growth hormone therapy and retinal changes mimicking diabetic retinopathy. *J Clin Endocrinol Metab* 85:923 (Letter)
154. **Malozowski S, Tanner LA, Wysowski D, Fleming GA, Stadel BV** 1995 Benign intracranial hypertension in children with growth hormone deficiency treated with growth hormone. *J Pediatr* 126:996-999
155. **Malozowski S, Tanner LA, Wysowski D, Fleming GA** 1993 Growth hormone, insulin-like growth factor I, and benign intracranial hypertension. *N Engl J Med* 329:665-666
156. **Cohn L, Feller AG, Draper MW, Rudman IW, Rudman D** 1993 Carpal tunnel syndrome and gynaecomastia during growth hormone treatment of elderly men with low circulating IGF-I concentrations. *Clin Endocrinol (Oxf)* 39:417-425
157. **Blackman MR, Sorkin JD, Münzer T, Bellantoni MF, Busby-Whitehead J, Stevens TE, Jayme J, O'Connor KG, Christmas C, Tobin JD, Stewart KJ, Cottrell E, St Clair C, Pabst KM, Harman SM** 2002 Growth hormone and sex steroid administration in healthy aged women and men: a randomized controlled trial. *JAMA* 288:2282-2292
158. **Tuffli GA, Johanson A, Rundle AC, Allen AB** 1995 Lack of increased risk for extracranial, nonleukemic neoplasms in recipients of recombinant deoxyribonucleic acid growth hormone. *J Clin Endocrinol Metab* 80:1416-1422
159. **Hatrack AG, Boghalo P, Bingham JB, Ayres AB, Sonksen PH, Russell-Jones DL** 2002 Does GH replacement therapy in adult GH-deficient patients result in recurrence or increase in size of pituitary tumours? *Eur J Endocrinol* 146:807-811
160. **Fradkin JE, Mills JL, Schonberger LB, Wysowski DK, Thomson R, Durako SJ, Robison LL** 1993 Risk of leukemia after treatment with pituitary growth hormone. *JAMA* 270:2829-2832
161. **Swerdlow AJ, Higgins CD, Adlard P, Preece MA** 2002 Risk of cancer in patients treated with human pituitary growth hormone in the UK, 1959-85: a cohort study. *Lancet* 360:273-277
162. **Porretti S, Giavoli C, Ronchi C, Lombardi G, Zaccaria M, Valle D, Arosio M, Beck-Peccoz P** 2002 Recombinant human GH replacement therapy and thyroid function in a large group of adult GH-deficient patients: when does L-T(4) therapy become mandatory? *J Clin Endocrinol Metab* 87:2042-2045
163. **Giavoli C, Libé R, Corbetta S, Ferrante E, Lania A, Arosio M, Spada A, Beck-Peccoz P** 2004 Effect of recombinant human growth hormone (GH) replacement on the hypothalamic-pituitary-adrenal axis in adult GH-deficient patients. *J Clin Endocrinol Metab* 89:5397-5401
164. **Burman P, Johansson AG, Siegbahn A, Vessby B, Karlsson FA** 1997 Growth hormone (GH)-deficient men are more responsive to GH replacement than women. *J Clin Endocrinol Metab* 82:550-555
165. **Cook DM, Ludlam WH, Cook MB** 1999 Route of estrogen administration helps to determine growth hormone (GH) replacement dose in GH-deficient adults. *J Clin Endocrinol Metab* 84:3956-3960
166. **Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, Guyatt GH, Harbour RT, Haugh MC, Henry D, Hill S, Jaeschke R, Leng G, Liberati A, Magrini N, Mason J, Middleton P, Mrukowicz J, O'Connell D, Oxman AD, Phillips B, Schunemann HJ, Edejer TT, Varonen H, Vist GE, Williams Jr JW, Zaza S; for the GRADE Working Group** 2004 Grading quality of evidence and strength of recommendations. *BMJ* 328:1490

Acknowledgments

The members of the Task Force thank the members of the Clinical Guidelines Subcommittee, the Clinical Affairs Committee, and The Endocrine Society Council for their careful, critical review of earlier versions of this manuscript and their helpful comments and suggestions. In addition, we thank the many members of The Endocrine Society who reviewed the draft version of this manuscript when it was posted on the Society Web site and who sent a great number of additional comments and suggestions, most of which were incorporated into the final version of the manuscript. We thank Patricia A. Stephens, Ph.D., medical writer on this Guideline, who meticulously checked the references and formatted the Guideline into its current form. Finally, we thank the staff at The Endocrine Society office for their helpful support during the development of this Guideline.

Financial Disclosure of Task Force: Mark E. Molitch, M.D., Consultant or Adviser: Abbott Laboratories, Novo Nordisk, Novartis, Pfizer, Sanofi-Aventis; Research Support: Eli Lilly & Co., Novartis, Pfizer, Genentech, Amgen, Sanofi-Aventis; David R. Clemmons, M.D., Consultant or Adviser: Eli Lilly & Co., Pfizer; Research Support: Pfizer; Other: reviewed grants for Genentech; Saul Malozowski, M.D., Ph.D., None; George R. Merriam, M.D., Consultant or Adviser: Elixir, Genentech, LG Life Sciences, Theratechnologies, Tokai; Research Support: Genentech, Eli Lilly & Co., LG Life Sciences Pfizer; Stephen M. Shalet, M.D., Consultant or Adviser: Transpharma, Skyepharma; Research Support: Pfizer, Novo Nordisk, Novartis; Mary Lee Vance, M.D., Consultant or Adviser: Genentech, Novartis, Pfizer; Research Support: Eli Lilly & Co., Novartis, Genentech, Pfizer.

What goes into our Clinical Guidelines *is a story worth telling*

The extensive process that goes into creating The Endocrine Society's Clinical Guidelines not only provides validation and assurance, but also raises the standard for the development of guidelines everywhere.

The guidelines are developed using a multi-step process that reflects the standards of excellence embraced by The Endocrine Society.

Other Endocrine Society Guidelines Coming Soon:

- 1. Testosterone Therapy in Adult Men with Androgen Deficiency Syndromes**
- 2. Therapeutic Use of Androgens in Women**
- 3. Prevention and Treatment of Pediatric Obesity**
- 4. Management of Thyroid Dysfunction During Pregnancy and Postpartum**
- 5. Metabolic Syndrome**
- 6. Diagnosis and Evaluation of Women with Hirsutism**
- 7. Lipids**
- 8. Vitamin D & Bone**



For more information on The Endocrine Society's Clinical Guidelines, visit www.endo-society.org/quickcontent/clinicalpractice/clinical-guidelines/.





8401 Connecticut Avenue, Suite 900
 Chevy Chase, MD 20815-5817
 Phone (301) 941-0210; Fax (301) 941-0257
 societyservices@endo-society.org
 FEIN 73-0521256

THE ENDOCRINE SOCIETY GUIDELINE ORDER FORM

(Single reprint request for orders of 100 and less)

PRODUCTS	QUANTITY	PRICE		SUBTOTAL
		Member	Non-Member	
Evaluation and Treatment of Adult Growth Hormone Deficiency: An Endocrine Society Clinical Practice Guideline		\$15.00	\$20.00	
Miscellaneous				
TOTAL		All prices include sales tax		\$

PAYMENT INFORMATION: Check MasterCard Visa

Card Number	Expiration Date
Billing Address	Signature

Are you a member of The Endocrine Society? Yes No

If you are a member and you know your member ID, please provide: _____

Prefix:	First Name (Given):	Middle:	Last (Surname):												
Institution/Company:		Dept/Div:													
Street/PO:															
City:	State/Province:	Zip/Mail Code:	Country:												
Telephone:	Fax:	Email:													
Degree(s) that you would like listed after your name:		Professional Title:	Date of Birth: Gender: <input type="radio"/> Male <input type="radio"/> Female												
Which of the following best describes your primary professional role? (Please mark only one) <table style="width: 100%; margin-top: 5px;"> <tr> <td><input type="radio"/> Administrator</td> <td><input type="radio"/> Retired</td> </tr> <tr> <td><input type="radio"/> Basic Researcher</td> <td><input type="radio"/> Teacher/Educator</td> </tr> <tr> <td><input type="radio"/> Clinical Practitioner</td> <td><input type="radio"/> Fellow (Clinical)</td> </tr> <tr> <td><input type="radio"/> Clinical Researcher</td> <td><input type="radio"/> Fellow (Postdoctoral/ Research)</td> </tr> <tr> <td><input type="radio"/> Industry/Corporate Professional</td> <td><input type="radio"/> Student</td> </tr> <tr> <td><input type="radio"/> Nurse/Healthcare Professional</td> <td><input type="radio"/> Other _____</td> </tr> </table>			<input type="radio"/> Administrator	<input type="radio"/> Retired	<input type="radio"/> Basic Researcher	<input type="radio"/> Teacher/Educator	<input type="radio"/> Clinical Practitioner	<input type="radio"/> Fellow (Clinical)	<input type="radio"/> Clinical Researcher	<input type="radio"/> Fellow (Postdoctoral/ Research)	<input type="radio"/> Industry/Corporate Professional	<input type="radio"/> Student	<input type="radio"/> Nurse/Healthcare Professional	<input type="radio"/> Other _____	Race or Ethnic Affiliation (voluntary) <ul style="list-style-type: none"> <input type="radio"/> African American, Black <input type="radio"/> Asian <input type="radio"/> Hispanic <input type="radio"/> Native American, Eskimo, Aleut <input type="radio"/> Pacific Islander <input type="radio"/> White, Caucasian <input type="radio"/> Other _____
<input type="radio"/> Administrator	<input type="radio"/> Retired														
<input type="radio"/> Basic Researcher	<input type="radio"/> Teacher/Educator														
<input type="radio"/> Clinical Practitioner	<input type="radio"/> Fellow (Clinical)														
<input type="radio"/> Clinical Researcher	<input type="radio"/> Fellow (Postdoctoral/ Research)														
<input type="radio"/> Industry/Corporate Professional	<input type="radio"/> Student														
<input type="radio"/> Nurse/Healthcare Professional	<input type="radio"/> Other _____														

(Commercial reprint information on inside back cover)

Commercial Reprint Information

For information on reprint requests of more than 101 and commercial reprints contact:

Heather Edwards
Reprint Sales Specialist
Cadmus Professional Communications

Phone: 410.691.6214
Fax: 410.684.2789
Email: endoreprints@cadmus.com

Single Reprint Information

To order reprints of 100 and less, complete the guideline order form and return using one of the following methods:

Mail: The Endocrine Society
c/o Bank of America
P.O. Box 630721
Baltimore, MD 21263-0736

Phone: 301.941.0210
Fax: 301.941.0257
Email: Societyservices@endo-society.org

Questions & Correspondences

The Endocrine Society
Attn: Government & Professional Affairs Department
8401 Connecticut Avenue, Suite 900
Chevy Chase, MD 20815

Phone: 301.941.0200
Email: govt-prof@endo-society.org
Web: www.endo-society.org

For more information on The Endocrine Society's Clinical Practice Guidelines, visit www.endo-society.org/quickcontent/clinicalpractice/clinical-guidelines/.



The Endocrine Society
8401 Connecticut Avenue, Suite 900
Chevy Chase, MD 20815

301.941.0200
www.endo-society.org