

# Exhibit 465

Report 65: In the First Three Months of Pfizer's mRNA  
"Vaccine" Rollout, Nine Patients Died of Anaphylaxis.  
79% of Anaphylaxis Adverse Events Were  
Rated as "Serious."

<https://dailyclout.io/report-65-in-the-first-three-months-of-pfizers-mrna-vaccine-rollout-nine-patients-died-of-anaphylaxis/>

(<https://dailyclout.io/>)



(<https://dailyclout.io/cart/>)

(<https://www.facebook.com/dailyclout/>)

(<https://twitter.com/DailyClout?lang=en>)

(<https://www.youtube.com/channel/UCU-FMBZNtCdSiYBgJdvJmXw>)

(<https://www.instagram.com/dailyclout.io/>)

[ALL POSTS](#)

[BULLETIN BOARD](#)

[DAILYCLOUT LAWSUIT](#)

[OPINION](#)

[PEZIZER REPORTS](#)

Search



DailyClout



Pfizer Reports

## Report 65: In the First Three Months of Pfizer's mRNA "Vaccine" Rollout, Nine Patients Died of Anaphylaxis. 79% of Anaphylaxis Adverse Events Were Rated as "Serious."

March 31, 2023 • by Joseph Gehrett, MD; Barbara Gehrett, MD; Chris Flowers, MD; and Loree Britt

The War Room/DailyClout Pfizer Documents Analysis Project Post-Marketing Group (Team 1) – Joseph Gehrett, MD; Barbara Gehrett, MD; Chris Flowers, MD; and Loree Britt – produced an shocking review of anaphylaxis (severe allergy reaction) adverse events found in Pfizer document

*5.3.6 Cumulative Analysis of Post-Authorization Adverse Event Reports of PF-07302048 (BNT162B2) Received Through 28-FEB-2021* ([https://www.phmpt.org/wp-content/uploads/2022/04/reissue\\_5.3.6-postmarketing-experience.pdf](https://www.phmpt.org/wp-content/uploads/2022/04/reissue_5.3.6-postmarketing-experience.pdf))

(a.k.a., "5.3.6"). Severe allergic reaction is typically triggered by latex, foods such as peanuts, bee stings, or medications (injected or taken by mouth), among other things, and is generally considered a **medical emergency**.

It is important to note that the adverse events (AEs) in the 5.3.6 document were reported to Pfizer for **only a 90-day period** starting on December 1, 2020, the date of the United Kingdom's public rollout of Pfizer's COVID-19 experimental mRNA

"vaccine" product.

Key points in this report include:

- There were **nine reported deaths**.
  - Only four patients who died were reported to have serious, underlying medical conditions that "likely contributed to their deaths."
- There were 1,833 potential anaphylaxis patients reported; but, after screening using a tool called [Brighton Collaboration \(https://brightoncollaboration.us/\)](https://brightoncollaboration.us/), 831 did not meet anaphylaxis criteria, leaving **1,002 cases reported in 90 days**. Pfizer reported 2,958 "potentially relevant events" from those 1,002 individuals that included the signs and symptoms of anaphylaxis.
  - Where did the other 831 patients and their allergic adverse events go/get assigned in the post-marketing data, if anywhere?
- **Pfizer reported only the most frequent anaphylaxis signs and symptoms:** anaphylactic reaction (435), shortness of breath (356), rash (190), redness of the skin (159), hives (133), cough (115), **respiratory distress (97)**, throat tightness (97), swollen tongue (93), low blood pressure (72), **low blood pressure severe enough to threaten organ function (shock) (80)**, chest discomfort (71), swelling face (70), throat swelling (68), and lip swelling (64).
- The events were rated as **serious in 2,341 (79%)** and non-serious in 617 (21%).
- The **ratio of females to males affected was over 8:1**. Of those cases with gender specified, **876 (89%) were female**, 106 (11%) were male.
- Half of patients with this adverse event were **younger than 43.5 years old**.
- Pfizer's Conclusion: Evaluation of BC (*Brighton Collaborative*) cases Level 1-4 **did not reveal any significant new safety information**. Anaphylaxis is appropriately described in the product labeling as are non-anaphylactic hypersensitivity events. Surveillance will continue.

Read this important two-page report below:



## War Room/DailyClout Pfizer Document Analysis

### SOURCE:

<https://www.phmpt.org/wp-content/uploads/2022/04/reissue-5.3.6-postmarketing-experience.pdf>

### 5.3.6 AE REPORTING PERIOD:

"Since the first temporary authorization for emergency supply under Regulation 174 in the UK (01 December 2020) and through 28 February 2021."

### 5.3.6 AE CASES/EVENTS:

TOTAL AE CASES: 42,086  
TOTAL AE EVENTS: 158,893

### ABBREVIATIONS:

5.3.6 : Pfizer source document

SOC : System Organ Class

AE : Adverse Event

AESI : Adverse Event of Special Interest

EUA : Emergency Use Authorization by FDA

PM : Post-Marketing

BNT162b2 : Pfizer's mRNA COVID-19 vaccine

SEQUELAE: an abnormal condition resulting from a previous disease, injury, or other trauma

AGE GROUPS defined in 5.3.6 (p. 25 footnote) :

Adult	18 - 64
Elderly	≥ 65
Child	2 - 11
Adolescent	12 - < 18
Infant	1 - 23 months

### AUTHORS:

Dr Joseph Gehrett MD  
Dr Barbara Gehrett MD  
Dr Chris Flowers MD  
Loree Britt



27Mar23

## Post-Marketing Team Micro-Report 10:

### Anaphylaxis – Important Identified Risk Review of 5.3.6

**Anaphylaxis** Since the first temporary authorization for emergency supply under Regulation 174 in the UK (01 December 2020) and through 28 February 2021, 1,833 potentially relevant cases were retrieved from the Anaphylactic reaction SMQ (Narrow and Broad) search strategy, applying the MedDRA algorithm. These cases were individually reviewed and assessed according to Brighton Collaboration (BC) definition and level of diagnostic certainty as shown in the Table below:

Brighton Collaboration Level	Number of cases
BC 1	299
BC 2	311
BC 3	10
BC 4	391
BC 5	831
Total	1,833

Level 1 indicates a case with the highest level of diagnostic certainty of anaphylaxis, whereas the diagnostic certainty is lowest for Level 3. Level 4 is defined as "reported event of anaphylaxis with insufficient evidence to meet the case definition" and Level 5 as not a case of anaphylaxis.

There were 1,002 cases (54.0% of the potentially relevant cases retrieved), 2,958 potentially relevant events, from the Anaphylactic reaction SMQ (Broad and Narrow) search strategy, meeting BC Level 1 to 4.

Country of incidence: UK (261), US (184), Mexico (99), Italy (82), Germany (67), Spain (38), France (36), Portugal (22), Denmark (20), Finland, Greece (19 each), Sweden (17), Czech Republic, Netherlands (16 each), Belgium, Ireland (13 each), Poland (12), Austria (11); the remaining 57 cases originated from 15 different countries.

Relevant event seriousness: Serious (2,341), Non-Serious (617);

Gender: Female (876), Male (180), Unknown (20);

Age (n=961) ranged from 16 to 98 years (mean = 54.8 years, median = 42.5 years);

Relevant event outcome\*: final (97), resolved/resolving (1,922), not resolved (229), resolved with sequelae (48), unknown (754).

Most frequently reported relevant PFs (≥2%), from the Anaphylactic reaction SMQ (Broad and Narrow) search strategy: Anaphylactic reaction (435), Depression (156), Rash (106), Pruritus (73), Erythema (39), Urticaria (133), Cough (115), Respiratory distress, Throat tightness (97 each), Swollen tongue (93), Anaphylactic shock (80), Hypotension (72), Chest discomfort (71), Swelling face (70), Periorbital swelling (68), and Lip swelling (64).

Conclusion: Evaluation of BC cases Level 1 - 4 did not reveal any significant new safety information. Anaphylaxis is appropriately described in the product labeling as are non-anaphylactic hypersensitivity events. Surveillance will continue.

[https://www.phmpt.org/wpcontent/uploads/2022/04/reissue\\_5.3.6-postmarketexperience.pdf](https://www.phmpt.org/wpcontent/uploads/2022/04/reissue_5.3.6-postmarketexperience.pdf)

Anaphylaxis is a condition most often referred to as a "severe allergic reaction" and is triggered by latex, foods such as peanuts, bee stings, or medications (injected or taken by mouth), among other things. Immunoglobulin E (IgE, a human antibody) is involved in this dangerous and explosive reaction. It causes mast cells to immediately release histamine and other chemicals.

The life-threatening symptoms that follow can include swelling such as hives, respiratory difficulties, drop in blood pressure and rapid heart rate, and abdominal symptoms such as pain, nausea and vomiting. These symptoms are typically rapid in both onset and progression. Treatment similarly has to be rapid (e.g., Epi-Pen) and often requires emergency room or hospital treatment.

Pfizer has used a tool, the Brighton Collaboration (BC), to assess whether the symptoms of a patient are correctly identified as anaphylaxis. In the three-month data collection, there were 1,833 potential anaphylaxis patients reported; but, after screening, 831 did not meet criteria, leaving 1,002 cases reported in this time period. Pfizer reported 2,958 "potentially relevant events" from those 1,002 individuals that included the signs and symptoms of anaphylaxis.

This report was written exclusively for DailyClout by the members of the War Room/DailyClout Pfizer Documents Analysis Project team. It may not be copied or republished without written permission from DailyClout or a full credit and link to DailyClout.io.

- Adverse Events were reported to Pfizer during a 90-day period, following the December 1, 2020, public rollout of its COVID-19 experimental "vaccine" product.
- In the Pfizer 5.3.6 document, these AEs were categorized by System Organ Classes (SOC) – in other words, by systems in the body.
- Of those with gender specified, 876 (89%) were female, 106 (11%) were male.
- The reported age range was 16 to 98 years old with half being less than 43.5 years old.

Pfizer reported only the most frequent anaphylaxis signs and symptoms: anaphylactic reaction (435), shortness of breath (356), rash (190), redness of the skin (159), hives (133), cough (115), respiratory distress (97), throat tightness (97), swollen tongue (93), low blood pressure (72), low blood pressure severe enough to threaten organ function (shock) (80), chest discomfort (71), swelling face (70), throat swelling (68), and lip swelling (64).

The events were rated as serious in 2,341 (79%) and non-serious in 617 (21%). 1,922 events were reported as resolved or resolving (65%), 229 not resolved (8%), 48 resolved with sequelae (1.6%), and 754 had no outcome known (25%).

### There were nine deaths reported.

#### IMPORTANT NOTE:

There are 831 patients, 45% of the total 1,833, who were determined not to be anaphylaxis and dropped from this category. Where did those patients and their adverse events go?

<https://dailyclout.io/wp-content/uploads/Post-Marketing-Anaphylaxis-microreport-p1.pdf> (<https://dailyclout.io/wp-content/uploads/Post-Marketing-Anaphylaxis-microreport-p1.pdf>)



**SOURCE:**  
[https://www.phmpt.org/wp-content/uploads/2022/04/reissue\\_5\\_3.6-postmarketing-experience.pdf](https://www.phmpt.org/wp-content/uploads/2022/04/reissue_5_3.6-postmarketing-experience.pdf)



Pfizer had 1,002 case reports in the anaphylaxis category, or 2.4% of the total 42,086 adverse event patients from all causes. They categorized anaphylaxis under "Important identified risks." Anaphylaxis is generally considered a medical emergency. In this report, **79% of the events were considered serious**. In spite of anaphylaxis being treatable, it remains a potentially fatal condition as demonstrated by the **nine deaths** reported.

Pfizer noted that four of the patients who died had "serious underlying medical conditions" that "likely contributed to their deaths." **Adverse events in the high-risk patients offset the potential benefits of the immunization and must be part of the consideration for approval.** Of course, those who are medically fragile will be at greater risk with any adverse event.

**Females comprised 89% of the anaphylaxis reports** compared to 77% of the total cases in the post-authorization compilation. There is no comment from Pfizer on the marked female predominance in this adverse event category, let alone any stated plan to address it.

---

**Pfizer's Conclusion:**  
 Evaluation of BC (*Brighton Collaborative*) cases Level 1 - 4 did not reveal any significant new safety information. Anaphylaxis is appropriately described in the product labeling as are non-anaphylactic hypersensitivity events. Surveillance will continue.



---

**Post-Marketing Team's CONCLUSION**  
**RECALL this unsafe "vaccine."**

---

This report was written exclusively for DailyClout by the members of the War Room/DailyClout Pfizer Documents Analysis Project team. It may not be copied or republished without written permission from DailyClout or a full credit and link to [DailyClout.io](https://dailyclout.io).

<https://dailyclout.io/wp-content/uploads/Post-Marketing-Anaphylaxis-microreport-p2-1.pdf> (<https://dailyclout.io/wp-content/uploads/Post-Marketing-Anaphylaxis-microreport-p2-1.pdf>)

**Pre-Order Your Copy of the Pfizer Reports Paperback Book.**  
(<https://dailyclout.io/product/war-room-dailyclout-pfizer-documents-analysis-volunteers-reports-book-paperback/>)

**Please support DailyClout.** (<https://dailyclout.io/donate/>)  
**Become a DailyClout Member.** (<https://dailyclout.io/memberships/>)

**Spread the love**

(<https://www.facebook.com/sharer/sharer.php?u=https%3A%2F%2Fdailyclout.io%2Freport-65-in-the-first-three-months-of-pfizers-mrna-vaccine-rollout-nine-patients-died-of-anaphylaxis%2F>)

(<http://twitter.com/intent/tweet?text=Report%2065%3A%20In%20the%20First%2065-in-the-first-three-months-of-pfizers-mrna-vaccine-rollout-nine-patients-died-of-anaphylaxis%2F>)

(<https://gettr.com/share?text=Report%2065%3A%20In%20the%20First%2065-in-the-first-three-months-of-pfizers-mrna-vaccine-rollout-nine-patients-died-of-anaphylaxis/>)

**NEXT STORY**

**PREVIOUS STORY**

**This is The New Beginning of Real Medicine**

(<https://dailyclout.io/this-is-the-new-beginning-of-real-medicine/>)

**Vax Death Count = COVID Death Count, According to Latest Rasmussen Reports Poll**

(<https://dailyclout.io/vax-death-count-covid-death-count/>)