

Mastering Chemistry With Ravi Arora

DPP - Daily Practice Problems

Chapter-wise Sheets

Date :

Start Time :

End Time :

CHEMISTRY (CC17)

SYLLABUS : Electrochemistry

Max. Marks : 120

Marking Scheme : + 4 for correct & (-1) for incorrect

Time : 60 min.

INSTRUCTIONS : This Daily Practice Problem Sheet contains 30 MCQ's. For each question only one option is correct. Darken the correct circle/ bubble in the Response Grid provided on each page.

1. Which of the following expressions correctly represents the equivalent conductance at infinite dilution of $\text{Al}_2(\text{SO}_4)_3$,
Given that $\Lambda_{\text{Al}^{3+}}^\circ$ and $\Lambda_{\text{SO}_4^{2-}}^\circ$ are the equivalent conductances at infinite dilution of the respective ions?
- (a) $\frac{1}{3}\Lambda_{\text{Al}^{3+}}^\circ + \frac{1}{2}\Lambda_{\text{SO}_4^{2-}}^\circ$ (b) $2\Lambda_{\text{Al}^{3+}}^\circ + 3\Lambda_{\text{SO}_4^{2-}}^\circ$
(c) $\Lambda_{\text{Al}^{3+}}^\circ + \Lambda_{\text{SO}_4^{2-}}^\circ$ (d) $(\Lambda_{\text{Al}^{3+}}^\circ + \Lambda_{\text{SO}_4^{2-}}^\circ) \times 6$
2. The equivalent conductance of $\frac{M}{32}$ solution of a weak monobasic acid is 8.0 mho cm^2 and at infinite dilution is 400 mho cm^2 . The dissociation constant of this acid is:
- (a) 1.25×10^{-6} (b) 6.25×10^{-4}
(c) 1.25×10^{-4} (d) 1.25×10^{-5}
3. Aqueous solution of which of the following compounds is the best conductor of electric current ?
- (a) Acetic acid, $\text{C}_2\text{H}_4\text{O}_2$
(b) Hydrochloric acid, HCl
(c) Ammonia, NH_3
(d) Fructose, $\text{C}_6\text{H}_{12}\text{O}_6$
4. The standard EMF of Daniell cell is 1.10 volt. The maximum electrical work obtained from the Daniell cell is
- (a) 212.3 kJ
(b) 175.4 kJ
(c) 106.15 kJ
(d) 53.07 kJ

RESPONSE GRID

1. (a)(b)(c)(d) 2. (a)(b)(c)(d) 3. (a)(b)(c)(d) 4. (a)(b)(c)(d)


Space for Rough Work

5. Which of the following reaction occurs at the cathode during the charging of lead storage battery?
- (a) $\text{Pb}^{2+} + 2\text{e}^- \longrightarrow \text{Pb}$
 (b) $\text{Pb}^{2+} + \text{SO}_4^{2-} \longrightarrow \text{PbSO}_4$
 (c) $\text{Pb} \longrightarrow \text{Pb}^{2+} + 2\text{e}^-$
 (d) $\text{PbSO}_4 + 2\text{H}_2\text{O} \longrightarrow 2\text{PbO}_2 + 4\text{H}^+ + \text{SO}_4^{2-} + 2\text{e}^-$
6. Molar ionic conductivities of a two-bivalent electrolytes x^{2+} and y^{2-} are 57 and 73 respectively. The molar conductivity of the solution formed by them will be
- (a) $130 \text{ S cm}^2 \text{ mol}^{-1}$
 (b) $65 \text{ S cm}^2 \text{ mol}^{-1}$
 (c) $260 \text{ S cm}^2 \text{ mol}^{-1}$
 (d) $187 \text{ S cm}^2 \text{ mol}^{-1}$
7. Kohlrausch's law states that at :
- (a) finite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte, whatever be the nature of the other ion of the electrolyte.
 (b) infinite dilution each ion makes definite contribution to equivalent conductance of an electrolyte depending on the nature of the other ion of the electrolyte.
 (c) infinite dilution, each ion makes definite contribution to conductance of an electrolyte whatever be the nature of the other ion of the electrolyte.
 (d) infinite dilution, each ion makes definite contribution to equivalent conductance of an electrolyte, whatever be the nature of the other ion of the electrolyte.
8. Standard free energies of formation (in kJ/mol) at 298 K are -237.2 , -394.4 and -8.2 for $\text{H}_2\text{O}(l)$, $\text{CO}_2(g)$ and pentane (g), respectively. The value E°_{cell} for the pentane-oxygen fuel cell is :
- (a) 1.968 V
 (b) 2.0968 V
 (c) 1.0968 V
 (d) 0.0968 V
9. If the E°_{cell} for a given reaction has a negative value, then which of the following gives the correct relationships for the values of ΔG° and K_{eq} ?
- (a) $\Delta G^\circ > 0$; $K_{\text{eq}} > 1$ (b) $\Delta G^\circ < 0$; $K_{\text{eq}} > 1$
 (c) $\Delta G^\circ < 0$; $K_{\text{eq}} < 1$ (d) $\Delta G^\circ > 0$; $K_{\text{eq}} < 1$
10. Standard electrode potentials are : Fe^{2+}/Fe [$E^\circ = -0.44$]; $\text{Fe}^{3+}/\text{Fe}^{2+}$ $E^\circ = +0.77$; If Fe^{2+} , Fe^{3+} and Fe blocks are kept together, then
- (a) Fe^{3+} increases
 (b) Fe^{3+} decreases
 (c) $\frac{\text{Fe}^{2+}}{\text{Fe}^{3+}}$ remains unchanged
 (d) Fe^{2+} decreases
11. An electrolytic cell contains a solution of Ag_2SO_4 and has platinum electrodes. A current is passed until 1.6 gm of O_2 has been liberated at anode. The amount of silver deposited at cathode would be
- (a) 107.88 gm (b) 1.6 gm
 (c) 0.8 gm (d) 21.60 gm
12. If ϕ denotes reduction potential, then which is true?
- (a) $E^\circ_{\text{cell}} = \phi_{\text{right}} - \phi_{\text{left}}$ (b) $E^\circ_{\text{cell}} = \phi_{\text{left}} + \phi_{\text{right}}$
 (c) $E^\circ_{\text{cell}} = \phi_{\text{left}} - \phi_{\text{right}}$ (d) $E^\circ_{\text{cell}} = -(\phi_{\text{left}} + \phi_{\text{right}})$.
13. In a fuel cell methanol is used as fuel and oxygen gas is used as an oxidizer. The reaction is
- $$\text{CH}_3\text{OH}(l) + 3/2\text{O}_2(g) \longrightarrow \text{CO}_2(g) + 2\text{H}_2\text{O}(l)$$
- At 298 K standard Gibb's energies of formation for $\text{CH}_3\text{OH}(l)$, $\text{H}_2\text{O}(l)$ and $\text{CO}_2(g)$ are -166.2 , -237.2 and $-394.4 \text{ kJ mol}^{-1}$ respectively. If standard enthalpy of combustion of methanol is -726 kJ mol^{-1} , efficiency of the fuel cell will be:
- (a) 87% (b) 90%
 (c) 97% (d) 80%
14. For the cell reaction,
 $\text{Cu}^{2+}(\text{C}_1, \text{aq}) + \text{Zn}(s) = \text{Zn}^{2+}(\text{C}_2, \text{aq}) + \text{Cu}(s)$ of an electrochemical cell, the change in free energy, ΔG , at a given temperature is a function of
- (a) $\ln(\text{C}_1)$ (b) $\ln(\text{C}_2/\text{C}_1)$
 (c) $\ln(\text{C}_2)$ (d) $\ln(\text{C}_1 + \text{C}_2)$

**RESPONSE
GRID**

5. (a)(b)(c)(d) 6. (a)(b)(c)(d) 7. (a)(b)(c)(d) 8. (a)(b)(c)(d) 9. (a)(b)(c)(d)
 10. (a)(b)(c)(d) 11. (a)(b)(c)(d) 12. (a)(b)(c)(d) 13. (a)(b)(c)(d) 14. (a)(b)(c)(d)

Space for Rough Work

15. When electric current is passed through acidified water, 112 ml of hydrogen gas at STP collected at the cathode in 965 seconds. The current passed in amperes is
 (a) 1.0 (b) 0.5 (c) 0.1 (d) 2.0
16. The electrode potential $E_{(Zn^{2+}/Zn)}$ of a zinc electrode at 25°C with an aqueous solution of 0.1 M $ZnSO_4$ is
 $[E_{(Zn^{2+}/Zn)}^\circ = -0.76 \text{ V. Assume } \frac{2.303RT}{F} = 0.06 \text{ at } 298 \text{ K}.]$
 (a) +0.73 (b) -0.79
 (c) -0.82 (d) -0.70
17. A gas X at 1 atm is bubbled through a solution containing a mixture of 1 M Y^- and MZ^- at 25°C. If the reduction potential of $Z > Y > X$, then,
 (a) Y will oxidize X and not Z
 (b) Y will oxidize Z and not X
 (c) Y will oxidize both X and Z
 (d) Y will reduce both X and Z
18. For the electrochemical cell, $M | M^+ || X^- | X$,
 $E^\circ M^+ / M = 0.44 \text{ V}$ and $E^\circ (X/X^-) = 0.33 \text{ V}$.
 From this data one can deduce that
 (a) $M+X \rightarrow M^++X^-$ is the spontaneous reaction
 (b) $M^++X^- \rightarrow M+X$ is the spontaneous reaction
 (c) $E_{\text{cell}} = 0.77 \text{ V}$
 (d) $E_{\text{cell}} = -0.77 \text{ V}$
19. Standard electrode potential data are useful for understanding the suitability of an oxidant in a redox titration. Some half cell reactions and their standard potentials are given below:
 $MnO_4^-(\text{aq.}) + 8H^+(\text{aq.}) + 5e^- \rightarrow Mn^{2+}(\text{aq.}) + 4H_2O(l)$
 $E^\circ = 1.51 \text{ V}$
 $Cr_2O_7^{2-}(\text{aq.}) + 14H^+(\text{aq.}) + 6e^- \rightarrow 2Cr^{3+}(\text{aq.}) + 7H_2O(l)$
 $E^\circ = 1.38 \text{ V}$
 $Fe^{3+}(\text{aq.}) + e^- \rightarrow Fe^{2+}(\text{aq.})$ $E^\circ = 0.77 \text{ V}$
 $Cl_2(g) + 2e^- \rightarrow 2Cl^-(\text{aq.})$ $E^\circ = 1.40 \text{ V}$
 Identify the only incorrect statement regarding the quantitative estimation of aqueous $Fe(NO_3)_2$
 (a) MnO_4^- can be used in aqueous HCl
 (b) $Cr_2O_7^{2-}$ can be used in aqueous HCl
 (c) MnO_4^- can be used in aqueous H_2SO_4
 (d) $Cr_2O_7^{2-}$ can be used in aqueous H_2SO_4
20. A hypothetical electrochemical cell is shown below
 $A | A^+(xM) || B^+(yM) | B$
 The emf measured is +0.20 V. The cell reaction is
 (a) $A^+ + e^- \rightarrow A$; $B^+ + e^- \rightarrow B$
 (b) The cell reaction cannot be predicted
 (c) $A + B^+ \rightarrow A^+ + B$
 (d) $A^+ + B \rightarrow A + B^+$
21. Conductance of 0.1 M KCl (conductivity = $X \text{ Ohm}^{-1} \text{ cm}^{-1}$) filled in a conductivity cell is $Y \text{ Ohm}^{-1}$. If the conductance of 0.1 M NaOH filled in the same cell is $Z \text{ Ohm}^{-1}$, the molar conductance of NaOH will be
 (a) $10^3 \frac{XZ}{Y}$ (b) $10^4 \frac{XZ}{Y}$ (c) $10 \frac{XZ}{Y}$ (d) $0.1 \frac{XZ}{Y}$
22. On the basis of the following E° values, the strongest oxidizing agent is:
 $Fe(CN)_6^{4-} \rightarrow [Fe(CN)_6]^{3-} + e^-$; $E^\circ = -0.35 \text{ V}$
 $Fe^{2+} \rightarrow Fe^{3+} + e^-$; $E^\circ = -0.77 \text{ V}$
 (a) $[Fe(CN)_6]^{4-}$ (b) Fe^{2+}
 (c) Fe^{3+} (d) $[Fe(CN)_6]^{3-}$
23. The mathematical expression for law of independent migration of ions and Ostwald's dilution law are given by
 (a) $\Lambda = \Lambda_m^\circ - BC^{1/2}$ (b) $\Lambda^\circ = F(U_+ + U_-)$
 (c) $\Lambda_m^\circ = v_+ \lambda_+ + v_- \lambda_-$ (d) $\frac{\Lambda^\circ}{\Lambda_m} = \frac{1}{\Lambda_m^\circ} + \frac{\Lambda_m c}{K_a (\Lambda_m^\circ)^2}$
24. The variation of equivalent conductance of strong electrolyte with (concentration)^{1/2} is represented by


**RESPONSE
GRID**
15. (a)(b)(c)(d)
20. (a)(b)(c)(d)
16. (a)(b)(c)(d)
21. (a)(b)(c)(d)
17. (a)(b)(c)(d)
22. (a)(b)(c)(d)
18. (a)(b)(c)(d)
23. (a)(b)(c)(d)
19. (a)(b)(c)(d)
24. (a)(b)(c)(d)

Space for Rough Work

Mastering Chemistry With Ravi Arora

C-68

DPP/ CC17

25. A device that converts energy of combustion of fuels like hydrogen and methane, directly into electrical energy is known as :
- (a) Electrolytic cell (b) Dynamo
(c) Ni-Cd cell (d) Fuel Cell
26. In acidic medium MnO_2 is an oxidant as
- $$\text{MnO}_2(\text{s}) + 4\text{H}^+ + 2\text{e}^- \longrightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O}$$
- If the pH of solution is decreased by one unit, the electrode potential of the half cell Pt: MnO_2 , Mn^{2+} will change by
- (a) 0.236V (b) -0.236V
(c) -0.118V (d) 0.118V
27. Consider the following cell reaction:
- $$2\text{Fe}(\text{s}) + \text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) \rightarrow 2\text{Fe}^{2+}(\text{aq}) + 2\text{H}_2\text{O}(\text{l}); E^\circ = 1.67\text{V}$$
- At $[\text{Fe}^{2+}] = 10^{-3}\text{M}$, $p(\text{O}_2) = 0.1\text{atm}$ and $\text{pH} = 3$, the cell potential at 25°C is
- (a) 1.47V (b) 1.77V
(c) 1.87V (d) 1.57V
28. In a hydrogen-oxygen fuel cell, combustion of hydrogen occurs to
- (a) produce high purity water
(b) create potential difference between two electrodes
(c) generate heat
(d) remove adsorbed oxygen from electron surfaces
29. Consider the following relations for emf of an electrochemical cell:
- (i) $\text{emf of cell} = (\text{Oxidation potential of anode}) - (\text{Reduction potential of cathode})$
(ii) $\text{emf of cell} = (\text{Oxidation potential of anode}) + (\text{Reduction potential of cathode})$
(iii) $\text{emf of cell} = (\text{Reduction potential of anode}) + (\text{Reduction potential of cathode})$
(iv) $\text{emf of cell} = (\text{Oxidation potential of anode}) - (\text{Oxidation potential of cathode})$
- Which of the above relations are correct?
- (a) (ii) and (iv)
(b) (iii) and (i)
(c) (i) and (ii)
(d) (iii) and (iv)
30. A hydrogen gas electrode is made by dipping platinum wire in a solution of HCl of $\text{pH} = 10$ and by passing hydrogen gas around the platinum wire at one atm pressure. The oxidation potential of electrode would be ?
- (a) 0.59V (b) 0.118V
(c) 1.18V (d) 0.059V

**RESPONSE
GRID**

25. (a)(b)(c)(d) 26. (a)(b)(c)(d) 27. (a)(b)(c)(d) 28. (a)(b)(c)(d) 29. (a)(b)(c)(d)
30. (a)(b)(c)(d)

DAILY PRACTICE PROBLEM DPP CHAPTERWISE 17 - CHEMISTRY

Total Questions	30	Total Marks	120
Attempted		Correct	
Incorrect		Net Score	
Cut-off Score	37	Qualifying Score	53
Success Gap = Net Score – Qualifying Score			
Net Score = (Correct × 4) – (Incorrect × 1)			

Space for Rough Work