

# Mastering Physics With Physics Guruji

## DPP - Daily Practice Problems

### Chapter-wise Sheets

Date :  Start Time :  End Time :

# PHYSICS

# CP05

SYLLABUS : Work, Energy and Power

Max. Marks : 180    Marking Scheme : (+4) for correct & (-1) for incorrect answer    Time : 60 min.

**INSTRUCTIONS** : This Daily Practice Problem Sheet contains 45 MCQs. For each question only one option is correct. Darken the correct circle/ bubble in the Response Grid provided on each page.

- A spring of spring constant  $5 \times 10^3$  N/m is stretched initially by 5cm from the unstretched position. Then the work required to stretch it further by another 5 cm is  
(a) 12.50 Nm    (b) 18.75 Nm  
(c) 25.00 Nm    (d) 6.25 Nm
- A particle of mass 10 g moves along a circle of radius 6.4 cm with a constant tangential acceleration. What is the magnitude of this acceleration if the kinetic energy of the particle becomes equal to  $8 \times 10^{-4}$  J by the end of the second revolution after the beginning of the motion ?  
(a)  $0.1 \text{ m/s}^2$     (b)  $0.15 \text{ m/s}^2$     (c)  $0.18 \text{ m/s}^2$     (d)  $0.2 \text{ m/s}^2$
- A body is moved along a straight line by a machine delivering a constant power. The distance moved by the body in time 't' is proportional to  
(a)  $t^{3/4}$     (b)  $t^{3/2}$     (c)  $t^{1/4}$     (d)  $t^{1/2}$
- A ball is thrown vertically downwards from a height of 20 m with an initial velocity  $v_0$ . It collides with the ground and loses 50% of its energy in collision and rebounds to the same height. The initial velocity  $v_0$  is : (Take  $g = 10 \text{ ms}^{-2}$ )  
(a)  $20 \text{ ms}^{-1}$     (b)  $28 \text{ ms}^{-1}$   
(c)  $10 \text{ ms}^{-1}$     (d)  $14 \text{ ms}^{-1}$
- A cord is used to lower vertically a block of mass M, a distance d at a constant downward acceleration of  $g/4$ . The work done by the cord on the block is  
(a)  $Mg \frac{d}{4}$     (b)  $3Mg \frac{d}{4}$     (c)  $-3Mg \frac{d}{4}$     (d)  $Mg d$
- A rubber ball is dropped from a height of 5m on a plane, where the acceleration due to gravity is not shown. On bouncing it rises to 1.8 m. The ball loses its velocity on bouncing by a factor of  
(a)  $\frac{16}{25}$     (b)  $\frac{2}{5}$     (c)  $\frac{3}{5}$     (d)  $\frac{9}{25}$
- A ball of mass m moving with a constant velocity strikes against a ball of same mass at rest. If e = coefficient of restitution, then what will be the ratio of velocity of two balls after collision?  
(a)  $\frac{1-e}{1+e}$     (b)  $\frac{e-1}{e+1}$     (c)  $\frac{1+e}{1-e}$     (d)  $\frac{2+e}{e-1}$
- A particle of mass m is driven by a machine that delivers a constant power of k watts. If the particle starts from rest the force on the particle at time t is  
(a)  $\sqrt{mk} t^{-1/2}$     (b)  $\sqrt{2mk} t^{-1/2}$   
(c)  $\frac{1}{2}\sqrt{mk} t^{-1/2}$     (d)  $\sqrt{\frac{mk}{2}} t^{-1/2}$


RESPONSE  
GRID

1. (a)(b)(c)(d)    2. (a)(b)(c)(d)    3. (a)(b)(c)(d)    4. (a)(b)(c)(d)    5. (a)(b)(c)(d)  
6. (a)(b)(c)(d)    7. (a)(b)(c)(d)    8. (a)(b)(c)(d)

Space for Rough Work

# Mastering Physics With Physics Guruji


9. A body of mass 2 kg moving under a force has relation between displacement  $x$  and time  $t$  as  $x = \frac{t^3}{3}$  where  $x$  is in metre and  $t$  is in sec. The work done by the body in first two second will be  
 (a) 1.6 joule (b) 16 joule  
 (c) 160 joule (d) 1600 joule
10. A sphere of mass 8m collides elastically (in one dimension) with a block of mass 2m. If the initial energy of sphere is  $E$ . What is the final energy of sphere?  
 (a)  $0.8E$  (b)  $0.36E$   
 (c)  $0.08E$  (d)  $0.64E$
11. Two similar springs P and Q have spring constants  $K_P$  and  $K_Q$ , such that  $K_P > K_Q$ . They are stretched, first by the same amount (case a), then by the same force (case b). The work done by the springs  $W_P$  and  $W_Q$  are related as, in case (a) and case (b), respectively  
 (a)  $W_P = W_Q$ ;  $W_P = W_Q$  (b)  $W_P > W_Q$ ;  $W_Q > W_P$   
 (c)  $W_P < W_Q$ ;  $W_Q < W_P$  (d)  $W_P = W_Q$ ;  $W_P > W_Q$
12. In the figure, the variation of potential energy of a particle of mass  $m = 2$  kg is represented w.r.t. its  $x$ -coordinate. The particle moves under the effect of this conservative force along the  $x$ -axis.




If the particle is released at the origin then

- (a) it will move towards positive  $x$ -axis  
 (b) it will move towards negative  $x$ -axis  
 (c) it will remain stationary at the origin  
 (d) its subsequent motion cannot be decided due to lack of information
13. The potential energy of a certain spring when stretched through distance  $S$  is 10 joule. The amount of work done (in joule) that must be done on this spring to stretch it through an additional distance  $s$ , will be  
 (a) 20 (b) 10 (c) 30 (d) 40
14. A force applied by an engine of a train of mass  $2.05 \times 10^6$  kg changes its velocity from 5m/s to 25 m/s in 5 minutes. The power of the engine is  
 (a) 1.025 MW (b) 2.05 MW  
 (c) 5 MW (d) 6 MW

15. The relationship between the force  $F$  and position  $x$  of a body is as shown in figure. The work done in displacing the body from  $x = 1$  m to  $x = 5$  m will be



- (a) 30 J (b) 15 J (c) 25 J (d) 20 J
16. A body is allowed to fall freely under gravity from a height of 10m. If it loses 25% of its energy due to impact with the ground, then the maximum height it rises after one impact is  
 (a) 2.5m (b) 5.0m (c) 7.5m (d) 8.2m
17. A block C of mass  $m$  is moving with velocity  $v_0$  and collides elastically with block A of mass  $m$  and connected to another block B of mass  $2m$  through spring constant  $k$ . What is  $k$  if  $x_0$  is compression of spring when velocity of A and B is same?



- (a)  $\frac{mv_0^2}{x_0^2}$  (b)  $\frac{mv_0^2}{2x_0^2}$   
 (c)  $\frac{3mv_0^2}{2x_0^2}$  (d)  $\frac{2mv_0^2}{3x_0^2}$
18. Two springs of force constants 300 N/m (Spring A) and 400 N/m (Spring B) are joined together in series. The combination is compressed by 8.75 cm. The ratio of energy stored in A and B is  $\frac{E_A}{E_B}$ . Then  $\frac{E_A}{E_B}$  is equal to :  
 (a)  $\frac{4}{3}$  (b)  $\frac{16}{9}$  (c)  $\frac{3}{4}$  (d)  $\frac{9}{16}$
19. A body of mass 1 kg begins to move under the action of a time dependent force  $\vec{F} = (2t\hat{i} + 3t^2\hat{j})$  N, where  $\hat{i}$  and  $\hat{j}$  are unit vectors along  $x$  and  $y$  axis. What power will be developed by the force at the time  $t$ ?  
 (a)  $(2t^2 + 3t^3)W$  (b)  $(2t^2 + 4t^4)W$   
 (c)  $(2t^3 + 3t^4)W$  (d)  $(2t^3 + 3t^5)W$
20. A bullet of mass 20 g and moving with 600 m/s collides with a block of mass 4 kg hanging with the string. What is the velocity of bullet when it comes out of block, if block rises to height 0.2 m after collision?  
 (a) 200 m/s (b) 150 m/s (c) 400 m/s (d) 300 m/s

**RESPONSE GRID**

- |                  |                  |                  |                  |
|------------------|------------------|------------------|------------------|
| 9. (a)(b)(c)(d)  | 10. (a)(b)(c)(d) | 11. (a)(b)(c)(d) | 12. (a)(b)(c)(d) |
| 14. (a)(b)(c)(d) | 15. (a)(b)(c)(d) | 16. (a)(b)(c)(d) | 17. (a)(b)(c)(d) |
| 19. (a)(b)(c)(d) | 20. (a)(b)(c)(d) | 18. (a)(b)(c)(d) |                  |

Space for Rough Work

# Mastering Physics With Physics Guruji

21. A body of mass  $m$  kg is ascending on a smooth inclined plane of inclination  $\theta$  ( $\sin \theta = \frac{1}{x}$ ) with constant acceleration of a  $m/s^2$ . The final velocity of the body is  $v$  m/s. The work done by the body during this motion is (Initial velocity of the body = 0)

- (a)  $\frac{1}{2}mv^2(g + xa)$       (b)  $\frac{mv^2}{2}\left(\frac{g}{2} + a\right)$   
 (c)  $\frac{2mv^2x}{a}(a + gx)$       (d)  $\frac{mv^2}{2ax}(g + xa)$

22. A glass marble dropped from a certain height above the horizontal surface reaches the surface in time  $t$  and then continues to bounce up and down. The time in which the marble finally comes to rest is

- (a)  $e^{nt}$       (b)  $e^{2t}$       (c)  $t \left[ \frac{1+e}{1-e} \right]$       (d)  $t \left[ \frac{1-e}{1+e} \right]$

23. The potential energy of a 1 kg particle free to move along

the  $x$ -axis is given by  $V(x) = \left( \frac{x^4}{4} - \frac{x^2}{2} \right)$  J.

The total mechanical energy of the particle is 2 J. Then, the maximum speed (in m/s) is

- (a)  $\frac{3}{\sqrt{2}}$       (b)  $\sqrt{2}$       (c)  $\frac{1}{\sqrt{2}}$       (d) 2

24. Water falls from a height of 60 m at the rate of 15 kg/s to operate a turbine. The losses due to frictional force are 10% of energy. How much power is generated by the turbine? ( $g = 10 \text{ m/s}^2$ )

- (a) 8.1 kW      (b) 10.2 kW      (c) 12.3 kW      (d) 7.0 kW


24. A car of mass  $m$  starts from rest and accelerates so that the instantaneous power delivered to the car has a constant magnitude  $P_0$ . The instantaneous velocity of this car is proportional to:

- (a)  $t^2 P_0$       (b)  $t^{1/2}$       (c)  $t^{-1/2}$       (d)  $\frac{t}{\sqrt{m}}$

25. When a 1.0kg mass hangs attached to a spring of length 50 cm, the spring stretches by 2 cm. The mass is pulled down until the length of the spring becomes 60 cm. What is the amount of elastic energy stored in the spring in this condition. if  $g = 10 \text{ m/s}^2$ .

- (a) 1.5 joule      (b) 2.0 joule      (c) 2.5 joule      (d) 3.0 joule

26. A block of mass  $m$  rests on a rough horizontal surface (Coefficient of friction is  $\mu$ ). When a bullet of mass  $m/2$  strikes horizontally, and get embedded in it, the block moves a distance  $d$  before coming to rest. The initial velocity of the bullet is  $k\sqrt{2\mu gd}$ , then the value of  $k$  is



- (a) 2      (b) 3      (c) 4      (d) 5

27. A force acts on a 30 gm particle in such a way that the position of the particle as a function of time is given by  $x = 3t - 4t^2 + t^3$ , where  $x$  is in metres and  $t$  is in seconds. The work done during the first 4 seconds is

- (a) 576mJ      (b) 450mJ      (c) 490mJ      (d) 530mJ

28. A particle of mass  $m_1$  moving with velocity  $v$  strikes with a mass  $m_2$  at rest, then the condition for maximum transfer of kinetic energy is

- (a)  $m_1 \gg m_2$       (b)  $m_2 \gg m_1$       (c)  $m_1 = m_2$       (d)  $m_1 = 2m_2$

29. A mass  $m$  is moving with velocity  $v$  collides inelastically with a bob of simple pendulum of mass  $m$  and gets embedded into it. The total height to which the masses will rise after collision is

- (a)  $\frac{v^2}{8g}$       (b)  $\frac{v^2}{4g}$       (c)  $\frac{v^2}{2g}$       (d)  $\frac{2v^2}{g}$

30. A 10 H.P. motor pumps out water from a well of depth 20 m and fills a water tank of volume 22380 litres at a height of 10 m from the ground. The running time of the motor to fill the empty water tank is ( $g = 10 \text{ ms}^{-2}$ )

- (a) 5 minutes      (b) 10 minutes  
 (c) 15 minutes      (d) 20 minutes

31. A particle of mass  $m_1$  is moving with a velocity  $v_1$  and another particle of mass  $m_2$  is moving with a velocity  $v_2$ . Both of them have the same momentum but their different kinetic energies are  $E_1$  and  $E_2$  respectively. If  $m_1 > m_2$  then

- (a)  $E_1 = E_2$       (b)  $E_1 < E_2$       (c)  $\frac{E_1}{E_2} = \frac{m_1}{m_2}$       (d)  $E_1 > E_2$

32. A block of mass 10 kg, moving in  $x$  direction with a constant speed of  $10 \text{ ms}^{-1}$ , is subject to a retarding force  $F = 0.1 \times J$  m during its travel from  $x = 20$  m to 30 m. Its final KE will be:

- (a) 450J      (b) 275J      (c) 250J      (d) 475J

33. Identify the false statement from the following

- (a) Work-energy theorem is not independent of Newton's second law.  
 (b) Work-energy theorem holds in all inertial frames.  
 (c) Work done by friction over a closed path is zero.  
 (d) No potential energy can be associated with friction.

34. A one-ton car moves with a constant velocity of  $15 \text{ ms}^{-1}$  on a rough horizontal road. The total resistance to the motion of the car is 12% of the weight of the car. The power required to keep the car moving with the same constant velocity of  $15 \text{ ms}^{-1}$  is [Take  $g = 10 \text{ ms}^{-2}$ ]

- (a) 9 kW      (b) 18 kW      (c) 24 kW      (d) 36 kW

35. A ball is released from the top of a tower. The ratio of work done by force of gravity in first, second and third second of the motion of the ball is

- (a) 1:2:3      (b) 1:4:9      (c) 1:3:5      (d) 1:5:3

**RESPONSE  
GRID**

21. (a)(b)(c)(d)	22. (a)(b)(c)(d)	23. (a)(b)(c)(d)	24. (a)(b)(c)(d)	25. (a)(b)(c)(d)
26. (a)(b)(c)(d)	27. (a)(b)(c)(d)	28. (a)(b)(c)(d)	29. (a)(b)(c)(d)	30. (a)(b)(c)(d)
31. (a)(b)(c)(d)	32. (a)(b)(c)(d)	33. (a)(b)(c)(d)	34. (a)(b)(c)(d)	35. (a)(b)(c)(d)

Space for Rough Work

# Mastering Physics With Physics Guruji

36. Two spheres A and B of masses  $m_1$  and  $m_2$  respectively collide. A is at rest initially and B is moving with velocity  $v$  along x-axis. After collision B has a velocity  $\frac{v}{2}$  in a direction perpendicular to the original direction. The mass A moves after collision in the direction.
- Same as that of B
  - Opposite to that of B
  - $\theta = \tan^{-1}(1/2)$  to the x-axis
  - $\theta = \tan^{-1}(-1/2)$  to the x-axis
37. A 2 kg block slides on a horizontal floor with a speed of 4m/s. It strikes an uncompressed spring, and compresses it till the block is motionless. The kinetic friction force is 15N and spring constant is 10,000 N/m. The spring compresses by
- 8.5 cm
  - 5.5 cm
  - 2.5 cm
  - 11.0 cm
38. An engine pumps water through a hose pipe. Water passes through the pipe and leaves it with a velocity of 2 m/s. The mass per unit length of water in the pipe is 100 kg/m. What is the power of the engine?
- 400 W
  - 200 W
  - 100 W
  - 800 W
39. A uniform chain of length 2 m is kept on a table such that a length of 60 cm hangs freely from the edge of the table. The total mass of the chain is 4 kg. What is the work done in pulling the entire chain on the table?
- 12 J
  - 3.6 J
  - 7.2 J
  - 1200 J
40. A mass ' $m$ ' moves with a velocity ' $v$ ' and collides inelastically with another identical mass. After collision the 1<sup>st</sup> mass moves

with velocity  $\frac{v}{\sqrt{3}}$  in a direction perpendicular to the initial direction of motion. Find the speed of the 2<sup>nd</sup> mass after collision.




- $\sqrt{3}v$
  - $v$
  - $\frac{v}{\sqrt{3}}$
  - $\frac{2}{\sqrt{3}}v$
41. A spherical ball of mass 20 kg is stationary at the top of a hill of height 100 m. It rolls down a smooth surface to the ground, then climbs up another hill of height 30 m and finally rolls down to a horizontal base at a height of 20 m above the ground. The velocity attained by the ball is
- 20 m/s
  - 40 m/s
  - $10\sqrt{30}$  m/s
  - 10 m/s

42. A block of mass M is kept on a platform which is accelerated upward with a constant acceleration 'a' during the time interval T. The work done by normal reaction between the block and platform is



- $-\frac{MgaT^2}{2}$
- $\frac{1}{2}M(g+a)aT^2$
- $\frac{1}{2}Ma^2T$
- Zero

43. A spring lies along an x axis attached to a wall at one end and a block at the other end. The block rests on a frictionless surface at  $x = 0$ . A force of constant magnitude F is applied to the block that begins to compress the spring, until the block comes to a maximum displacement  $x_{max}$ .



During the displacement, which of the curves shown in the graph best represents the kinetic energy of the block?

- 1
  - 2
  - 3
  - 4
44. The K.E. acquired by a mass m in travelling a certain distance d, starting from rest, under the action of a constant force is directly proportional to
- m
  - $\sqrt{m}$
  - $\frac{1}{\sqrt{m}}$
  - independent of m
45. A vertical spring with force constant k is fixed on a table. A ball of mass m at a height h above the free upper end of the spring falls vertically on the spring so that the spring is compressed by a distance d. The net work done in the process is
- $mg(h+d) - \frac{1}{2}kd^2$
  - $mg(h-d) - \frac{1}{2}kd^2$
  - $mg(h-d) + \frac{1}{2}kd^2$
  - $mg(h+d) + \frac{1}{2}kd^2$

<b>RESPONSE GRID</b>	36. (a)(b)(c)(d)	37. (a)(b)(c)(d)	38. (a)(b)(c)(d)	39. (a)(b)(c)(d)	40. (a)(b)(c)(d)
	41. (a)(b)(c)(d)	42. (a)(b)(c)(d)	43. (a)(b)(c)(d)	44. (a)(b)(c)(d)	45. (a)(b)(c)(d)

DAILY PRACTICE PROBLEM DPP CHAPTERWISE CP05 - PHYSICS			
Total Questions	45	Total Marks	180
Attempted		Correct	
Incorrect		Net Score	
Cut-off Score	50	Qualifying Score	70
Success Gap = Net Score – Qualifying Score			
Net Score = (Correct × 4) – (Incorrect × 1)			

Space for Rough Work