

Recommendations for Preventive Pediatric Health Care

Bright Futures/American Academy of Pediatrics

Each child and family is unique; therefore, these **Recommendations for Preventive Pediatric Health Care** are designed for the care of children who are receiving competent parenting, have no manifestations of any important health problems, and are growing and developing in satisfactory fashion. **Additional visits may become necessary** if circumstances suggest variations from normal.

Developmental, psychosocial, and chronic disease issues for children and adolescents may require frequent counseling and treatment visits separate from preventive care visits.

These guidelines represent a consensus by the American Academy of Pediatrics (AAP) and Bright Futures. The AAP continues to emphasize the great importance of **continuity of care** in comprehensive health supervision and the need to avoid **fragmentation of care**.

The recommendations in this statement do not indicate an exclusive course of treatment or standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

Copyright © 2008 by the American Academy of Pediatrics.

No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.

AGE	INFANCY									EARLY CHILDHOOD						MIDDLE CHILDHOOD						ADOLESCENCE											
	PRENATAL ¹	NEWBORN ²	3-5 d ³	By 1 mo	2 mo	4 mo	6 mo	9 mo	12 m	15 mo	18 mo	24 mo	30 mo	3 y	4 y	5 y	6 y	7 y	8 y	9 y	10 y	11 y	12 y	13 y	14 y	15 y	16 y	17 y	18 y	19 y	20 y	21 y	
HISTORY																																	
Initial/Interval	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
MEASUREMENTS																																	
Length/Height and Weight		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Head Circumference		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Weight for Length		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	
Body Mass Index																																	
Blood Pressure ⁴		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★		
SENSORY SCREENING																																	
Vision		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★		
Hearing		●	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★		
DEVELOPMENTAL/BEHAVIORAL ASSESSMENT																																	
Developmental Screening ⁵									●			●																					
Autism Screening ⁶																																	
Developmental Surveillance ⁷		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		
Psychosocial/Behavioral Assessment		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		
Alcohol and Drug Use Assessment																							★	★	★	★	★	★	★	★	★		
PHYSICAL EXAMINATION⁸		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		
PROCEDURES¹																																	
Newborn Metabolic/Hemoglobin Screening ²		←	●	→																													
Immunization ³		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●		
Hematocrit or Hemoglobin ⁴						★					★	★		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★			
Lead Screening ⁵							★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★		
Tuberculin Test ⁶				★			★		★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★			
Dyslipidemia Screening ⁷												★										★	★	★	★	★	★	★	★	★			
STI Screening ⁸																						★	★	★	★	★	★	★	★	★			
Cervical Dysplasia Screening ⁹																						★	★	★	★	★	★	★	★	★			
ORAL HEALTH¹⁰								★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★	★			
ANTICIPATORY GUIDANCE¹¹	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●			

- If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
- A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding per AAP statement "The Prenatal Visit" (2007) [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;107/9/1456>].
- Every infant should have a newborn evaluation after birth, breastfeeding encouraged, and instruction and support offered.
- Every infant should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital to include evaluation for feeding and jaundice. Breastfeeding infants should receive formal breastfeeding evaluation, encouragement, and instruction as recommended in AAP statement "Breastfeeding and the Use of Human Milk" (2005) [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;115/2/398>]. For newborns discharged in less than 48 hours after delivery, the infant must be examined within 48 hours of discharge per AAP statement "Hospital Stay for Healthy Term Newborns" (2004) [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;113/3/434>].
- Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
- If the patient is uncooperative, reexamine within 6 months per the AAP statement "Eye Examination in Infants, Children, and Young Adults by Pediatricians" (2007) [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;111/4/902>].
- All newborns should be screened per AAP statement "New 2000 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs" (2000) [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;106/4/798>]. Joint Committee on Infant Hearing, Year 2007 position statement: principles and guidelines for early hearing detection and intervention programs. *Pediatrics*. 2007;120:989-997.

- AAP Council on Children With Disabilities, AAP Section on Developmental Behavioral Pediatrics, AAP Bright Futures Steering Committee, AAP Medical Home Initiatives for Children With Special Needs Project Advisory Committee. Identifying infants and young children with developmental disabilities in the medical home: an algorithm for developmental surveillance and screening. *Pediatrics*. 2006;118:405-420 [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;118/4/405>].
- Gupta VB, Hyman SL, Johnson CP et al. Identifying children with autism early? *Pediatrics*. 2007;119:152-153 [URL: <http://pediatrics.aappublications.org/cgi/content/full/119/1/152>].
- At each visit, age-appropriate physical examination is essential, with infant totally undressed, older child undressed and suitably draped.
- These may be modified, depending on entry point into schedule and individual need.
- Newborn metabolic and hemoglobinopathy screening should be done according to state law. Results should be reviewed at visits and appropriate monitoring or referral done as needed.
- Schedule per the Committee on Infectious Diseases, published annually in the January issue of *Pediatrics*. Every visit should be an opportunity to update and complete a child's immunizations.
- See AAP Pediatric Nutrition Handbook, 5th Edition (2003) for a discussion of universal and selective screening options. See also Recommendations to prevent and control iron deficiency in the United States, *MMWR*, 1998;47(RR-3):1-36.
- For children at risk of lead exposure, consult the AAP statement "Lead Exposure in Children: Prevention, Detection, and Management" (2006) [URL: <http://aappolicy.aappublications.org/cgi/content/full/pediatrics;118/4/1038>]. Additionally, screening should be done in accordance with state law where applicable.

- Perform risk assessments or screens as appropriate, based on universal screening requirements for patients with Medicaid or high prevalence areas.
- Tuberculosis testing per recommendations of the Committee on Infectious Diseases, published in the current edition of *Red Book: Report of the Committee on Infectious Diseases*. Testing should be done on recognition of high-risk factors.
- "Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report" (2002) [URL: <http://circ.ahajournals.org/cgi/content/full/106/25/3143>] and "The Expert Committee Recommendations on the Assessment, Prevention, and Treatment of Child and Adolescent Overweight and Obesity." Supplement to *Pediatrics*. In press.
- All sexually active patients should be screened for sexually transmitted infections (STIs).
- All sexually active girls should have screening for cervical dysplasia as part of a pelvic examination beginning within 3 years of onset of sexual activity or age 21 (whichever comes first).
- Referral to dental home, if available. Otherwise, administer oral health risk assessment. If the primary water source is deficient in fluoride, consider oral fluoride supplementation.
- At the visits for 3 years and 6 years of age, it should be determined whether the patient has a dental home. If the patient does not have a dental home, a referral should be made to one. If the primary water source is deficient in fluoride, consider oral fluoride supplementation.
- Refer to the specific guidance by age as listed in Bright Futures Guidelines, Hagan JJ, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008.

KEY
● = to be performed ★ = risk assessment to be performed, with appropriate action to follow, if positive ← ● → = range during which a service may be provided, with the symbol indicating the preferred age