

# Croatia Norval S.C.

## **Concussion Return to Play Protocol**

| Policy Type | <u>Author</u> | Approved By | Approval Date |
|-------------|---------------|-------------|---------------|
| Tech/Admin. | A. Radosevic  | CNSC B.O.D  | Dec 1, 2022   |





#### What is a concussion?

A concussion is a neurologic injury that causes a temporary disturbance in brain cells. It is the result of extreme acceleration or deceleration of the brain within the skull. The common misconception is that you must be hit in the head to cause a concussion injury – this is not true! Because a concussion is the result of acceleration or deceleration of the brain, a concussion can happen with a significant blow to anywhere on the body, provided sufficient acceleration/deceleration is transmitted to the brain tissue, head or face. There may be no other visible signs of a brain injury. It can cause changes in the brain that may not be seen in a CT scan or X-Ray. Balance, reaction time and cognitive skills can also be impacted.

#### What are the signs and symptoms of a concussion?

Symptoms of a concussion range from mild to severe and can last for or appear within hours, days, weeks, or months. In some cases, there may be no symptoms at all. If you suspect a concussion or notice any of the symptoms listed below, contact your doctor immediately. In severe cases, proceed to an Urgent Care Centre or Emergency Room immediately.

#### Visible Symptoms:

| Loss of consciousness or responsiveness | Loss of motion/slow to rise   |
|---|-------------------------------|
| Unsteady on feet/balance issues         | Grabbing/Clutching the head   |
| Dazed, vacant or blank look             | Vomiting                      |
| Disoriented                             | Bleeding from the nose/ears   |
| Sensitive to light                      | Unequal pupils (eye)          |
| Seizures or convulsions                 | Deteriorating conscious state |

#### Non-Visible Symptoms

| Confusion/disorientation               | Complaints of pain/headache                                      |
|--|--|
| Complaints of dizziness/nausea         | Complaints of fuzzy/blurred vision                               |
| Inability to concentrate               | Easily upset/angered (emotional)                                 |
| Nervous/Anxious                        | Complaints of tingling/burning feeling in arms & legs            |
| Tired/sleeping more or less than usual | Memory loss-unable to remember events leading up to the incident |
| Trouble falling asleep                 | Deteriorating conscious state                                    |





\*\*Please note that complaints of neck pain should be addressed immediately as a head injury may also result in a neck / spinal injury\*\*

#### **CNSC Emergency Protocol – Suspected Player Concussion**

A player showing any signs/symptoms as listed on the *Pocket Concussion Recognition Tool* or as listed in this document must be removed from play immediately. All Coaches/Trainers are required to be familiar with the use of this tool and follow the Club's protocol without question. The player is not to re-enter the game or practice: <a href="https://sirc.ca/wp-content/uploads/2019/10/crt5\_pdf2.pdf">https://sirc.ca/wp-content/uploads/2019/10/crt5\_pdf2.pdf</a>

- In the event of a loss of consciousness call 911 for assistance immediately. Do not move the player unless necessary for safety reasons.
- > Ensure that a member of the team staff stays with the player.
- > Notify any family members immediately if they are present.
- The player must be evaluated by a physician before returning to play. Documentation from the physician is required. The average recovery period is 10 to 14 days but can sometimes be longer.
- A graduated <u>"Return to Play Process"</u> using Ontario Soccer's Return to Play policy that can be found at: <u>https://cdn2.sportngin.com/attachments/document/2cec-1895166/Return-to-</u> <u>Sport\_Protocol\_Tracking\_-\_Soccer\_2019.pdf#\_ga=2.29790207.110022128.1643121993-</u> <u>1567226926.1620322605</u>
- Coaches must complete the required injury report and submit it to CNSC within 24 hours of the incident, see the CNSA website for download.





|  | EXERCISE ALLOWED   | % MAX HEART RATE | DURATION             | OBJECTIVE   |
|--|--|------------------|----------------------|---|
| REST AND<br>RECOVER                    | <ul> <li>None</li> <li>"Rest the body, Rest the brain"</li> </ul>  | No training      | Until symptoms clear | <ul> <li>Recovery</li> <li>Symptom free</li> </ul>  |
| STAGE 1<br>LIGHT EXERCISE              | <ul> <li>Walking, light jogging, swimming, stationary cycling or equivalent</li> <li>No football, resistance training, weight lifting, jumping or hard running</li> </ul>  | < 70%            | < 15 min             | Increase heart rate   |
| STAGE 2<br>SOCCER-SPECIFIC<br>EXERCISE | <ul> <li>Simple movement activities ie. running drills</li> <li>Limit body and head movement</li> <li>NO head impact activities</li> <li>NO heading</li> </ul>   | < 80%            | < 45 min             | Add movement  |
| STAGE 3<br>NON-CONTACT<br>TRAINING     | <ul> <li>Progression to more complex training activities</li> <li>with increased intensity,</li> <li>coordination and attention e.g. passing, change of direction, shooting, small-sided game</li> <li>May start resistance training</li> <li>NO head impact activities including NO heading</li> <li>goalkeeping activities should avoid diving and any risk of the head being hit by a ball</li> </ul> | < 90%            | < 60 min             | <ul> <li>Exercise,<br/>coordination and<br/>skills/tactics</li> </ul>                             |
| STAGE 4<br>FULL CONTACT<br>PRACTICE    | » Normal training activities ie tackling, heading diving saves   |                  |                      | <ul> <li>Restore confidence<br/>and assess<br/>functional skills by<br/>coaching staff</li> </ul> |
| STAGE 5<br>GAME PLAY                   | » Player rehabilitated   |                  |                      | <ul> <li>Return to game<br/>play</li> </ul>   |

#### Ontario Soccer 4 Stage Remove from Sport Protocol

A suspected concussion has been identified and the player is removed from play. Head Coaches hold the final decision to remove players with a suspected concussion, which can be found here:

https://cdn2.sportngin.com/attachments/document/787f-1895165/Remove-from-Sport\_Protocol\_Summary\_-\_Soccer\_2019.pdf#\_ga=2.214299668.2090528831.1647219489-334875778.1647022291

#### **Ontario Soccer Suspected Concussion report**

A suspected concussion must be filled out when a coach suspects a player has suffered a concussion & the player is removed from play. Head Coaches hold the final decision to remove players with a suspected concussion, which can be found here: <u>https://cdn3.sportngin.com/attachments/document/f9af-</u> 1895168/Suspected Concussion Report Form -

Soccer\_2019.pdf#\_ga=2.171302051.2090528831.1647219489-334875778.1647022291

#### Ontario Soccer Return to Play 6 Stages Protocol

This is a multi-part process that requires supervision from the player's healthcare provider at each stage.

Ontario Soccer's 6-stage Return to Play protocol can be found by clicking here:

https://cdn2.sportngin.com/attachments/document/2cec-1895166/Return-to-Sport\_Protocol\_Tracking\_-

<u>Soccer\_2019.pdf#\_ga=2.29790207.110022128.1643121993-1567226926.1620322605</u>





### 6 Stage Protocol

| 1) Limit activity              | Physical & cognitive rest        |
|--------------------------------|----------------------------------|
| 2) Light aerobic exercise      | Walking, swimming, cycling       |
| 3) Sport-specific exercise     | Light running skills, no heading |
| 4) Non-contact training drills | Passing/Movement                 |
| 6) Full contact practice       | Normal training cycle            |
| 7) Return to Play              | Normal Game Play                 |

#### \*\*A minimum of 24 hours is required between each stage.\*\*

Any recurrence of symptoms requires the player to return to the previous level. CNSC Coaches, Assistant Coaches, Managers & Trainers are not authorized to oversee Return to Play processes.

#### **Base Line Testing**

Many soccer clubs recommend that players involved in competitive programs undergo Base Line Testing before the beginning of the outdoor season. This should be done in a controlled environment under the supervision of an approved concussion testing facility.

#### Caution

CNSC is committed to providing a safe environment for all players regardless of age and gender. It is, however, the responsibility of the parent to ensure that appropriate professional care is provided to players under the age of 18 in the event of injury. In the event of a concussion diagnosis, premature return to play can result in serious and sometimes permanent injury, particularly in the event of a second impact. Parents should seek ongoing medical evaluation for the benefit of the player's long-term health and safety and find an approved concussion testing facility.

#### More Information can be found at the following:

https://www.ontariosoccer.net/concussion-resources

#### If you suspect an athlete or your child suffered a concussion, contact your healthcare provider or:

https://hollandbloorview.ca/services/programs-services/concussion-centre/concussion-

#### services/clinical-services

| <u>Rev. #</u> | Date        | <u>Revisor</u> | <u>Comments</u> |
|---------------|-------------|----------------|-----------------|
| .001          | Dec 1, 2022 | A. Radosevic   | New Document    |