

**Screening for Lifetime History of TBI (from the OSU TBI-ID)  
and other Acquired Brain Injuries**

1. Please think about injuries you have had during your entire lifetime, especially those that affected your head or neck. It might help to remember times you went to the hospital or Emergency room. Think about injuries you may have received from a car or motorcycle wreck, bicycle crash, being hit by something, falling down, being hit by someone, playing sports or an injury during military service.
  - a. Thinking about any injuries you have had in your lifetime, were you ever knocked out or did you lose consciousness?  
 Yes  
 No (IF NO, GO TO QUESTION 2)
  - b. What was the longest time you were knocked out or unconscious? (Choose just one; if you are not sure please make your best guess.)  
 knocked out or lost consciousness for *less than 30 min*  
 knocked out or lost consciousness *between 30 min and 24 hours*  
 knocked out or lost consciousness for *24 hours or longer*
  - c. How old were you the first time you were knocked out or lost consciousness?  
\_\_\_\_\_ years old
2. Have you ever lost consciousness from a drug overdose or being strangled or choked?  
 Yes  
 No (IF NO, GO TO QUESTION 3)
  - a. How many times from a drug overdose? \_\_\_\_\_ # overdose
  - b. How many time from being strangled or choked? \_\_\_\_\_ # choked
3. Have you EVER been told by a doctor or other health professional that you had any of the following?  
 epilepsy or seizures?  
 a stroke or a transient ischemic attack?  
 cerebral palsy?  
 brain cancer?  
 a brain infection like meningitis or encephalitis?  
 toxic exposure, like to lead or pesticides?  
 dementia, like Alzheimer's Disease?  
 a progressive disease like AIDS, multiple sclerosis, Parkinson's Disease or Huntington's Disease? (if yes, which one \_\_\_\_\_)

**NOTES:**

Question 1 allows 3 indicators of lifetime history of TBI to be computed:

1. Positive for a lifetime history for TBI with loss of consciousness (yes/no)
2. Worst TBI with loss of consciousness (LOC) was mild, moderate or severe (no TBI with LOC, mild TBI with LOC, moderate TBI, severe TBI)
3. Age at first TBI with loss of consciousness (in years)