## performance evaluation two

## INFANT - ONE-PROVIDER CPR



Student Name Class Date

You are a trained CPR AED provider responding to repeated, panic-stricken shouts of "Help!" As you reach the scene, an obviously distraught teenager hands you a limp infant, pleading, "Help me...she's not breathing!" You have appropriate PPE. There is no AED nearby. You have an infant CPR mask with a one-way valve with your first aid kit. Demonstrate what actions you would take next.

Provider Action (Performance Criteria)	Instructor Prompt	Check Off
<ul> <li>Assesses scene safety.</li> <li>Takes Standard Precautions.</li> <li>Assesses responsiveness.</li> <li>Activates EMS and/or EAP.</li> <li>After activating, and unless they are readily available, sends someone to get the first aid kit and an AED.</li> </ul>	"Scene is safe." "Infant is unresponsive." "EMS has been activated.	
<ul> <li>Assesses breathing for no more than 10 seconds.</li> </ul>	"The infant is not breathing."	
<ul> <li>Immediately starts CPR, beginning with chest compressions.</li> <li>Performs 30 high-quality chest compressions.</li> <li>Correctly uses one of three hand-position techniques: 2-Finger, 2-Thumb Encircling-Hands, or Heel of One Hand.</li> <li>Does not press the tip of the breastbone.</li> <li>Compresses at least 1/3 depth of chest or about 1½ inches (4 cm).</li> <li>Compresses at a rate of 100–120 times per minute (30 compressions in no less than 15 and no more than 18 seconds).</li> <li>Allows chest to fully recoil at the top of compressions.</li> </ul>		
<ul> <li>Seals a CPR mask against the infant's face.</li> <li>Opens the airway using the head-tilt, chin-lift maneuver.</li> <li>Gives two rescue breaths.</li> <li>Ensures each breath is 1 second in length. Creates visible rise of chest, but no more.</li> <li>Immediately resumes high-quality chest compressions in less than 10 seconds.</li> </ul>		
<ul> <li>Repeats a second CPR cycle of 30:2 compressions-to-breaths.</li> <li>Completes a third cycle of 30:2 compressions-to-breaths.</li> </ul>		
	<ul> <li>Assesses scene safety.</li> <li>Takes Standard Precautions.</li> <li>Assesses responsiveness.</li> <li>Activates EMS and/or EAP.</li> <li>After activating, and unless they are readily available, sends someone to get the first aid kit and an AED.</li> <li>Assesses breathing for no more than 10 seconds.</li> <li>Immediately starts CPR, beginning with chest compressions.</li> <li>Performs 30 high-quality chest compressions.</li> <li>Correctly uses one of three hand-position techniques: 2-Finger, 2-Thumb Encircling-Hands, or Heel of One Hand.</li> <li>Does not press the tip of the breastbone.</li> <li>Compresses at least 1/3 depth of chest or about 1½ inches (4 cm).</li> <li>Compresses at a rate of 100–120 times per minute (30 compressions in no less than 15 and no more than 18 seconds).</li> <li>Allows chest to fully recoil at the top of compressions.</li> <li>Seals a CPR mask against the infant's face.</li> <li>Opens the airway using the head-tilt, chin-lift maneuver.</li> <li>Gives two rescue breaths.</li> <li>Ensures each breath is 1 second in length. Creates visible rise of chest, but no more.</li> <li>Immediately resumes high-quality chest compressions in less than 10 seconds.</li> <li>Repeats a second CPR cycle of 30:2 compressions-to-breaths.</li> <li>Completes a third cycle of 30:2</li> </ul>	Performance Criteria)  Assesses scene safety. Takes Standard Precautions. Assesses responsiveness. Activates EMS and/or EAP. After activating, and unless they are readily available, sends someone to get the first aid kit and an AED.  Assesses breathing for no more than 10 seconds.  Immediately starts CPR, beginning with chest compressions. Correctly uses one of three hand-position techniques: 2-Finger, 2-Thumb Encircling-Hands, or Heel of One Hand. Compresses at least 1/3 depth of chest or about 1½ inches (4 cm). Compresses at a rate of 100–120 times per minute (30 compressions in no less than 15 and no more than 18 seconds). Allows chest to fully recoil at the top of compressions.  Seals a CPR mask against the infant's face. Opens the airway using the head-tilt, chin-lift maneuver. Giives two rescue breaths. Ensures each breath is 1 second in length. Creates visible rise of chest, but no more. Immediately resumes high-quality chest compressions in less than 10 seconds.  Repeats a second CPR cycle of 30:2 compressions-to-breaths. Completes a third cycle of 30:2

Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Check Off
Removes PPE	<ul> <li>Correctly removes and disposes of gloves.</li> <li>Verbalizes washing hands immediately after removing gloves.</li> </ul>	"EMS arrives and takes over. Please demonstrate how to remove your gloves and what actions you would take next."	

## **END PERFORMANCE EVALUATION**

Successfully Completed.	Not Successfully Completed. Remediation Required.	
Signature of HSI Authorized Ins	structor	HSI Instructor Registry Number: