performance evaluation five

ADULT FIRST AID - SEVERE ALLERGIC REACTION



Student Name	Class Date

You are a trained first aid provider responding to a call for help received over your workplace mobile device for a team member with trouble breathing. As you reach the scene, you see a person sitting at a break table. You have appropriate PPE and a first aid kit. Demonstrate what actions you would take next.

	Burth Arthur		
Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Check Off
Performs Assessment	 Assesses scene safety. Takes (or verbalizes taking) standard precautions. Assesses responsiveness. Activates EMS and/or EAP. After activating, and unless they are readily available, sends someone to get the first aid kit and an AED. 	"The scene is safe." "Person is responsive." "EMS/EAP activated." "Other first aid providers are on the way, bringing an AED and an additional first aid kit."	
Assesses Breathing	 Assesses breathing for no more than 10 seconds. 	"Person is responsive and breathing."	
Obtains Consent	► Introduces self and asks, "May I help you?"	"Person consents."	
Assesses for Life- Threatening Conditions	 Quickly scans the person for life-threatening conditions. 	"The person is in obvious distress - trembling, wheezing, and coughing. You see swelling of their lips, eyelids, and face. With difficulty the person says, "Nut allergiesatepretzel bite had peanut butter inside."	
Immediately Provides Appropriate First Aid	 Allows the person to find the most comfortable position in which to breathe. 	"Person wants to sit up."	
Performs a Secondary Assessment	► Looks for medical identification jewelry.	"Person is wearing a Medical Alert Bracelet on wrist that states, 'Nut Allergy - Use EpiPen®.'" [Instructor Note: Hand student epinephrine autoinjector trainer. Say "Here's their EpiPen®. They forgot how to use it."	

Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Check Off
Correctly Uses Epinephrine Autoinjector Trainer	 Grasps epinephrine autoinjector trainer in fist with the orange tip pointing downward. Removes blue safety release. Positions trainer near middle of person's outer thigh. Swings and firmly pushes orange tip at a 90-degree angle against thigh until it clicks. Holds device firmly on thigh for 3 seconds. Removes the trainer from the thigh and massages the injection area for 10 seconds. 	[Instructor Note: Use instructions at the left are for the correct use of an EpiPen® epinephrine autoinjector trainer. If using a different manufacturer's training device, students should follow the manufacturer's instructions for use].	
	END PERFORMANCE EVALUATION		

Successfully Completed.	Not Successfully Con	mpleted. Remediation Required.	
Signature of HSI Authorized Ins	structor	HSI Instructor Registry Number:	