

ADHIKAANSH ACADEMY (IITJEE NEET IX X XI XII)

RUN BY:

DEEPAK SAINI SIR

B.TECH, M.TECH (N.S.I.T. DELHI UNIVERSITY)

Ex. Faculty of

Resonance Kota, Career Point Kota

Aakash Institute Mumbai

PHYSICS NOTES (CLASS 12TH)



**FOR MORE FREE QUESTION
BANKS AND SAMPLE PAPER
WT'S UP ON : 7665186856**

A Best Faculty Group in Meerut....

ADHIKAANSH ACADEMY
IITJEE NEET IX X XI XII

सही समय और सही दिशा में कड़ी मेहनत, तो सफलता पक्की!

*So why
to wait...*



DIRECTOR
DEEPAK SAINI (DSA SIR)
B.TECH, M.TECH
NSIT DELHI UNIVERSITY

**और Class 11th से ही शुरू करें,
Board के साथ-साथ,
NEET और IITJEE की तैयारी**

Special Classes For Students Studying in std. 9th & 10th.

EX. FACULTY OF RESONANCE KOTA, AAKASH INSTITUTE MUMBAI
225/5, 1ST FLOOR, PANCHSHEEL COLONY, BEHIND PINNACLE TOWER, GARH ROAD, MEERUT. ☎ 8057870069
www.adhikanshiitjeemedical.com | adhikanshiitjeemedical@gmail.com

**Take Dummy Admission
in Class 11th & Prepare For..**

IITJEE & NEET

**For Free Counselling &
Career Guidance contact on:**

7665186856

CBSE Class-12 Physics Quick Revision Notes
Chapter-04: Moving Charges and Magnetism

- **Force on a Straight Conductor:**

Force F on a straight conductor of length l and carrying a steady current I placed in a uniform external magnetic field B ,

$$\vec{F} = I\vec{l} \times \vec{B}$$

- **Lorentz Force:**

Force on a charge q moving with velocity v in the presence of magnetic and electric fields B and E .

$$\vec{F} = q(\vec{v} \times \vec{B} + \vec{E})$$

- **Magnetic Force:**

The magnetic force $\vec{F}_B = q(\vec{v} \times \vec{B})$ is normal to \vec{V} and work done by it is zero.

- **Cyclotron:**

A charge q executes a circular orbit in a plane normal with frequency called the cyclotron frequency given by,

$$v_c = \frac{qB}{2\pi m}$$

This cyclotron frequency is independent of the particle's speed and radius.

- **Biot – Savart Law:**

It asserts that the magnetic field $d\vec{B}$ due to an element $d\vec{l}$ carrying a steady current I at a point P at a distance r from the current element is,

$$d\vec{B} = \frac{\mu_0}{4\pi} I \frac{d\vec{l} \times \vec{r}}{r^3}$$

- **Magnetic Field due to Circular Coil:**

Magnetic field due to circular coil of radius R carrying a current I at an axial distance X from the centre is

$$B = \frac{\mu_0 IR^2}{2(X^2 + R^2)^{3/2}}$$

At the centre of the coil,

$$B = \frac{\mu_0 I}{2R}$$

- **Ampere's Circuital Law:**

For an open surface S bounded by a loop C , then the Ampere's law states that

$$\oint_C \vec{B} \cdot d\vec{l} = \mu_0 I$$

where I refers to the current passing through S .

- If B is directed along the tangent to every point on the perimeter then

$$BL = \mu_0 I_e$$

Where I_e is the net current enclosed by the closed circuit.

- **Magnetic Field:**

Magnetic field at a distance R from a long, straight wire carrying a current I is given by,

$$B = \frac{\mu_0 I}{2R}$$

The field lines are circles concentric with the wire.

- **Magnetic field B inside a long Solenoid carrying a current I :**

$$B = \mu_0 n I$$

Where n is the number of turns per unit length.

- For a toroid,

$$B = \frac{\mu_0 N I}{2\pi r}$$

Where N is the total numbers of turns and r is the average radius.

- **Magnetic Moment of a Planar Loop:**

Magnetic moment m of a planar loop carrying a current I , having N closely wound turns, and an area A , is

$$\vec{m} = NI \vec{A}$$

- **Direction of \vec{m} is given by the Right – Hand Thumb Rule:**

Curl and palm of your right hand along the loop with the fingers pointing in the direction of the current, the thumb sticking out gives the direction of

$$\vec{m}(\text{and } \vec{A})$$

- **Loop placed in a Uniform Magnetic Field:**

a) When this loop is placed in a uniform magnetic field B ,

Then, the force F on it is, $F = 0$

And the torque on it is, $\vec{\tau} = \vec{m} \times \vec{B}$

In a moving coil galvanometer, this torque is balanced by a counter torque due to a spring, yielding.

$$k\phi = NI AB$$

where ϕ is the equilibrium deflection and k the torsion constant of the spring.

- **Magnetic Moment in an Electron:**

An electron moving around the central nucleus has a magnetic moment μ_l , given by

$$\mu_l = \frac{e}{2m} l$$

where l is the magnitude of the angular momentum of the circulating electron about the central nucleus.

- **Bohr Magnetron:**

The smallest value of μ_l is called the Bohr magnetron μ_B

$$\mu_B = 9.27 \times 10^{-24} \text{ J/T}$$

FOR MORE FREE QUESTION
BANKS AND SAMPLE PAPER
WT'S UP ON : 7665186856

A Best Faculty Group in Meerut....

ADHIKAANSH  **ACADEMY**
IITJEE NEET IX X XI XII

सही समय और सही दिशा में कड़ी मेहनत, तो सफलता पक्की!

So why
to wait...



DIRECTOR
DEEPAK SAINI (DSA SIR)
B.TECH, M.TECH
NSIT DELHI UNIVERSITY

और **Class 11th** से ही शुरू करें,
Board के साथ-साथ,
NEET और **IITJEE** की तैयारी

Special Classes For Students Studying in std. 9th & 10th.

EX. FACULTY OF RESONANCE KOTA, AAKASH INSTITUTE MUMBAI
225/5, 1ST FLOOR, PANCHSHEEL COLONY, BEHIND PINNACLE TOWER, GARH ROAD, MEERUT. ☎ 8057870069
www.adhikanshiitjeemedical.com | adhikanshiitjeemedical@gmail.com

Take Dummy Admission
in Class 11th & Prepare For..

IITJEE & NEET

For Free Counselling &
Career Guidance contact on:

7665186856