### PERIODIC CLASSIFICATION OF ELEMENTS

#### **CONTENTS**

- > INTRODUCTION
- EARLY ATTEMPTS OF CLASSIFICATION
- ➤ MENDELEEV'S PERIODIC

  LAW & PERIODIC TABLE
- > MODERN PERIODIC TABLE

#### > Introduction

One hundred and fifteen elements are known till date and many more may be known in future. All elements differ in their properties due to the presence of different kinds of atoms in them. Elements combine to form millions of compounds.

#### **♦** Important Terms and Concepts :

- 1. Need of Classification: It is difficult to study each and every element individually and to know its properties and uses. Therefore, they have been classified into groups on the basis of their similarities in properties.
- 2. Classification: Classification means grouping of elements on the basis of similarities in properties, e.g., All of you belong to class X because you resemble in you properties, e.g., All of you have passed class IX, all of you are in age group of 13-15 years.

**3. Basis of Classification :** Classification is done on the basis of similarities in properties so that the systematic study could be made about them

## **Early attempts of classification**

#### **♦** Lavoisier's Classification:

Lavoisier classified elements into metals and nonmetals. This classification was based on certain distinctive physical properties such as hardness, malleability and luster. On the basis of these properties, sodium and lead were classed together as belonging to the group of metals.

#### Limitations

- (1) Hardness, malleability and luster were found to be the only common properties of sodium and lead, otherwise the two elements were entirely different.
- (2) In such a classification there was no place for elements with properties resembling those of metals as well as nonmetals.

There fore, Lavoisier's classification was found to be inadequate.

#### **Dobereiner's Classification:**

Law of triads In 1817, German chemist Johann Dobereginer classified elements having similar chemical properties into groups of three. These groups were called triads. He proposed a law known as Dobereiner's law of triads. According to this law, when elements are arranged in the order of increasing atomic mass in a triad, the atomic mass of the middle element was found to be approximately equal to the arithmetic mean of the atomic masses of the other two elements.

Set	I	Set	II	Set	-III
Element	Atomic mass	Element	Atomic mass	Element	Atomic mass
Calcium	40	Lithiu m	7	Chlorin e	35.5
Strontiu m	87.5	Sodium	23	Bromin e	80
Barium	137	Potassi um	39	Iodine	127
Average atomic of calcium barium $= \frac{40 + 13}{2}$	masses m and	Average atomic of lithiu potassium $= \frac{7+39}{2}$	masses m and n	Average atomic in chlorine a $= \frac{35.5 + 1}{2}$	nasses of nd iodine
Atomic r strontium		Atomic r sodium =		Atomic bromine =	

The classification of elements into triads was very successful in predicting the atomic mass and properties of the middle element. Further, this classification showed that there exists some relationship between the properties of elements and their atomic masses. This paved the way for future attempts at classification of elements.

**Limitation**: All the elements could not be grouped into triads.

#### **♦** Newlands' Classification:

Law of octaves In 1864, John Newlands, and English chemist, showed that when elements are arranged in the order of their increasing atomic masses, the eighth element, starting from a given element, was a kind of repetition of the first one, like the eighth note in an octave of music, i.e.,

sa re ga ma pa dha ni sa, where the first and the eighth note are same.

A part of Newlands' classification is given below where the figures under the symbols show the atomic masses

Octaves of music and Newlands' arrangement of elements												
Indian :	sa	re	ga	ma	pa	dha	ni					
Octaves	sa											
Western:	do	re	mi	fa	so	la	ti					
	do											
Newlands's	Н	Li	Be	В	C	N	O					
arrangement of	1.0	7.0	9.0	11.0	12.0	14.0	16.0					
elements with	F	Na	Mg	Al	Si	P	S					
atomic masses	19.0	23.0	24.0	27.0	28.0	31.0	32.0					
	Cl	K	Ca	Cr	Ti	Mn	Fe					
	35.5	39.0	40.0	52.0	48.0	55.0	56.0					
	Co and Ni	Cu	Zn	Y	In	As	Se					
	58.93 and 58.71	63.54	65.37	88.90	114.82	74.92	78.96					
	Br	Rb	Sr	Ce and La	Zr							
	79.90	85.47	87.62	140.12 and 138.91	91.22							

Starting from lithium (Li) the eight element is sodium (Na). The eight element starting from sodium is potassium. The properties of lithium, sodium and potassium are similar. The properties of beryllium, magnesium and calcium are similar too.

#### Limitation:

(i) This law worked well for lighter elements (up to calcium), but it could not be applied to heavier ones (elements of higher atomic masses) because starting from calcium every eight element was found to have properties different from those of the first element. (iii) In arranging elements in the form of a table, Newlands clubbed two elements together at the same place and in the same column. Not only this, he also placed some dissimilar elements in the same column. For example, cobalt (Co) and nickel (Ni) were clubbed together in the column of fluorine (F), chlorine (Cl) and bromine (Br) (under sa/do). We know that cobalt and nickel have properties entirely different from those of fluorine, chlorine and bromine. It is also known that cobalt and nickel have properties similar to those of iron. But iron (Fe) was placed in a column (under ni/ti) different from the column of cobalt and nickel.

However, this law support to the idea that the properties of elements depend upon the atomic masses. It also showed that the properties of elements are repeated after a certain interval, i.e., the properties of elements are periodic in nature.

## Þ

# Mendeleev's periodic law and periodic table

While working systematically on the physical and chemical properties of elements, Dmitri Invanovich Mendeleev noticed that properties of elements varied regularly with the atomic mass. He arranged the 63 elements then known in a table on the basis of similarities in properties. It was found that most of the elements occupied places in the table in order of their increasing atomic masses. In 1869, *Mendeleeve* formulated a law, now known as the periodic law. The law is stated as follows.

The properties of elements are periodic functions of their atomic masses. This means, if the elements are arranged in order of increasing atomic masses then those with similar properties are repeated at regular intervals.

On the basis of the periodic law, Mendeleev presented his classification in the form of a table, now known as *Mendeleev's periodic table*. A simplified version of this periodic table is given below. In this table, copper, silver and gold find places in groups I as well as VIII.

the proper			1								
Groups → Periods↓	I	II	Ш	IV	V	VI	VII		V]	Ш	
1	H 1										
2	Li	Be	В	С	N	0	F				
	7	9.4	11	12	14	16	19				
3	Na	Mg	Al	Si	P	S	Cl				
3	23	24	27.3	28	31	32	35.5				
4	K	Ca	?	Ti	V	Cr	Mn	Fr	Co	Ni	Cu
4	39	40		48	51	52	55	56	59	59	63
5	Cu	Zn	?	?	As	Se	Br				
3	63	65	68	72	75	78	80				
6	Rb	Sr	Yt	Zr	Nb	Mo	?	Ru	Rh	Pd	Ag
6	85	87	88	90	94	96	100	104	104	106	108
7	Ag	Cd	In	Sn	Sb	Te	I				
/	108	112	113	118	122	125	127				
8	Cs	Ba	Di	Ce	?	?	?			?	
O	133	137	138	140	'	'				4	
9	?	?	?	?	?	?	?				
10	?	?	Er	La	Ta	W	?	Os	Ir	Pt	Au
10	•		178	180	182	184		195	197	198	199
11	Au	Hg	T1	Pb	Bi	?	?		•		·
11	199	200	204	207	208	· ·					
12	9	?	?	Th	9	U					
12	?			231	?	240					

This table consists of vertical columns called groups and horizontal rows called periods. There are only eight groups in the table. Mendeleev left some vacant places (shown by question marks) for the yet undiscovered elements. Noble gases were not discovered then. So, he did not provide any place for them in his periodic table.

Mandeleev's idea was remarkable in that he used a fundamental atomic property (atomic mass) as the basis of classification. While classifying elements he laid special emphasis on tow factors.

- 1. Similar elements were grouped together.
- 2. Elements were arranged in order of increasing atomic masses.

#### ♦ Modified version of Mendeleev's Periodic Table:

The elements which were undiscovered and for whom Mendeleev had left vacant places were discovered later. Some of these are scandium (Sc), gallium (Ga) and germanium (Ge). These elements were accommodated in their proper places in the table. The elements helium (He), neon (Ne), argon (Ar), Krypton (Kr), Xenon (Xe) and radon (Rn) became known only towards the end of the nineteenth century. These elements, called noble gases, were placed in the table as a separate group, called 0 group. The periodic table had to be modified then. The modified version of the table is shown below.

Group →		I	I	I	III		ľ	V	V	V	V	Ί	V]	Π	VIII	0
Periods↓	A	В	A	В	A	В	A	В	A	В	A	В	A	В		
1	]	Н														Не
2	Li		Be			В		С		N		О		F		Ne
3	Na		Mg			Al		Si		P		S		Cl		Ar
4	K		Ca		Sc		Ti		V		Cr		Mn		Fe Co Ni	Kr
		Cu		Zn		Ga		Ge		As		Se		Br		
5	Rb		Sr		Y		Zr		Nb		Mo		Тс		Ru Rh Pd	Xe
		Ag		Cd		In		Sn		Sb		Te		I		
6	Cs		Ba		La*		Hf		Ta		W		Re		Os Ir Pt	Rn
		Au		Hg		Tl		Pb		Bi		Po		At		
7	Fr		Ra		Ac**											

Lanthanide series*	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Но	Er	Tm	Yb	Lu
(along with langhanum)														
Actinide series**	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
(along with actinium)														

# **♦** Features of the modified version of Mendeleev's periodic table :

1. Groups into subgroups: Each group of this periodic table is further divided into two subgroups A and B. The properties of elements within a subgroup resemble more markedly but they differ from those of the elements of the other subgroups. For example., lithium (Li), sodium (Na), potassium (K), etc., of subgroups IA have close resemblance of properties but they have hardly any resemblance to the coinage metals (Cu, Ag and Au) of subgroup IB.

Mendeleev allowed the subgroups to be represented within the same group.

2. Prediction of errors: This periodic table could predict errors in the atomic masses of some elements on the basis of their position in the periodic table. For example, when the periodic table was published, experimental value of the atomic mass of beryllium (Be) we was supposed to be 13.65 and its valency, 3. So, the position of Be should have been somewhere else, but Mendeleev placed it at its appropriate position on the basis of its properties. He further suggested that the atomic mass of Be needed correction. Mendeleev predicted its

atomic mass to be 9.1 and valency, 2. Latter investigations proved him right.

Similarly, the atomic mass of uranium was corrected from 120 to 240. Corrections were also made in the atomic masses of gold, platinum, etc.

3. Predictions of properties of higher to undiscovered element: We know that Mendeleev classified the elements in order of their increasing atomic masses. However, this order had to be ignored at some places to make sure that the elements with similar properties fell in the same group. In doing so, he left some vacant places in the table. These vacant places were kept reserved for elements not discovered till then. Mendeleev was confident that these elements would be discovered later and they would occupy these vacant places. Not only this, he also predicted the properties of these undiscovered elements on the basis of this study of his the properties of the neighboring elements. Amazingly, when the missing elements of Mendeleev's periodic table were discovered subsequently, their properties were found to be very similar to those predicted by Mendeleev.

The elements scandium, gallium and germanium were not known in 1871 but their existence was predicted by mendelev. He named these elements as eka-boron, eka-Aluminium and eka silicon when these elements were discovered, they were named scandium, gallium and germanium respectively and their properties were found to be in good agreement with those predicted by Mendeleev. Properties of ka-aluminium (predicted by Mendeleev) and those of the gallium (discovered later) are given in table.

Prope	rty	Eka-aluminium	Gallium
Atomic ma	SS	68	69.7
Formula oxide	of	$E_2O_3$	Ga <sub>2</sub> O <sub>3</sub>
Formula chloride	of	ECl <sub>3</sub>	GaCl <sub>3</sub>

Considering its atomic mass, titanium (Ti) should have been placed below aluminium in the periodic table, but Mendeleev placed is below silicon (Si) because the properties of titanium were similar to those of silicon. Thus, a gap was left below aluminium in the periodic table. This gap was filled up by

- gallium which was discovered later. The properties of gallium (Ga) were found to be similar to those of boron and aluminium.
- 4. Basic features intact: All the basic features of Mendeleev's periodic table are intact even today. Even when a new class of elements, i.e., noble gases, were discovered, they found place in a separate group called the zero group. The existing order of the periodic table was not at all disturbed.

#### **Discrepancies in Mendeleev's periodic table :**

Mendeleev's periodic table has the following defects.

- 1. Position of hydrogen: The position of hydrogen in the periodic table is anomalous. Hydrogen resembles alkali metals (Li, Na, K, etc). So it may be placed in the group of the halogens (VII A).
- 2. Position of lanthanides and actinides: The elements from atomic number 57 to 71 are collectively known as lanthanides. They do not have a proper place in the periodic table. They all have been placed at the same position in group III and period 6. Similarly, the actinides (atomic numbers 89-103) also have no proper place in the periodic table. These elements have also been placed in the same position, in group III and period 7.
- 3. Some similar elements are separated, while some dissimilar elements have been placed in the group: Some similar elements are separated in the periodic table. For example, copper (Cu) and mercury (Hg), silver (Ag) and thallium (Tl), and barium (Ba) and lead (Pb). On the other hand, some dissimilar elements have been placed together in the same group. For example, copper (Cu), silver (Ag) and gold (Au) have been placed in group I along with the alkali metals. Similarly, manganese (Mn) is placed in the group of the halogens.
- 4. Presence of some anomalous pairs of elements: In Mendeleev's periodic table, elements are arranged in order of increasing atomic mass. In some places, this order has been ignored.
  - (a) The atomic mass of argon is 40 and that of potassium is 39. But argon is placed before potassium in the periodic table.

- (b) The positions of cobalt and nickel are not in proper order. Cobalt (at. mass = 58.9) is placed before nickel (at. mass = 58.6).
- (c) Tellurium (at. mass = 127.6) is placed before iodine (at. mass = 126.9).
- (d) Thorium (at. mass = 232.12) is placed before protactinium (at. mass = 231)
- **5. Position of isotopes :** The isotopes of an element have no place in the periodic table.

The failure of Mendeleev's periodic law to explain the wrong order of the atomic masses of some elements and the position of isotopes led scientists working in this field to conclude that atomic mass cannot be the basis for the classification of elements. There must be a more fundamental property of elements which can be the basis of classification.

#### Anomalous pairs of elements

Element →	Ar	K	Со	Ni	Te	Ι	Th	Pa
Atomic mass	40	39	59.9	58.6	127.6	126.9	232.12	231
Group	0	IA	VIII	VIII	VIB	VII B	III B	III B

#### **Modern Periodic Table :**

Henry Moseley, an English physicist found that the atomic number (Z) was the fundamental property of an elements and not the atomic mass for classification of elements.

#### **♦** Modern Periodic Law:

"Properties of elements are periodic functions of their atomic numbers, i.e., the number of protons or electrons present in the neutral atom of an element."

#### **Long form of Periodic Table :**

Arranged in increasing order of their atomic numbers.

The prediction of properties elements and their compounds can be made with precision. All drawbacks of Mendeleev's Periodic Table vanish when the elements are arranged on the basis of increasing atomic numbers.

#### **Elements in a Group :**

(1) They show similar chemical properties due to similar outer electronic configuration, i.e., same number of valence electrons.

(2) They have gradation in properties due to gradually varying attraction of the nucleus and the outer valence electrons as we go down the group.

# **♦** Main Features of the Long Form of the Periodic Table:

- (1) It shows arrangement of elements based on modern periodic law.
- (2) There are 18 vertical columns known as groups.
- (3) There are 7 horizontal rows known as periods.
- (4) Elements having similar outer electronic configurations, i.e., having same valence electrons have been placed in same groups, e.g.,

M

O

Q

#### Group-1

K

L

H	(1)	1						
Li	(3)	2,	1					
Na	(11)	2,	8,	1				
$\square$ K	(19)	2,	8,	8,	1			
Rb	(37)	2,	8,	18,	8,	1		
Cs	(55)	2,	8,	18,	18,	8,	1	
Fr	(87)	2	8	18	32	18	8	1

Grou	ıp-2								■Bi	(83)	2,	8,	18,	32,	18,	5
		K	L	M	N	О	P	Q	Gro	up-16						
Ве	(4)	2	2						Gro	ир-10	K	L	M	N	O	P
Mg	(12)	2,	8,	2					Гр	(8)	2	6	171	11	Ü	•
□Ca	(20)	2,	8,	8,	2				s	(16)	2,	8,	6			
\$r	(38)	2,	8,	18,	8,	2			∟					6		
Ва	(56)	2,	8,	18,	18,	8,	2		_	(34)	2,	8,	18,	6	(	
Ra	(88)	2,	8,	18,	32,	18,	8,	2		(52)	2,	8,	18,	18,	6	
									□Po	(84)	2,	8,	18,	32,	18,	6
Grou	ıp-13								Gro	up-17						
		K	L	M	N	О	P				K	L	M	N	O	P
□₿	(5)	2	3						F	(9)	2	7				
A1	(13)	2,	8,	3					C1	(17)	2,	8,	7			
Ga	(31)	2,	8,	18,	3				Br	(35)	2,	8,	18,	7		
tn	(49)	2,	8,	18,	18,	3				(53)	2,	8,	18,	18,	7	
<u></u>	(81)	2,	8,	18,	32,	18,	3		At	(85)	2,	8,	18,	32,	18,	7
Grou	up-14								Cro	up-18						
		K	L	M	N	O	P		GIO	up-10	K	L	M	N	О	P
□C □c:	(6)	2	4	4					Ш	(2)		L	1 <b>V1</b>	11	O	1
Si   Ge	(14) (32)	2, 2,	8, 8,	4 18,	4				Не	(2)	2	0				
Sn	(50)	2,	8,	18,	18,	4			Ne	(10)	2,	8				
□₽b	(82)	2,	8,	18,	32,	18,	4		Ar	(18)	2,	8,	8			
Grou	ıp-15								<u></u> Kr	(36)	2,	8,	18,	8		
		K	L	M	N	О	P		Xe	(54)	2,	8,	18,	18,	8	
□N	(7)	2	5						Rn	(86)	2,	8,	18,	32,	18,	8
<u> </u>	(15)	2,	8,	5					(5)	_						number of
As	(33)	2,	8,	18,	5					gradua						ll increases
\$b	(51)	2,	8,	18,	18,	5						-				

Period 1	Н							Не
(K-shell)	1							2
Second Period	Li(3)	Be(4)	B(5)	C(6)	N(7)	O(8)	F(9)	Ne(10)
(K, L, shells)	2, 1	2, 2	2, 3	2, 4	2, 5	2, 6	2, 7	2, 8

- (6) Each group in the table signifies identical outer shell electronic configuration i.e., same valence electrons, e.g., group 1 has 1 valence electron, group 2 has 2 valence electrons, group 13 has 3, group 14 has 4 valence electrons.
- (7) Each period starts with filling of new shell, e.g.,

1st Period – K shell (1st shell) starts filling with Hydrogen and ends at Helium.

2<sup>nd</sup> Period – L shell (2<sup>nd</sup> shell) starts filling from Li (3) upto Ne (10)

3<sup>rd</sup> Period – M shell (3<sup>rd</sup> shell) start filling from Na (11) upto Ar (18)

4<sup>th</sup> Period – N shell (4<sup>th</sup> shell) starts filling from K (19) upto Kr (36) and so on.

- (8) The periodic table is divided in four blocks:
  - (a) s-block elements: Group 1 and 2 elements are called s-block elements.
  - (b) p-block elements : Group 13 to 18 elements are called p-block elements
  - (c) d-block elements: Group 3 to group 12 are called d-block elements or transition elements (in between s- block and p-block elements)
  - (d) f-block elements: The elements placed at the bottom of the periodic table are known as f-block elements. The fourteen elements after La(57) (Lanthanum) are called Lanthanoides and 14 elements after Actinium Ac (89) are called Actinoides.

#### **Naturally occurring Elements:**

Elements upto atomic number 92 occur in nature except Technetium, Tc (Z = 43) and Promethium, Pm (Z = 61) which are formed from radioactive elements where 'Z' represents atomic number.

**Synthetic Elements :** Elements beyond atomic number 92 are man-made elements. They are also called synthetic elements

#### **Groups:**

- (1) Elements in a group have same number of valence electrons.
- (2) The chemical properties of valence electrons, e.g., all the group 1 elements have 1 valence electron. They form positively charged ions by losing one electron, when required amount of energy is supplied to them i.e., Li<sup>+</sup>, Na<sup>+</sup>, K<sup>+</sup>.

# Group 1 elements are called alkali metals. Group 2 elements are called alkaline earth metals.

Group 2 elements when 2 valence electrons in the outermost shells. They can lose both the valence electrons to form dipositive cations, i.e., Be<sup>2+</sup>, Mg<sup>2+</sup>, Ca<sup>2+</sup>, etc. Positively charged ions are called cations.

Group 13 elements belong to boron family, 14 to carbon family, 15 to Nitrogen family, 16 to Oxygen family.

Group 16 elements contain 6 valence electrons in their outermost shells, i.e., two electrons less than the maximum number of electrons that can be present in the outermost shell. They can gain 2 electrons more easily rather than lose 6 electrons. They change into dinegative ions such as O<sup>2-</sup>, S<sup>2-</sup>. Group 17 elements called Halogens contain 7 valence electrons. They can gain one electron to

acquire stable electronic configuration, i.e., 8 electrons in the outermost shell and form uni-negative (single negative) ions such as F<sup>-</sup>, Cl<sup>-</sup>, Br<sup>-</sup>, I<sup>-</sup>.

Negatively charged ions are called anions.

Group 18 elements called noble gases, have their outermost, shell completely filled. The elements of this group have no tendency to lose or gain electrons. Thus, the elements of this group have zero valency and are almost unreactive. Hence they are called Noble gases. However, nowadays, compounds of Kr, Xe and Rn have been prepared.

In any particular group, the number of shells increase but the number of valence electrons remains the same.

#### Periods:

- (1) The horizontal rows in the periodic table are called periods.
- (2) There are 7 periods in the long form of periodic table
- (3) The first period contains 2 elements, Hydrogen and Helium. They have only one shell.

- (4) The second period contains 8 elements: Lithium Li(3), Beryllium Be(4), Boron B(5), Carbon C(6), Nitrogen N(7), Oxygen O (8), Fluorine F(9) and Neon Ne (10). The second period has 2 shells (K and L) and L shell is progressive filled.
- (v) The elements of 3<sup>rd</sup> period are:

3 <sup>rd</sup>	Na	Mg	Al	Si	P	S	Cl	Ar
Period	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
(KII M	2, 8, 1	2, 3, 2	2, 8, 3	2, 8, 4	2, 8, 5	2, 8, 6	2, 8, 7	2, 8, 8
shells)								

In 3<sup>rd</sup> period, 3<sup>rd</sup> shell (M-shell) is being progressively filled and there are three shells.

4th period has 18 elements

5<sup>th</sup> period has 18 elements

6<sup>th</sup> period has 32 elements

7<sup>th</sup> period has 32 elements

In periods, the number of valence electrons increases from left to right in s and p-blocks

18) 2 2 4.0026 Helium	10 Neon	18 Ar 39.948 Argon	36 <b>Kr</b> 33.80 rypton	54 Xe 131.29 Xenon	<b>86 Rn</b> 222 Radon		<b>↑</b>	17
4 <u>5</u>			35 Br 79.904 Bromine K		<b>85 A1</b> 210 Astatine			10
40	8 15.999 Nitrogen		-		<b>84 Po</b> 210 Polonium			
p-Block Elements VA VI/ (15) (16	7 14.007 Nitrogen	15 P 30.974 Phosporus	33 <b>AS</b> 74.922 Arsenic	<b>51 Sb</b> 121.76 Arsenic				0
5£			32 <b>Ge</b> 72.61 Germanium	50 <b>Sn</b> 118.71 Tin	<b>82</b> <b>Pb</b> 207.2 Lead			1.0
139A	5 10.811 Boron	13 <b>AI</b> 6.982 Aluminium	31 <b>Ga</b> 62.723 Gallium	49 In 114.82 Indium	<b>81 Ti</b> 204.59 Thallium			2
,	1	_	30 <b>Zn</b> 65.39 Zinc	<b>48</b> <b>Cd</b> 112.41 Cadmium	80 <b>Hg</b> 200.59 Mercury		lements	10
		<b>8</b> €	29 <b>Cu</b> 63.546 Copper	47 <b>Ag</b> 107.87 Silver	79 <b>Au</b> 196.97 Gold	<b>111 Uuu</b> 272 Unununium	f-Block Elements	73
		(10)	28 Nickel	<b>46 Pd</b> 106.42 Palladium	<b>78 Pt</b> 195.08 Platinum			63
		- <b>[6</b> ]	27 <b>Co</b> 58.933 Cobolt	45 <b>Rh</b> 102.91 Rhodium	77 	109 Une 268 Unnilennium		5
	:lements	<u>[8</u>	26 <b>Fe</b> 55.847 Iron	<b>44 Ru</b> 101.07 Ruthenium	76 <b>0s</b> 190.23 Osmium	108 <b>Uno</b> 269 Unniloctium		2
Noble gases	d-Block Elements	VIIB	5	43 <b>Tc</b> 98 Technetium	75 <b>Re</b> 186.21 Rhenium	107 <b>Uns</b> 264 Unnilseptium		9
N N		VIB (6)	24 <b>Cr</b> 51.996 Chrominum	42 <b>Mo</b> 95.94 Molybdenum	74 W 183.84 Tungsten	106 Unh 266 Unnilhexium		í
Metals Non metals Metalloids	3	VB (5)	<b>23 V</b> 50.941 Vanadium	41 Nb 92.906 Molybdenum	73 <b>Ta</b> 180.95 Tantalum	104 105 106		2
Metals  Non metal		<b>5</b>	<b>22 Ti</b> 47.867 Titanium	<b>40 Zr</b> 91.224 Zirconium	72 <b>Hf</b> 178.49 Hafnium	104 Unq 261 Unnilquadiun		
		3 3 3	<b>Sc</b> 44.956 Scandium	39 <b>X</b> 88.936 Yttrium	<b>57 La</b> * 138.91 Lanthanum	89 AC** 227 Actinium (		
S-Block Elements  (1)  1  1  1  IIA  1  Close (2)  Close (2)	<b>Be</b> 9.0122 Beryllium	12 Mg 24.305 Magnesium	20 <b>Ca</b> 40.078 Calcium	38 Sr 87.62 Strontium	56 <b>Ba</b> 137.33 Barium	<b>88 Ra</b> 226 Radium		
S-Block  LT  1.0079  Hydrogen	3 6.940 Lithium	22.990 Sodium	19 <b>K</b> 39.098 Potassium	37 <b>Rb</b> 85.468 Rubidium	<b>CS</b> 132.91 Cesium	<b>87</b> <b>Fr</b> 223 Francium		
Group Period	7	က	4	5	9	7		

		<b>\</b>						f-Block Ele	ements						<b>↑</b>
ď	**************************************	58 <b>C</b>		09	61 Pm	62 <b>Sm</b>	63 Fu	64 <b>Gd</b>	65 <b>Tb</b>	99	<b>6</b> 4	68 F	69 <b>E</b>	02 <b>X</b>	17
٥	Familianiue Series	140.12 Cerium	140.91 Praseodymium	144.24 Neodymium	145 Promethium			157.25 Gadolinium	158.93 Terbium	162.50 Oysprosium	0.00	167.26 Erbium	168.93 Thulium	173.07 Ytterbium	174.97 Lutetium
		90		92	93	94	92	96	26	86	66	100	101	102	103
7	** Actinide Series	۲ ۲	Pa	<b>&gt;</b>	S Z	Pu	Am	E	짪	<u>უ</u>	Es	E	S N	ŝ	Ľ
•		232.04	231.04	238.03	237	244	243	247	247	251	252	257	258	259	262
		Thorium	Protactinium	Uranium	Neptunium	Prutonium	Americium	Curium	Berkelium	Califonium	Einsteinium	Fermium	Mendelevium	Nobelium	_awrencium

#### **Periodicity in Properties :**

The properties of elements depends upon the electronic configuration which changes along a period and down a group in periodic table.

There is periodicity in properties, i.e., repetition of properties after a regular interval due to similarity in electronic configuration.

## **♦** Atomic Size (Atomic radii) :

Atomic size means radius of an atom. It is defined as distance between centre of nucleus and outermost shell (valence shell) of an isolated atoms.

#### **Ovalent Radii:**

It is defined as half of the distance between the centres of nuclei of two atoms (bond length) bonded by a single covalent bond, e.g., Bond length in case of H—H (Hydrogen molecule) is 74 pm.

Covalent radius =  $1/2 \times 74$  pm = 37 pm (picometre)

$$[1 \text{ pm} = 10^{-12} \text{ m}]$$

It can be measured in case of diatomic molecules of non-metals.

Metallic Radii: If is defined as half of the internuclear distance between the two metal ions in a metallic crystal. It is measured in case of metals.

#### **Variation of Atomic size in a Group:**

Size generally increases from top to bottom in a group.

**Reason :** It is due to addition of a new shell, i.e., number of shells go one increasing, e.g., pm stands for picometre, i.e.,  $10^{-12}$  m.

Group I	Electronic Configuration	No. of Shells	Atomic radius
			(pm)
Li(3)	2, 1	(2 shells)	133
Na(11)	2, 8, 1	(3 shells)	154
K(19)	2, 8, 8, 1	(4 shells)	201
Rb(37)	2, 8, 18, 8, 1	(5 shells)	216
Cs(55)	2, 8, 18, 18, 8, 1	(6 shells)	235

## **♦** Variation of Atomic size along a Period :

Atomic size goes on decreasing along a period from left to right

**Reason:** It is due to increase in nuclear charge (number of protons in nucleus) which pulls the electrons towards it, i.e., force of attraction between nucleus and valence electrons increase, therefore atomic size decreases, e.g.,

Elements of 2 <sup>nd</sup> Period Atomic radius (pm)	Elements of 3 <sup>rd</sup> Period Atomic radius (pm)	
Li (3)	Na(11)	
Be(4) 89	Mg(12)	
B(5) 80	Al(13) 125	
C(6)	Si(14) 117	

N(7)	P(15)	
70	110	
O(8)	S(16)	
66	104	
F(9)	Cl(17)	0
64	99	

## **Solution** In Energy and Electron Affinity:

Chemical nature and reactivity of an element depend upon the ability of its atoms to donate or accept electrons. This can be measured quantitatively with the help of ionisation energy and electron affinity of its atoms:

**Ionisation energy:** It is defined as the energy required to remove an electron completely from an isolated gaseous atom of an element. The energy required to remove the 1<sup>st</sup> electron is called first ionisation energy.

$$A(g) + I.E_1 \longrightarrow A^+(g) + e^-$$

♦ Second Ionisation Energy: he energy required to remove an electron from a unipositive ion is called the second ionisation energy.

$$A^{\scriptscriptstyle +}(g) + I.E_2 {\longrightarrow} A^{2^{\scriptscriptstyle +}}(g) + e^{\scriptscriptstyle -}$$

Te second ionisation energy is greater than the first ionisation energy due to increase in positive charge, hence increase in force of attraction between the nucleus and the valence electron.

Ist I.E. 
$$< 2^{nd}$$
 I.E.  $< 3^{rd}$  I.E.

#### **Variation of Ionisation energy down a Group:**

Ionisation energy goes on decreasing down a group.

Reason: It is due to the increase in the distance between the valence electrons and the nucleus as the atomic size increase down a group, the force of attraction between the nucleus and the valence electrons decrease, therefore, the energy required to remove the electrons, i.e., the ioisation energy goes on decreasing

#### **Example:**

Group I	Ionisation Energy	Group 2	Second Ionisation	First Ionisation
	(in kJ mol <sup>-1</sup> )		Energy	Energy (in
			(in kJ mol <sup>-1</sup> )	kJ mol <sup>-1</sup> )
Li	500	Be	1757	899
Na	496	Mg	1450	737
K	420	Ca	1146	590
Rb	403	Sr	1064	549
Cs	376	Ba	965	503

#### **♦** Variation of Ionisation energy along a Period :

It goes on increasing generally along a period from left to right with decrease in atomic size.

**Reason:** Due to decrease in atomic size, the force of attraction between the valence electrons and the nucleus increase and, therefore, more energy is required to remove electron.

#### Example:

Elements of 2 <sup>nd</sup> Period	I.E. in kJ mol <sup>-1</sup>
Li	500
Ве	900
В	801
C	1085
N	1400
О	1314
F	1680
Ne	2080

There is a decrease in ionisation energy from Be to B and from N to O, the reason of which you will study in higher classes.

Group 18 elements (noble gases) have the highest ionisation energy in respective periods due to stable electronic configuration, i.e., 8 electrons in their valence shells except He which has 2 electrons.

#### **Electron Affinity:**

It is the energy change when an electron is gained by a neutral gaseous atom converting it into a negatively charged ion.

It is a measure of attraction or affinity of the gaseous atom for an extra electron to be added.

$$Cl(g) + e^{-} \longrightarrow Cl^{-}(g) + E.A.$$

#### **Solution** Factors:

It depends upon atomic size as well as *electronic* configuration.

#### **Variation down the Group:**

Electron affinity goes on decreasing down the group in general.

**Reason:** Due to the increase in atomic size, the force of attraction between the nucleus and the electron to e added becomes less.

#### **♦** Variation along a Period :

Electron affinity increase from left to right in period.

**Reason:** It is due to decrease in atomic size which leads to an increase in the force of attraction between the nucleus and the electrons to be added.

#### Example:

Group 17	E.A. (kJ mol <sup>-1</sup> )
F	333
Cl	348
Br	324
I	295

However, deviations to this rule are observed in variation of electron affinity.

#### **♦** Metallic and Non-metallic Character:

Group 1 to 12 are metals. Group 13 to 18 comprise non-metals, metalloids and metals.

**Metalloids:** Those elements which resemble both metals and non-metals are called metalloids. They are also called semi-metals, e.g., Boron, Silicon, Germinaium, Arsenic, Antimony, Tellurium and Polonium.

#### **Properties of Metals:**

- (i) The are malleable.
- (ii) They are ductile.
- (iii) They are good conductors of heat and electricity.
- (iv) They have generally 1 to 3 valence electrons.
- (v) They have the same or less number of electrons in their outermost shell than the number of shells.
- (vi) They are mostly solids.

#### **Properties of Non-metals:**

- (i) They exist in solid, liquid or gaseous state.
- (ii) Non-metals are generally brittle.
- (iii) They are non conductors.
- (iv) They have 4 to 8 valence electrons.

#### **Variation of Metallic Character:**

Metallic character increases down a group due to decrease in ionisation energy. It decrease along a period due to increase in ionisation energy from left to right

#### **♦** Variation of Non-metallic Character:

Non-metallic character decreases down a group because of decrease in electron affinity which is due to increase in atomic size.

Along a period, non-metallic character increases from left to right due to increase in electron affinity which is due to decrease in atomic size

## **POINTS TO REMEMBER**

- ➤ Dobereiner's law of triads states that in a given set of three elements (triad) the atomic mass of the middle element is approximately equal to the average of the atomic masses of the other two elements.
- ➤ Newland's law of octaves states that if the elements are arranged in order of increasing atomic mass, the eighth element starting from a given element shows a repetition of the properties of the first element.
- ➤ According to Mendeleev's periodic law, the physical and chemical properties of elements are a periodic function of their atomic masses.
- ➤ Mendeleev's periodic table (i) helped correct the wrongly assigned values of atomic masses of some elements, and (ii) predicted the properties of some undiscovered elements.
- ➤ In Mendeleev's periodic table (i) isotopes had no place, and (ii) some similar elements are separated, while some dissimilar elements are placed in the same group.

- ➤ In Mendeleev's periodic table there are eight groups (I to VIII). Each group is divided into subgroups A and B, except group VIII. In the modern periodic table there are 18 groups.
- ➤ The vertical columns of the periodic table are called groups.
- ➤ The horizontal rows of the periodic table are called periods.
- ➤ Modern periodic law. The physical and chemical properties of elements are a periodic function of their atomic numbers.
- ➤ The modern periodic table (long form) is based on atomic number.
- ➤ There are seven periods in the periodic table. The first six periods consists of 2, 8, 8, 18, 18 and 32 elements respectively. The seventh periods is still incomplete.
- > Elements in a group have similar properties.
- ➤ The regular occurrence of similar chemical properties of elements with increasing atomic number is known as chemical periodicity.
- ➤ A series of transition elements starting from actinium are called actinides or actinoids.
- ➤ A family of elements headed by helium in the periodic table constitutes the noble gases. They are unrecactive.
- An element that is intermediate between metals and nonmetals is known as a metalloid.
- ➤ The elements of group 1 are called alkali metals.
- ➤ The elements of group 2 are called alkaline earth metals.

## **Exercise-I**

## A. Very Short Answer Type Questions

- Q.1 Classify as acidic, basic and amphoteric oxide Na<sub>2</sub>O, Al<sub>2</sub>O<sub>3</sub> and SO<sub>2</sub>
- Q.2 Pick the odd one out
  - (i) H, N, P, Na
- (ii) H, He, O, C
- (iii) C, N, O, Cs
- (iv) Cs, Cl, Ca, Cd
- Q.3 Give one example each of the following
  - (i) Metal belonging to Group 12.
  - (ii) Metal belonging to Group 2.
  - (iii) Non-metal belonging to the halogen group. (iv) Most reactive halogen.
  - (v) Alkali metal which is radioactive.
- **Q.4** Give one example each of transition element, lanthanide, actinide and radioactive element.
- Q.5 What are the vertical columns in the periodic table called?
- **Q.6** Give an example of an element discovered after Mendeleev gave the periodic table
- Q.7 An element of group 14 has an atomic number of 14. State whether this element will have metallic properties
- **Q.8** (i) What is the position of hydrogen in the periodic table?
  - (ii) Where are lanthanides and actinides placed in the periodic table?
- **Q.9** Two atoms A and B have the following distribution of protons and neutrons

Atoms	Protons	Neutrons
A	6	6
В	6	7

Which element do they represent? How are A and B related?

## **B.** Short Answer Type Questions

- **Q.10** State the modern periodic law
- Q.11 Explain Newland's law of octaves
- **Q.12** Give reasons for the following:
  - (i) Atomic size decreases as we move from left to right across a period
  - (ii) Atomic size increases as we move from top to bottom along a group.
- Q.13 Fluorine, chlorine, bromine and iodine were put in one group on the basis of their similar properties.
  - (A) What is the name of this group or family?
  - (B) State two of these similar properties.
- Q.14 Explain why potassium is more reactive than lithium and chlorine less reactive than fluorine.
- Q.15 State two defects in Mendeleev's periodic table
- Q.16 For the main groups of the periodic table, the metallic properties of the elements vary approximately with their position as shown in the chart below.

1	2	3	4	5	6	7	0
Н							Не
A							В
C							D

Find A, B, C, D

## C. Long Answer Type Questions

- Q.17 What were the early attempts to classify the elements?
- Q.18 What is meant by the statement, 'properties of elements are a periodic function of atomic number'?

- **Q.19** Define the following:
  - (A) group, (B) period, (C) representative elements' (D) transition elements,
  - (e) lanthanides and (F) actinides
- Q.20 Sodium and aluminium have atomic numbers 11 and 13 respectively. They are separated by one element in the periodic table and have valencies of 1 and 3 respectively. Chlorine and potassium are also separated by one element in the periodic table. Their atomic numbers are 17 and 19 respectively, but both have a valency of 1.Explain why.
- Q.21 How does the modern periodic table differ from the Mendleeve's periodic table
- **Q.22** Write a short note on the following:
  - (i) Dobereiner's triads
  - (ii) Newland's law of octaves
  - (iii) Lothar Meyer's curves
- Q.23 Discuss the variation in the following along a group and a period:
  - (i) Atomic size
- (ii) Ionization energy
- (iii) Metallic character (iv) Electron affinity
- (v) Chemical reactivity

## **Exercise-II**

#### A. Fill in the blanks

- Q.1 Vertical columns of elements in the periodic table are called......
- Q.2 The number of electrons in the valence shell of the atom of an element indicates the ......to which the element belongs.
- **Q.3** The first period contains ......elements.
- **Q.4** The oxides of the elements of groups 1 and 2 are ..... in nature.
- Q.5 Electronegativity .....along a group from top to bottom.
- **Q.6** Alkali metals belong to group ...... and halogens belong to group ......in the periodic table.
- Q.7 The basis of modern periodic table is...........

#### **B.** True /False Type Questions

- **Q.8** Elements of a group have the same chemical properties.
- **Q.9** Element with least ionization energy is Ceasuim.
- **Q.10** The element with configuration ns<sup>2</sup>np<sup>5</sup> will belong to group 17.
- Q.11 Helium is the noble gas with least atomic size
- Q.12 Elements having atomic numbers 57 to 71 constitute the lanthanide series.
- Q.13 The nature of an element can be predicated on the basis of its position in the periodic table.
- Q.14 Within a group, various elements show variation in properties
- Q.15 From top to bottom in a group, electronegativity increases.