

3/10/2016

## Matthew's prime producing polynomial project

### A logical analysis of the trinomial $n^2 + n + 41$

We consider the composite values of  $n^2 + n + 41$ . We assume that  $n$  is an integer. We only consider positive integer values for  $n$  in this project.

The story so far

We analyze the behavior of the factorization of integers of the form  $h(n) = n^2 + n + 41$  where  $n$  is a non-negative integer. It was shown by Legendre, in 1798 that if  $0 \leq n < 40$  then  $h(n)$  is a prime number.

Certain patterns become evident when considering points  $(a, n)$  where  $h(n) \equiv 0 \pmod{a}$ . The collection of all such point produces what we are calling a "bifurcation" graph due to certain self-similar features. From experimental data we find that the integer points in this bifurcation graph lie on a collection of parabolic curves indexed by pairs of relatively prime integers. Each such pair yields (again determined experimentally and by observation of calculations) an integer polynomial  $a*z^2 + b*z + c$ , and the quartic  $h(a*z^2 + b*z + c)$  then factors non-trivially over the integers into two quadratic expressions. We call this our "parabola conjecture". Certain symmetries in the bifurcation graph are due to elementary relationships between pairs of co-prime integers. For instance if  $m < n$  are co-prime integers, then there is an observable relationship between the parabola it determines that that formed from  $(n-m, n)$ .

We conjecture that all composite values of  $h(n)$  arise by substituting integer values of  $z$  into  $h(a*z^2 + b*z + c)$ , where this quartic factors algebraically over  $\mathbf{Z}$  for  $a*z^2 + b*z + c$  a quadratic polynomial determined by a pair of relatively prime integers. We name this our "no stray points conjecture" because all the points in the bifurcation graph appear to lie on a parabola.

We further conjecture that the minimum  $x$ -values for parabolas corresponding to  $(m, n)$  with  $\gcd(m, n) = 1$  are equal for fixed  $n$ . Further, these minimum  $x$ -values line up at  $163*d^2/4$  where  $d = 1, 2, 3, \dots$ . The numerical evidence seems to support this. This is called our "parabolas line up" conjecture.

The notation  $\gcd(m, n)$  used above is defined here. The greatest common divisor of two integers is the smallest whole number that divides both of those integers.

Theorem 1 - Consider  $h(n)$  with  $n$  a non negative integer.  
 $h(n)$  never has a factor less than 41.

We prove Theorem 1 with a modular construction. We make a residue table with all the prime factors less than 41. The fundamental theorem of arithmetic states that any integer greater than one is either a prime number, or can be written as a unique product of prime numbers (ignoring the order). So if  $h(n)$  never has a prime factor less than 41, then by extension it never has an integer factor less than 41.

For example, to determine that  $h(n)$  is never divisible by 2, note the first column of the residue table. If  $n$  is even, then  $h(n)$  is odd. Similarly, if  $n$  is odd then  $h(n)$  is also odd. In either case,  $h(n)$  does not have factorization by 2.

Also, for divisibility by 3, there are 3 cases to check. They are  $n = 0, 1,$  and  $2 \pmod 3$ .  $h(0) \pmod 3$  is 2.  $h(1) \pmod 3$  is 1. and  $h(2) \pmod 3$  is 2. Due to these three cases,  $h(n)$  is never divisible by 3. This is the second column of the residue table.

The number 0 is first found in the residue table for the cases  $h(0) \pmod{41}$  and  $h(40) \pmod{41}$ . This means that if  $n$  is congruent to  $0 \pmod{41}$  then  $h(n)$  will be divisible by 41. Similarly, if  $n$  is congruent to  $40 \pmod{41}$  then  $h(n)$  is also divisible by 41. After the residue table, we observe a bifurcation graph which has points when  $h(y) \pmod x$  is divisible by  $x$ . The points  $(x,y)$  can be seen on the bifurcation graph.

\*see residue table page\*

Thus we have shown that  $h(n)$  never has a factor less than 41.

Theorem 2

Since  $h(a) = a^2 + a + 41$ , we want to show that  $h(a) = h(-a - 1)$ .

Proof of Theorem 2

Because  $h(a) = a^2 + a + 41$ ,

Now  $h(-a - 1) = (-a - 1)^2 + (-a - 1) + 41$ .

So  $h(-a - 1) = (-a - 1)^2 + (-a - 1) + 41$ ,

And  $h(-a - 1) = h(a)$

Which was what we wanted

End Proof of Theorem 2

Corollary 1


Further if  $h(b) \pmod c \equiv 0$  then  $h(c - b - 1) \pmod c \equiv 0$ .

We can observe broken bilateral symmetry in the bifurcation graph on a following page.




The function  $h(n) \sim n^2 + n + 41$  has interesting properties. Especially when  $n$  is restricted to the integers. As we know  $h(n)$  is a prime number for as  $n$  goes from 0 to 39.  $h(40)$  is  $41^2$ , a composite number.

Also  $h(n)$  can be generated recursively as  $h(0) = 0$  and  $h(n) = h(n-1) + 2*n$ . We checked this pair arithmetically.



Bifurcation Graph

These are pairs  $(x,y)$  such that  $h(y) \bmod x \equiv 0$ .



Here is a zoomed out iteration of the same graph.

There seems to be an apparent regular structure in this bifurcation graph.

Also there is a pattern about a slanted line that almost goes through the origin.