



Student Name _____

Date of Test _____

Hospital Scenario: "You are working in a hospital or clinic when a woman runs through the door, carrying an infant. She shouts, 'Help me! My baby's not breathing.' You have gloves and a pocket mask. You send your coworker to activate the emergency response system and to get the emergency equipment."

Prehospital Scenario: "You arrive on the scene for an infant who is not breathing. No bystander CPR has been provided. You approach the scene and ensure that it is safe. Demonstrate what you would do next."

Assessment and Activation

Checks responsiveness Shouts for help/Activates emergency response system

Checks breathing Checks pulse

Once student shouts for help, instructor says, "Here's the barrier device."

Cycle 1 of CPR (30:2) *CPR feedback devices are required for accuracy

Infant Compressions

Performs high-quality compressions*:

- Placement of 2 fingers or 2 thumbs in the center of the chest, just below the nipple line
- 30 compressions in no less than 15 and no more than 18 seconds
- Compresses at least one third the depth of the chest, approximately 1½ inches (4 cm)
- Complete recoil after each compression

Infant Breaths

Gives 2 breaths with a barrier device:

- Each breath given over 1 second
- Visible chest rise with each breath
- Resumes compressions in less than 10 seconds

Cycle 2 of CPR (repeats steps in Cycle 1) Only check box if step is successfully performed

Compressions Breaths Resumes compressions in less than 10 seconds

Rescuer 2 arrives with bag-mask device and begins ventilation while Rescuer 1 continues compressions with 2 thumb-encircling hands technique.

Cycle 3 of CPR

Rescuer 1: Infant Compressions

Performs high-quality compressions*:

- 15 compressions with 2 thumb-encircling hands technique
- 15 compressions in no less than 7 and no more than 9 seconds
- Compresses at least one third the depth of the chest, approximately 1½ inches (4 cm)
- Complete recoil after each compression

Rescuer 2: Infant Breaths

Gives 2 breaths with a barrier device:

- Each breath given over 1 second
- Visible chest rise with each breath
- Resumes compressions in less than 10 seconds

This rescuer is not evaluated.

Cycle 4 of CPR

Rescuer 2: Infant Compressions

This rescuer is not evaluated.

Rescuer 1: Infant Breaths

Gives 2 breaths with a bag-mask device:

- Each breath given over 1 second
- Visible chest rise with each breath
- Resumes compressions in less than 10 seconds

STOP TEST

Instructor Notes		
<ul style="list-style-type: none"> • Place a check in the box next to each step the student completes successfully. • If the student does not complete all steps successfully (as indicated by at least 1 blank check box), the student must receive remediation. Make a note here of which skills require remediation (refer to instructor manual for information about remediation). 		
Test Results	Check PASS or NR to indicate pass or needs remediation:	<input type="checkbox"/> PASS <input type="checkbox"/> NR
Instructor Initials _____	Instructor Number _____	Date _____