

2023 Recommended Immunizations for Children from Birth Through 6 Years Old

VACCINE	Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	19-23 MONTHS	2-3 YEARS	4-6 YEARS
HepB Hepatitis B	HepB	HepB				HepB					
RV* Rotavirus			RV	RV	RV*						
DTaP Diphtheria, Pertussis, & Tetanus			DTaP	DTaP	DTaP			DTaP			DTaP
Hib* Haemophilus influenzae type b			Hib	Hib	Hib*	Hib					
PCV13, PCV15 Pneumococcal disease			PCV	PCV	PCV	PCV					
IPV Polio			IPV	IPV		IPV					IPV
COVID-19** Coronavirus disease 2019									COVID-19**		
Flu[†] Influenza						Flu (One or Two Doses Yearly) [†]					
MMR Measles, Mumps, & Rubella						MMR					MMR
Varicella Chickenpox						Varicella					Varicella
HepA# Hepatitis A						HepA#		HepA#			

FOOTNOTES

RV* Administering a third dose at age 6 months depends on the brand of Hib or rotavirus vaccine used for previous dose.

Hib* recommended depends on your child's age and type of COVID-19 vaccine used.

COVID-19** Number of doses are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

Flu[†] Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.

HepA# Two doses of Hep A vaccine are needed for lasting protection. The 2 doses should be given between age 12 and 23 months. Both doses should be separated by at least 6 months. Children 2 years and older who have not received 2 doses of Hep A should complete the series.

ADDITIONAL INFORMATION


1. If your child misses a shot recommended for their age, talk to your child's doctor as soon as possible to see when the missed shot can be given.

2. If your child has any medical conditions that put them at risk for infection (e.g., sickle cell, HIV infection, cochlear implants) or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.

Talk with your child's doctor if you have questions about any shot recommended for your child.



FOR MORE INFORMATION
Call toll-free: 1-800-CDC-INFO (1-800-232-4636)
Or visit: [cdc.gov/vaccines/parents](https://www.cdc.gov/vaccines/parents)



American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

Diseases and the Vaccines that Prevent Them

BIRTH–6 YEARS OLD

DISEASE	VACCINE	DISEASE SPREAD BY	DISEASE SYMPTOMS	DISEASE COMPLICATIONS
Hepatitis B	HepB vaccine protects against hepatitis B.	Contact with blood or body fluids	May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain	Chronic liver infection, liver failure, liver cancer, death
Rotavirus	RV vaccine protects against rotavirus.	Through the mouth	Diarrhea, fever, vomiting	Severe diarrhea, dehydration, death
Diphtheria	DTaP* vaccine protects against diphtheria.	Air, direct contact	Sore throat, mild fever, weakness, swollen glands in neck	Swelling of the heart muscle, heart failure, coma, paralysis, death
Pertussis (whooping cough)	DTaP* vaccine protects against pertussis (whooping cough). ¹	Air, direct contact	Severe cough, runny nose, apnea (a pause in breathing in infants)	Pneumonia (infection in the lungs), death
Tetanus	DTaP* vaccine protects against tetanus.	Exposure through cuts in skin	Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever	Broken bones, breathing difficulty, death
Haemophilus influenzae type b (Hib)	Hib vaccine protects against Haemophilus influenzae type b.	Air, direct contact	May be no symptoms unless bacteria enter the blood	Meningitis (infection of the covering around the brain and spinal cord), intellectual disability, epiglottitis (life-threatening infection that can block the windpipe and lead to serious breathing problems), pneumonia (infection in the lungs), death
Pneumococcal disease (PCV13, PCV15)	PCV vaccine protects against pneumococcal disease.	Air, direct contact	May be no symptoms, pneumonia (infection in the lungs)	Bacteremia (blood infection), meningitis (infection of the covering around the brain and spinal cord), death
Polio	IPV vaccine protects against polio.	Air, direct contact, through the mouth	May be no symptoms, sore throat, fever, nausea, headache	Paralysis, death
Coronavirus disease 2019 (COVID-19)	COVID-19 vaccine protects against severe complications from coronavirus disease 2019.	Air, direct contact	May be no symptoms, fever, muscle aches, sore throat, cough, runny nose, diarrhea, vomiting, new loss of taste or smell	Pneumonia (infection in the lungs), respiratory failure, blood clots, bleeding disorder, injury to liver, heart or kidney, multi-system inflammatory syndrome, post-COVID syndrome, death
Influenza (Flu)	Flu vaccine protects against influenza.	Air, direct contact	Fever, muscle pain, sore throat, cough, extreme fatigue	Pneumonia (infection in the lungs), bronchitis, sinus infections, ear infections, death
Measles	MM/R** vaccine protects against measles.	Air, direct contact	Rash, fever, cough, runny nose, pink eye	Encephalitis (brain swelling), pneumonia (infection in the lungs), death
Mumps	MM/R** vaccine protects against mumps.	Air, direct contact	Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain	Meningitis (infection of the covering around the brain and spinal cord), encephalitis (brain swelling), inflammation of testicles or ovaries, deafness, death
Rubella	MM/R** vaccine protects against rubella.	Air, direct contact	Sometimes rash, fever, swollen lymph nodes	Very serious in pregnant women—can lead to miscarriage, stillbirth, premature delivery, birth defects
Chickenpox	Varicella vaccine protects against chickenpox.	Air, direct contact	Rash, tiredness, headache, fever	Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs), death
Hepatitis A	HePA vaccine protects against hepatitis A.	Direct contact, contaminated food or water	May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine	Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders, death

DTaP*

DTaP combines protection against diphtheria, tetanus, and pertussis.

MM/R**

MMR combines protection against measles, mumps, and rubella.

Last updated December 2022 • CS322257-A

2023 Recommended Immunizations for Children 7–18 Years Old



KEY
Indicates when the vaccine is recommended for all children unless your doctor tells you that your child cannot safely receive the vaccine.



Indicates the vaccine series can begin at this age.



Indicates the vaccine should be given if a child is catching up on missed vaccines. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses.



Indicates children not at increased risk may get the vaccine if they wish after speaking to a provider.

ADDITIONAL INFORMATION

1. If your child misses a shot recommended for their age, talk to your child's doctor as soon as possible to see when the missed shot can be given.
2. If your child has any medical conditions that put them at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that they may need.

Talk with your child's doctor if you have questions about any shot recommended for your child.

FOOTNOTES

COVID-19* Number of doses recommended depends on your child's age and type of COVID-19 vaccine used.

Flu** Two doses given at least 4 weeks apart are recommended for children age 6 months through 8 years of age who are getting an influenza (Flu) vaccine for the first time and for some other children in this age group.

HPV† Ages 11 through 12 years old should get a 2-shot series separated by 6 to 12 months. The series can begin at 9 years old. A 3-shot series is recommended for those with weakened immune systems and those who start the series after their 15th birthday.

*Originally recommended age ranges for missed childhood vaccinations: 2-dose series of MMR at 12–15 months and 4–6 years; 2-dose series of Varicella at 12–15 months and 4–6 years; 2-dose series of HepA (minimum interval: 6 months) at age 12–23 months; 3-dose series of HepB at birth, 1–2 months, and 6–18 months; and 4-dose series of Polio at 2 months, 4 months, 6–18 months, and 4–6 years.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

FOR MORE INFORMATION
Call toll-free: 1-800-CDC-INFO (1-800-232-4636)
Or visit: [cdc.gov/vaccines/parents](https://www.cdc.gov/vaccines/parents)



American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN™

Diseases and the Vaccines that Prevent Them

DISEASE	VACCINE	DISEASE SPREAD BY	DISEASE SYMPTOMS	DISEASE COMPLICATIONS
Coronavirus disease 2019 (COVID 19)	COVID-19	Air, direct contact	May be no symptoms, fever, muscle aches, sore throat, cough, runny nose, diarrhea, vomiting, new loss of taste or smell	Pneumonia (infection in the lungs), respiratory failure, blood clots, bleeding disorder, injury to liver, heart or kidney, multi-system inflammatory syndrome, post-COVID syndrome, death
Influenza (Flu)	Flu	Air, direct contact	Fever, muscle pain, sore throat, cough, extreme fatigue	Pneumonia (infection in the lungs), bronchitis, sinus infections, ear infections, death
Tetanus	Tdap* and Td** vaccines protect against tetanus.	Exposure through cuts in skin	Stiffness in neck and abdominal muscles, difficulty swallowing, muscle spasms, fever	Broken bones, breathing difficulty, death
Diphtheria	Tdap* and Td** vaccines protect against diphtheria.	Air, direct contact	Sore throat, mild fever, weakness, swollen glands in neck	Swelling of the heart muscle, heart failure, coma, paralysis, death
Pertussis (whooping cough)	Tdap*	Air, direct contact	Severe cough, runny nose, apnea (a pause in breathing in infants)	Pneumonia (infection in the lungs), death
Human papillomavirus	HPV	Direct skin contact	May be no symptoms, genital warts	Cervical, vaginal, vulvar, penile, anal, oropharyngeal cancers
Meningococcal disease	MenACWY and MenB	Air, direct contact	Sudden onset of fever, headache, and stiff neck, dark purple rash	Loss of limb, deafness, nervous system disorders, developmental disabilities, seizure disorder, stroke, death
Measles	MIMR†	Air, direct contact	Rash, fever, cough, runny nose, pink eye	Encephalitis (brain swelling), pneumonia (infection in the lungs), death
Mumps	MMR†	Air, direct contact	Swollen salivary glands (under the jaw), fever, headache, tiredness, muscle pain	Meningitis (infection of the covering around the brain and spinal cord), encephalitis (brain swelling), inflammation of testicles or ovaries, deafness, death
Rubella	MIMR†	Air, direct contact	Sometimes rash, fever, swollen lymph nodes	Very serious in pregnant women—can lead to miscarriage, stillbirth, premature delivery, birth defects
Chickenpox	Varicella	Air, direct contact	Rash, tiredness, headache, fever	Infected blisters, bleeding disorders, encephalitis (brain swelling), pneumonia (infection in the lungs), death
Hepatitis A	HePA	Direct contact, contaminated food or water	May be no symptoms, fever, stomach pain, loss of appetite, fatigue, vomiting, jaundice (yellowing of skin and eyes), dark urine	Liver failure, arthralgia (joint pain), kidney, pancreatic and blood disorders, death
Hepatitis B	HepB	Contact with blood or body fluids	May be no symptoms, fever, headache, weakness, vomiting, jaundice (yellowing of skin and eyes), joint pain	Chronic liver infection, liver failure, liver cancer, death
Polio	IPV	Air, direct contact, through the mouth	May be no symptoms, sore throat, fever, nausea, headache	Paralysis, death
Dengue	Dengue†	Bite from infected mosquito	May be no symptoms, fever, headache, pain behind the eyes, rash, joint pain, body ache, nausea, loss of appetite, feeling tired, abdominal pain	Severe bleeding, seizures, shock, damage to the liver, heart, and lungs, death

Tdap* Tdap combines protection against diphtheria, tetanus, and pertussis.

Td** Td combines protection against diphtheria and tetanus.

MMR†

MMR combines protection against measles, mumps, and rubella.

Dengue†

Recommended where dengue is common.

Last updated December 2022
CS32257-B