

Separable Differential Equations

Notes, Examples, and Practice Exercises (w/Solutions)

Topics include natural logarithms, integrals, direct and inverse variation, Newton's Law of Cooling, and more.

Mathplane.com

Differential Equations

Example: Find the particular equation for $\frac{dy}{dx} = \frac{-x}{y}$ at (2, 0)

Step 1: Separate the x's and y's

cross multiply $y \, dy = -x \, dx$

Step 2: Integrate each side

$$\int y \, dy = \int -x \, dx$$

$$\frac{y^2}{2} + C = \frac{-x^2}{2} + C$$

Step 3: Rewrite AND substitute the given point (to find C)

$$\frac{x^2}{2} + \frac{y^2}{2} = C$$

@ (2, 0): $\frac{(2)^2}{2} + \frac{(0)^2}{2} = C$

$$C = 2$$

Step 4: Write final equation with C value

$$\frac{x^2}{2} + \frac{y^2}{2} = 2$$

The particular equation is $x^2 + y^2 = 4$

Example: Find the general solution for $x \frac{dy}{dx} + y = 2x^2 y$

Step 1: Separate the x's and y's

$$x \frac{dy}{dx} = 2x^2 y - y$$

$$x \frac{dy}{dx} = y(2x^2 - 1)$$

$$\frac{x}{(2x^2 - 1)} \frac{dy}{dx} = y$$

$$\frac{x}{(2x^2 - 1)} \frac{1}{dx} = y \frac{1}{dy}$$

$$\frac{(2x^2 - 1)}{x} dx = \frac{1}{y} dy$$

Step 2: Integrate both sides

$$\int \frac{(2x^2 - 1)}{x} dx = \int \frac{1}{y} dy$$

$$\int \frac{2x^2}{x} - \frac{1}{x} dx = \ln y + C$$

$$\int 2x - \frac{1}{x} dx = \ln y + C$$

$$x^2 - \ln x + C = \ln y + C$$

Step 3: Rewrite (solve for y)

$$x^2 - \ln x + C = \ln y$$

rewrite in exponential form

$$\log_e y = x^2 - \ln x + C$$

$$y = e^{x^2 - \ln x + C}$$

exponent laws

$$y = \frac{e^{x^2}}{e^{\ln x}} \cdot e^C$$

since e^C can be any constant, we can substitute any constant C

$$y = \frac{C e^{x^2}}{x}$$

Example: Find a general solution for $x \frac{dy}{dx} + y = 0$

$$x \frac{dy}{dx} = -y$$

$$x \, dy = -y \, dx$$

$$\frac{1}{-y} dy = \frac{1}{x} dx$$

$$\int \frac{1}{-y} dy = \int \frac{1}{x} dx$$

$$-1 \ln y = \ln x + C$$

$$\ln y = -\ln x - C$$

$$\ln y = \ln \frac{C}{x}$$

$$y = \frac{C}{x}$$

Again, since C is any constant, it can be negative or positive...

Separable Differential Equations: Using Separation of Variables

Example: Find a general solution for the differential equation $\frac{dy}{dx} = e^{x-y}$

e^{x-y} can be re-written as $\frac{e^x}{e^y}$ using basic exponent properties

$$\frac{dy}{dx} = \frac{e^x}{e^y} \quad \text{If we cross-multiply, we separate the variables!}$$

$$e^y dy = e^x dx \quad \text{Then, integrate both sides}$$

$$\int e^y dy = \int e^x dx$$

$$e^y + C = e^x + C \quad \text{Combine the constants}$$

$$e^y = e^x + C \quad \text{Use logarithms to solve for y}$$

$$\ln e^y = \ln(e^x + C)$$

$$y = \ln(e^x + C)$$

Example: Solve the differential equation by using separation of variables

$$\frac{dy}{dx} = (x+2)(y+5)$$

$$dy(1) = dx(x+2)(y+5)$$

Separate the variables

$$dy \frac{1}{(y+5)} = dx(x+2)$$

$$\ln(y+5) + C = \frac{x^2}{2} + 2x + C$$

Since constants C are arbitrary, they can be combined...

$$\ln(y+5) = \frac{x^2}{2} + 2x + C$$

Change logarithm to exponential form

$$e^{\frac{x^2}{2} + 2x + C} = y + 5$$

$$C e^{\frac{x^2}{2} + 2x} = y + 5$$

Logarithm 'power rule' (exponent C can be moved to coefficient of term)

$$y = C e^{\frac{x^2}{2} + 2x} - 5$$

Exponential Equations $y = ab^x$

Examples: $A = Pe^{rt}$ $A = A_0 \left(1 + \frac{r}{n}\right)^{nt}$ $y = Ce^{kt}$



What is the rate of change?

$$\frac{dy}{dt} = Ce^{kt} \cdot k$$

$$= y \cdot k$$

$$\frac{dy}{dt} = ky$$

k is the constant of proportion
the rate of change varies directly with the output...

Direct Variation $y = kx$ where k is the constant of variation

Example: A model shows that $y = 14$ when $x = 7$.

If there is direct variation, then

if $x = 6/7$, what is y ?

{	$y = kx$	model for direct variation
	$14 = k(7)$	substitute x and y in order to find k
	$k = 2$	
	$y = 2x$	find y
	$y = 2(6/7)$	
	$y = 12/7$	

Indirect Variation $y = k \frac{1}{x}$ where k is the constant of variation
("Inverse" Variation)

Example: A guy drives 4 hours to his vacation spot at 40 miles/hour.
If the travel time (between home and vacation spot) varies inversely to speed,
how long will the return trip home take at 60 miles/hour.

step 1: write inverse variation general equation $y = \frac{k}{x}$

what is the constant of variation k? the distance!

$$y = \text{time of travel}$$

$$x = \text{speed of car}$$

step 2: find constant of variation, using given information

$$4 \text{ hours} = \frac{k}{40 \text{ miles/hour}} \quad k = 160 \text{ miles}$$

$$y = \frac{160 \text{ miles}}{x}$$

step 3: use equation to answer question $y = \frac{160 \text{ miles}}{60 \text{ m/h}} = 2 \frac{2}{3} \text{ hours or } 2 \text{ hours } 40 \text{ minutes}$

Example: Find the particular equation....

$$\frac{dy}{dx} = y + yx^2 \quad \text{for } y(0) = 5$$

separate the x's and y's

$$dy = y(1 + x^2) dx$$

$$\frac{1}{y} dy = (1 + x^2) dx$$

integrate each side...

$$\int \frac{1}{y} dy = \int (1 + x^2) dx$$

$$\ln|y| + C = x + \frac{x^3}{3} + C$$

consolidate, simplify, and solve for y

$$\ln|y| = x + \frac{x^3}{3} + C$$

$$y = e^{x + \frac{x^3}{3} + C}$$

$$y = e^{x + \frac{x^3}{3}} \cdot e^C$$

$$y = Ce^{x + \frac{x^3}{3}}$$

find C

$$y(0) = 5$$

$$5 = Ce^{0+0} \quad C = 5$$

$$y = 5e^{\left(x + \frac{x^3}{3}\right)}$$

Example: Bank earns interest *continuously* at a rate of 5%..

If the initial deposit is \$1000, write an equation representing the account balance...
(assume no deposit or withdrawal)

rate of change: $\frac{dy}{dt} = .05y$ where y is the account balance...

now, find the particular equation where $y(0) = 1000$

$$1 dy = .05y dt$$

$$\frac{20 dy}{y} = 1 dt$$

$$\int \frac{20 dy}{y} = \int 1 dt$$

$$20 \ln|y| + C = t + C$$

$$\ln|y| = \frac{t}{20} + C$$

$$y = e^{\frac{t}{20} + C}$$

$$y = Ce^{\frac{t}{20}}$$

and, since the original deposit is 1000

$$y = 1000e^{\frac{1}{20}t}$$

Example: At a local preserve, the rate of change of the coyote population "P" is directly proportional to (650 - P), where the time is measured in t years. At the beginning, the first census showed a population of 300. Two years later, the population increased to 500. What is the projected population after 5 years?

"The population growth is directly proportional to (650 - P)

$$\frac{dP}{dt} = k(650 - P)$$

We have the rate of change...

Now, we want to find the population (growth) function..

$$dP = k(650 - P) \cdot dt$$

$$dP \cdot \frac{1}{(650 - P)} = k \cdot dt$$

$$\int \frac{1}{(650 - P)} dP = \int k dt$$

$$-\ln(650 - P) + C = kt + C$$

$$\ln(650 - P) = -kt - C$$

$$650 - P = e^{-kt - C}$$

$$650 - P = e^{-kt} \cdot e^{-C} \text{ --- a constant...}$$

$$-P = Ce^{-kt} - 650$$

$$P = 650 - Ce^{-kt}$$

To find the specific solution...

$$P(t) = 650 - Ce^{-kt}$$

When $t = 0$, $P(t) = 300$

$$300 = 650 - Ce^{-k(0)}$$

$$C = 350$$

To find the growth (k)...

$$P(t) = 650 - 350e^{-kt}$$

When $t = 2$, $P(t) = 500$

$$500 = 650 - 350e^{-k(2)}$$

$$-150 = -350e^{-k(2)}$$

$$\frac{3}{7} = e^{-2k}$$

$$k = \frac{\ln(3/7)}{-2} = .4236$$

Finally, to find the number of coyotes at $t = 5$

$$P(t) = 650 - 350e^{-.4236t}$$


$$P(5) = 650 - 350e^{-.4236(5)}$$

= 607.9

Check: If $t = 0$... $P(0) = 650 - 350e^{-.4236(0)} = 300$

If $t = 2$... $P(2) = 650 - 350e^{-.4236(2)} = 499.98$

Here's a graph...



Example: Newton's Law of Cooling states that the rate of change in temperature of an object is directly proportional to the difference between its temperature and the temperature of the medium.
 An outdoor thermometer reading 85 degree F is brought into a 72 degree F room.
 After 2 minutes, the thermometer reads 83 degrees.
 When will the thermometer read 77 degrees? 75 degrees? 70 degrees?

$$\frac{dT}{dt} = k(T_0 - S)$$

T_0 = initial temperature
(i.e. temperature of the object)

S = temperature of the medium
(i.e. temperature of the Surroundings)

k = constant of proportion

$$\frac{dT}{dt} = k(T_0 - 72)$$

$$\frac{1}{(T_0 - 72)} dT = k dt$$

$$\int \frac{1}{(T_0 - 72)} dT = \int k dt$$

$$\ln(T_0 - 72) + C = kt + C$$

$$\ln(T_0 - 72) = kt + C$$

$$C e^{kt} = T_0 - 72$$

$$T_0 = C e^{kt} + 72$$

We know that the original temperature is 85 degrees when $t = 0$

$$85 = C e^{k(0)} + 72 \quad t = 0$$

$$C = 13 \quad T_0 = 85$$

$$T(t) = 13e^{kt} + 72$$

2 minutes later, the temperature is 83 degrees...

$$83 = 13e^{k(2)} + 72$$

$$\frac{11}{13} = e^{2k}$$

$$\ln(11/13) = 2k$$

$$k = -.0835 \text{ (approx)}$$

$$T(t) = 13e^{-.0835t} + 72$$

If temperature cools to 77 degrees...

$$77 = 13e^{-.0835t} + 72$$

$$\frac{5}{13} = e^{-.0835t}$$

$$t = 11.44 \text{ minutes}$$

If temperature is 75 degrees...

$$75 = 13e^{-.0835t} + 72$$


$$\frac{3}{13} = e^{-.0835t}$$

$$t = 17.56 \text{ minutes}$$

To check: let $t = 2$ minutes

$$T(0) = 13e^{-.0835(0)} + 72 = 85$$

$$T(2) = 13e^{-.0835(2)} + 72 = 83$$



Can the temperature reach 70 degrees?

$$70 = 13e^{-.0835t} + 72$$

$$-2 = 13e^{-.0835t}$$


It's an exponential equation...
The range is only positive numbers!

$$\frac{-2}{13} = e^{-.0835t}$$

$$\ln(-2/13) = -.0835t$$

NO SOLUTION

This is Benjamin: he's a little worried about his future.



*The (Math)
Graduate*

"Are you listening?
.... *Postulates*.
There's a great future in *postulates*.
Think about it..."

LanceAF #32 5-13-12
www.mathplane.com

This sequel to *The Graduate*
went straight to video.

Practice Exercises →

1) Find the general solution: $\frac{dy}{dx} = 6x^2 + 3$

2) Find the particular solution: $\frac{dy}{dx} = 6x^2 + 3$
if $y = 30$ when $x = 1$

3) Find the general solution: $\frac{dy}{dx} = 2xy$

4) $\frac{dy}{dx} = 3x^2(y - 3)$ and $y = 12$ when $x = 0$.

Find the particular solution.

A) A calf that weighs 60 pounds at birth has a rate of change in weight with respect to time (t years) that is directly proportional to the difference between 1200 and its weight.

$$\frac{dw}{dt} = k(1200 - w)$$

If $k = .8$, when will the calf's weight reach 800 pounds?

B) For my 30th birthday, my parents gave me \$10,000 to save... I placed it in an account that pays 4% per year... Each month, I deposit \$200 into the account...

a) If I received no interest, (just the original amount plus the \$200 per month), how much will I have when I'm 50?

b) Using calculus create an equation that describes how much money you have at any given time..

c) How much will you have on your 40th birthday? 50th birthday?

- C) At age 17, you won the lottery.
5,000,000 dollars go into a fund that returns 4%.
Meanwhile, you spend 300,000 each year.


How much money will you have 20 years later?

- D) Suppose a body is found at 4:00 am, and it is 30 degree C. At 6:00 am, the body temperature is 28 degrees.
If the room is 20 degrees C, then when was the murder?
(Common body temperature is 37 degrees)



Separable Differential Equations: Slope Fields

Sketch the slope field. Then, find the antiderivative to verify your sketch.



1) $\frac{dy}{dx} = 2x$




2) $\frac{dy}{dx} = y + 1$



3) $\frac{dy}{dx} = \frac{-x}{y}$





SOLUTIONS-→

1) Find the general solution: $\frac{dy}{dx} = 6x^2 + 3$

$\frac{dy}{dx} = \frac{6x^2 + 3}{1}$ (cross multiply) to separate the variables

$1 dy = 6x^2 + 3 dx$ integrate

$\int 1 dy = \int 6x^2 + 3 dx$ simplify

$y + C = 2x^3 + 3x + C$

$y = 2x^3 + 3x + C$

(note: since the constants are arbitrary, they can be combined to make one constant C)

2) Find the particular solution: $\frac{dy}{dx} = 6x^2 + 3$

$\frac{dy}{dx} = \frac{6x^2 + 3}{1}$

if $y = 30$ when $x = 1$

$1 dy = 6x^2 + 3 dx$

substitute x and y: $30 = 2(1)^3 + 3(1) + C$

$\int 1 dy = \int 6x^2 + 3 dx$

$30 = 5 + C$

$y = 2x^3 + 3x + C$

$C = 25$

$y = 2x^3 + 3x + 25$

3) Find the general solution: $\frac{dy}{dx} = 2xy$

$\frac{dy}{dx} = \frac{2xy}{1}$

(cross multiply)

$1 dy = 2xy dx$

separate the variables

$\frac{1}{y} dy = 2x dx$

integrate

$\int \frac{1}{y} dy = \int 2x dx$

$\ln|y| + C = x^2 + C$

$\ln|y| = x^2 + C$ (convert log function into exponential form)

$y = e^{x^2 + C}$

$y = e^{x^2} \cdot e^C$ (exponent laws)

$y = C e^{x^2}$

(note: since C can be any number, presumably e^C can be any constant.)

4) $\frac{dy}{dx} = 3x^2(y - 3)$ and $y = 12$ when $x = 0$.

cross multiply $3x^2(y - 3) dx = dy$

To change the general solution to a specific solution, we use the given point:

Find the particular solution.

separate the variables (x and y) $3x^2 dx = \frac{1}{(y-3)} dy$

$(0, 12)$

integrate $\int 3x^2 dx = \int \frac{1}{(y-3)} dy$

$12 = C e^{0^3} + 3$

rearrange and combine the constants C

$x^3 + C = \ln(y - 3) + C$

$12 = C(1) + 3$

$C = 9$

convert to exponential form

$\ln(y - 3) = x^3 + C$

$y = 9e^{x^3} + 3$

(exponent laws)

$e^{x^3 + C} = y - 3$

Check: find derivative...

Since C can be any constant e^C can be any number

$y = e^{x^3} \cdot e^C + 3$

$\frac{dy}{dx} = 9e^{x^3} \cdot 3x^2 + 0$

**So, we'll simplify and use C to be our constant, rather than e^C

$y = C e^{x^3} + 3$

we know that $9e^{x^3} = y - 3$

so, $\frac{dy}{dx} = (y - 3) \cdot 3x^2$ ✓

A) A calf that weighs 60 pounds at birth has a rate of change in weight with respect to time (t years) that is directly proportional to the difference between 1200 and its weight.

SOLUTIONS

$$\frac{dw}{dt} = k(1200 - w)$$

If $k = .8$, when will the calf's weight reach 800 pounds?

$$\frac{1}{(1200 - w)} dw = k dt$$

$$k = .8$$

$$w(0) = 60$$

$$\int \frac{1}{(1200 - w)} dw = \int k dt$$

$$w(t) = -Ce^{-kt} + 1200$$

When will the calf reach 800 pounds?

$$-\ln(1200 - w) + C = kt + C$$

$$w(0) = 60 = -Ce^0 + 1200$$

$$800 = -1140e^{-.8(t)} + 1200$$

$$\ln(1200 - w) = -kt - C$$

$$C = 1140$$

$$-400 = -1140e^{-.8(t)}$$

$$1200 - w = e^{-kt - C}$$

$$w(t) = -1140e^{-.8t} + 1200$$

$$\frac{400}{1140} = e^{-.8(t)}$$

$$1200 - w = e^{-kt} \cdot e^{-C}$$

$$w = -Ce^{-kt} + 1200$$

$$t = 1.309 \text{ years}$$

B) For my 30th birthday, my parents gave me \$10,000 to save... I placed it in an account that pays 4% per year... Each month, I deposit \$200 into the account...

a) If I received no interest, (just the original amount plus the \$200 per month), how much will I have when I'm 50?

$$\$10,000 + \$200(240 \text{ months}) = \boxed{\$58,000}$$

b) Using calculus create an equation that describes how much money you have at any given time..

$$\frac{dF}{dt} = 200 + \frac{.04}{12} F$$

F = amount in the Fund
t = time in months

$$dt = \frac{1}{200 + \frac{.04}{12} F} dF$$

$$\int dt = \int \frac{1}{200 + \frac{.04}{12} F} dF$$

$$t + C = \frac{12}{.04} \ln \left[200 + \frac{.04}{12} F \right]$$

$$\frac{.04}{12} t + C = \ln \left[200 + \frac{.04}{12} F \right]$$

$$e^{\frac{.04}{12} t + C} = \left[200 + \frac{.04}{12} F \right]$$

$$e^{\frac{.04}{12} t} \cdot e^C = 200 + \frac{.04}{12} F$$

$$C e^{\frac{.04}{12} t} = 200 + \frac{.04}{12} F$$

$$\frac{.04}{12} F = C e^{\frac{.04}{12} t} - 200$$

Since we have 10,000 dollars at the beginning $t = 0$
 $F = 10,000$

$$\frac{.04}{12} (10,000) = C e^0 - 200$$

$$C = 233.333$$

$$F = \frac{12}{.04} \left[233.333 e^{\frac{.04}{12} t} - 200 \right]$$

c) How much will you have on your 40th birthday? 50th birthday?

10 years later ----> 120 months...

$$F(120) = \frac{12}{.04} \left[233.333 e^{\frac{.04}{12} (120)} - 200 \right] = \boxed{\$44,427}$$

$$F(240) = \frac{12}{.04} \left[233.333 e^{\frac{.04}{12} (240)} - 200 \right] = \boxed{\$95,787}$$

Note: if we test 1 month: $t = 1$

\$10,000 initial fund

$\frac{.04}{12} (10,000) = \33.33 interest

\$200 contribution

\$10,233.33

then, using our formula:

$$F(1) = \frac{12}{.04} \left[233.333 e^{\frac{.04}{12} (1)} - 200 \right] = \$10,233.72$$

**Note, our formula assumes the interest is applied daily, rather than monthly... So, there is a slight difference!

C) At age 17, you won the lottery.
5,000,000 dollars go into a fund that returns 4%.
Meanwhile, you spend 300,000 each year.

How much money will you have 20 years later?

$$\frac{dF}{dt} = .04F - 300,000 \quad \text{rate of change each year (t)}$$

$F = \text{amount of money in the fund}$

$$dt = \frac{1}{(.04F - 300,000)} dF$$

$$\int dt = \int \frac{1}{(.04F - 300,000)} dF$$

$$t + C = \frac{1}{.04} \ln(.04F - 300,000) + C$$

$$.04t + C = \ln(.04F - 300,000)$$

$$Ce^{.04t} = .04F - 300,000$$

$$.04F = Ce^{.04t} + 300,000$$

When $t = 20$,

$$F = \frac{-100,000 e^{.04(20)} + 300,000}{.04}$$

$$F = 1,936,150$$

At $t = 0$, $F = 5,000,000$

$$F = \frac{Ce^{.04t} + 300,000}{.04}$$

$$5,000,000 = \frac{Ce^{.04(0)} + 300,000}{.04}$$

$$5,000,000 = \frac{C + 300,000}{.04}$$

$$C = -100,000$$

$$F = \frac{-100,000 e^{.04t} + 300,000}{.04}$$

$$t = 1 \quad 4,897,975$$

$$t = 2 \quad 4,791,775$$

$$t = 3 \quad 4,681,250$$

check for reasonableness:

$$A = Pe^{rt}$$

1 year:	$5,000,000e^{-.04(1)}$	=	5,204,055	
			- 300,000	300,000 spent
			490,4055	
			- 12243	4% of 300,000 compounded continuously
			4,891,812	

D) Suppose a body is found at 4:00 am, and it is 30 degree C. At 6:00 am, the body temperature is 28 degrees.
If the room is 20 degrees C, then when was the murder?
(Common body temperature is 37 degrees)

Murder committed, and body starts at 37 degrees C.

Newton's Law of cooling

find temperature of body (H) as function of (t) time since murder

$$\frac{dH}{dt} = k(H - 20)$$

$$dH = k(H - 20) dt$$

$$\frac{1}{(H - 20)} dH = k dt$$

$$\int \frac{1}{(H - 20)} dH = \int k dt$$

$$\ln |H - 20| + C = kt + C$$

$$\ln |H - 20| = kt + C$$

$$H - 20 = e^{kt + C}$$

$$H = Ce^{kt} + 20$$

When the murder occurred,

$$t = 0$$

$$37 = Ce^{k(0)} + 20$$

$$17 = C$$

$$H = 17e^{kt} + 20$$

$$\frac{dT_0}{dt} = k(T_0 - S)$$

$T_0 = \text{initial temperature (i.e. temperature of the object)}$

$S = \text{temperature of the medium (i.e. temperature of the Surroundings)}$

$k = \text{constant of proportion}$

At 4:00 AM, the body was 30 degrees

$$30 = 17e^{kt} + 20$$

At 6:00 AM, the body was 28 degrees...

$$28 = 17e^{k(t+2)} + 20$$

solve the system:

$$k = -.112$$

$$t = 4.76$$

11:15 PM (approx.)

$$T(t) = T_s + (T_0 - T_s)e^{-kt} \quad \text{(Using the two body temps at 4:00 and 6:00)}$$

2 hours later:

$$28 = 20 + (30 - 20)e^{-k(2)}$$

$$8 = 10e^{-2k}$$

$$\frac{8}{10} = e^{-2k}$$

$$k = .111572$$

$$\text{time of death: } 37 = 20 + (30 - 20)e^{-.111572 t}$$

$$\frac{17}{10} = e^{-.111572 t}$$

$$\text{time } t = -4.756$$

4 hours, 45 minutes, 21 seconds earlier ✓


Separable Differential Equations: Slope Fields

Sketch the slope field. Then, find the antiderivative to verify your sketch.

1) $\frac{dy}{dx} = 2x$

$y = x^2 + C$

at $x = 1$, the slopes of each segment are 2
 at $x = 2$, the slopes are 4
 at $x = 0$, the slopes are 0
 at $x = -1$, the slopes are -2
 etc..



2) $\frac{dy}{dx} = y + 1$

$(y + 1)dx = dy$

$\int dx = \int \frac{1}{(y + 1)} dy$

$x + C = \ln|y + 1| + C$

$x + C = \ln|y + 1|$


$y + 1 = e^{x + C}$

$y + 1 = e^x + e^C$

$y = Ce^x - 1$

at $y = -1$, all segments have slopes equal to 0 (horizontal)
 at $y = 0$, all segments have slopes equal to 1
 at $y = 1$, all segments have slopes equal to 2
 at $y = -2$, all segments have slopes equal to -1
 etc...

(if C is negative, then the curve opens downward...)



3) $\frac{dy}{dx} = \frac{-x}{y}$

$y dy = -x dx$


$\int y dy = \int -x dx$

$\frac{y^2}{2} + C = \frac{-x^2}{2} + C$

$\frac{x^2}{2} + \frac{y^2}{2} = C$

$x^2 + y^2 = 2C$


at $y = 0$, the slopes are undefined (vertical lines)
 at $x = 0$, the slopes are zero (horizontal lines)
 when $x = y$, the slopes are -1
 when $x = -y$, the slopes are 1
 etc...
 others include (3, 1) ---> -3
 (3, -1) ---> 3
 (-3, 1) ----> 3



Thanks for visiting. (Hope it helped!)

If you have questions, suggestions or requests, let us know.

Cheers.



Mathplane *Express* for mobile and tablets at Mathplane.org

Also, our store is at TeachersPayTeachers