



Pendimethalin 0.86% + Fertilizer

Preemergence Weed Control / FOR USE ON TURFGRASSES AND ESTABLISHED LANDSCAPE ORNAMENTALS
NOT FOR USE IN ORGANIC CROP AND FOOD PRODUCTION IN CALIFORNIA

Group

3

Herbicide

ACTIVE INGREDIENT:

Pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine..... 0.86%

OTHER INGREDIENTS:.....99.14%

Total:.....100.00%

19-0-5

GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N)19.00%
19.00% Urea Nitrogen*

SOLUBLE POTASH (K₂O)5.00%

TOTAL SULFUR (S)1.75%
1.75% Free Sulfur (S)

DERIVED FROM: Polymer Coated Sulfur-Coated Urea, Urea, Muriate of Potash.

CHLORINE (Cl) Max.4.00%

* 5.10% Slowly Available Nitrogen From Polymer Coated Sulfur-Coated Urea.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

In case of an emergency endangering life or property involving this product, call day or night 800-832-HELP (4357).

See inside for additional First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

FIRST AID

| | |
|------------------------|---|
| If swallowed | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person. |
| If on skin or clothing | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice. |
| If in eyes | <ul style="list-style-type: none">• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes.• Call a poison control center or doctor for advice. |
| If inhaled | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or ambulance; then give artificial respiration, preferably by mouth-to-mouth if possible.• Call a poison control center or doctor for advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

Precautionary Statements

Hazards to Humans and Domestic Animals

Caution. Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. If you want more options, follow the instructions for **Category A** on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of waterproof material, such as butyl rubber ≥14 mils, nitrile rubber ≥14 mils, or neoprene rubber ≥14 mils.
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

For Florida ONLY: DO NOT apply near water, storm drains or drainage ditches. **DO NOT** apply if heavy rain is expected. Apply only to lawn, and sweep any product that lands on the driveway, sidewalk or street, back onto lawn.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at time of pesticide application.

BASF Corporation does not recommend or authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application to turf or ornamentals.

DO NOT APPLY PENDIMETHALIN 0.86% + FERTILIZER in greenhouses, shadehouses or other enclosed structures.

NOT FOR USE on grass grown for commercial seed production.

DO NOT apply this product through any irrigation system. Keep children and pets out of treated area until dust has settled.

DO NOT allow dust to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, nontarget crops, aquatic and wetland areas, woodlands, pasture, rangeland, or animals.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of waterproof material, such as butyl rubber ≥14 mils, nitrile rubber ≥14 mils or neoprene rubber ≥14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter or allow others to enter the treated area until dusts have settled.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR RESIDUES.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: **DO NOT** store above 120°F. Store in original container and keep closed. Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER HANDLING (For Paper or Plastic Bags): Nonrefillable Container. **DO NOT** reuse or refill this container. After completely emptying container into application equipment, dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Observe all cautions and limitations on this label and the labels of products used in combination with **PENDIMETHALIN 0.86% + FERTILIZER**. The use of **PENDIMETHALIN 0.86% + FERTILIZER** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

PRODUCT INFORMATION

PENDIMETHALIN 0.86% + FERTILIZER is a combination preemergence herbicide and fertilizer formulated to provide control of most annual grasses and certain broadleaf weeds found in any turfgrass site (lawns, sod, turf areas). Examples of such sites include, but are not limited to: golf courses, lawns in residential, commercial establishments, or grounds around military or other institutions, parks, airports, roadsides, schools or cemeteries; athletic fields or recreational areas; prairiegrass areas; and sod farms. **PENDIMETHALIN 0.86% + FERTILIZER** can be used around established ornamentals in the landscape and in established wildflower plantings. **PENDIMETHALIN 0.86% + FERTILIZER** may be used for hardwood and conifer regeneration on conservation reserve program land or similar areas.

PENDIMETHALIN 0.86% + FERTILIZER is labeled for use on the following turf grass species: Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass, *Poa annua* (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, centipedegrass, zoysiagrass and blends of these grasses.

PENDIMETHALIN 0.86% + FERTILIZER controls weeds as they germinate but will not control established weeds. Therefore, areas to be treated should be free of established weeds. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

APPLICATION INSTRUCTIONS

Apply product uniformly at the specified rate using a properly calibrated spreader. The efficacy of **PENDIMETHALIN 0.86% + FERTILIZER** will be improved if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the turfgrass foliage and establish a preemergent herbicide barrier in the soil. If **PENDIMETHALIN 0.86% + FERTILIZER** is not activated by rainfall or irrigation within 30 days, erratic weed control may result.

Avoid contact of product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

PENDIMETHALIN 0.86% + FERTILIZER can be applied before or after postemergence herbicides used to control emerged weeds in turfgrasses. For annual grass control, applications can be made with **Segment® herbicide**, **Acclaim Extra® herbicide**, or MSMA-containing products to control emerged weeds. Broadleaf weeds can be controlled using **Trimec® herbicide**, **Three-Way™ herbicide**, 2,4-D; and other similar products.

Applied according to label directions and under normal growing conditions, **PENDIMETHALIN 0.86% + FERTILIZER** will not cause injury to desirable plants. Overapplication can result in stand loss, plant injury, or soil residues. Uneven application can decrease weed control or cause plant injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of plant damage from **PENDIMETHALIN 0.86% + FERTILIZER**.

Apply with properly calibrated spreader equipment to provide uniform distribution. Check application routinely to determine proper calibration. Avoid overlaps that will increase rates above those specified. Avoid application when winds may cause drift.

Spreader Settings

| Spreader Type | Setting |
|---|---------|
| LESCO Commercial | #17 |
| Vigoro 4300 | 4 |
| Scotts Broadcast/Rotary (Scotts® Basic, Standard™, Deluxe EdgeGuard® or SpeedyGreen®) | 7 |

Maximum Application Rate by Use Sites

| Use Site | lbs Product/A | lbs Product/1000 ft ² | lbs a.i./A |
|--|---------------|----------------------------------|------------|
| Residential and Sod Farm Turf | 233 | 5.3 | 2.0 |
| Golf Course, Commercial and Other Non-residential Turf Sites | 349 | 8.0 | 3.0 |

* Residential is defined as turf in any residential situation as well as home lawns, schools, parks and playgrounds.

Florida Specific Requirements for Fertilizer Use

For applications to sports turf at golf courses, parks and athletic fields: DO NOT exceed fertilizer rates recommended in the document titled SL 191, "Recommendations for N, P, K and Mg for Golf Course and Athletic Field Fertilization Based on Mehlich I Extractant" dated March 2007. Copies may be downloaded from <http://edi.ifas.ufl.edu/SS404>.

Follow "Best Management Practices for the Enhancement of Environmental Quality on Florida Golf Courses," published by the Florida Department of Environmental Protection, dated January 2007. Copies may be downloaded from <http://www.dep.state.fl.us/water/nonpoint/pubs.htm>.

For all fertilizer uses other than as a specialty fertilizer on urban turf: DO NOT exceed rates recommended in the document titled Best Management Practices for Protection of Water Resources in Florida, June 2002, Florida Green Industries. This document may be downloaded from http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book.

ACCEPTED

FOR REGISTRATION

Feb 4, 2014

Doc id: 537656

#900133

Net Weight 50 lbs. (22.67 kg)

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration



Pendimethalin 0.86% + Fertilizer

TURFGRASS

RESIDENTIAL, GOLF COURSE, COMMERCIAL AND OTHER NON-RESIDENTIAL SITES

Application Rates For Preemergence Weed Control¹

| | Turfgrass Species | Weed | Pendimethalin 0.86% + Fertilizer Rate | Comment |
|-------------------|---|---|---|---|
| Cool Season Grass | Bluegrass, Kentucky / Fescue, fine / Fescue, tall / Ryegrass, perennial. | crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover | All Turf Uses 175-233 lbs/acre as the initial application before weed germination in the spring | Apply a repeat application of 116-175 lbs/acre after 5-8 weeks for extended control or where heavy weed infestations are expected. |
| | | goosegrass | Residential and Sod Farm Turf Uses Only² 175-233 lbs/acre in the spring before weed germination | Apply a repeat application of 116 lbs/acre after 5-8 weeks if the lower rate was used initially or for extended goosegrass control. |
| | | | Golf Course, Commercial and Other Non-Residential Turf Sites 175-349 lbs/acre in the spring before weed germination | |
| | Bentgrass or established <i>Poa annua</i> ³ (1/2 inch or taller) | cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell | All Turf Uses 175-233 lbs/acre | Apply in late summer or early fall before weed germination. Apply a repeat application of 175-233 lbs/acre after 5-8 weeks for extended <i>Poa annua</i> control. |
| | | crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover | All Turf Uses (Non-Greens and Tees) 175 lbs/acre in the spring before weed germination. | Apply a repeat application of 116-175 lbs/acre after 5-8 weeks where heavy weed infestations are expected. |
| | | goosegrass | All Turf Uses (Non-Greens and Tees) 175 lbs/acre in the spring before weed germination. | Apply a repeat application of 175 lbs/acre after 5-8 weeks for extended control. |
| Warm Season Grass | Bahiagrass / Bermudagrass / Buffalograss / Centipedegrass / Fescue, tall / St. Augustinegrass / Zoysiagrass | crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed evening primrose hop clover | Residential and Sod Farm Turf Uses Only² 175-233 lbs/acre in the spring before weed germination. | Apply a repeat application of 116-175 lbs/acre after 5-8 weeks if necessary. |
| | | goosegrass | Golf Course, Commercial and Other Non-Residential Turf Sites 175-349 lbs/acre in the spring before weed germination. | |
| | | | All Turf Uses (Non-Greens and Tees) 175 lbs/acre before weed germination in the spring. Apply a repeat application of 175-233 lbs/acre after 5-8 weeks. | An additional application of 175 lbs/acre may be applied for extended goosegrass control 8 weeks after the second application. |
| | | cudweed <i>Poa annua</i> chickweed lawn burweed henbit corn speedwell | All Turf Uses 175-233 lbs/acre | Apply in late summer or early fall before weed germination. Apply a repeat application of 175-233 lbs/acre after 5-8 weeks for extended <i>Poa annua</i> control. |

¹ **DO NOT** exceed a maximum rate of 233 pounds **per acre per application** for use on residential turfgrass and sod farm turfgrass.
DO NOT exceed a maximum rate of 349 pounds **per acre per application** for use on golf course turfgrass, commercial or other non-residential turfgrass sites.

² Residential is defined as turf in any residential situation as well as home lawns, schools, parks and playgrounds.

³ **Not for use** on bentgrass or *Poa annua* greens or tees.

TURFGRASS PRECAUTIONS

Use on well-established turf grass with a dense and uniform stand. On newly planted areas, application should not be made until turfgrass has filled in and been mowed at least 4 times. On turf thinned or damaged because of winter injury, excessive moisture, etc., allow for turf recovery before making an application.

Applications made to overseeded warm-season turfgrass may cause thinning or injury of the overseeded species.

DO NOT use on bentgrass or *Poa annua* greens and tees or injury may occur.

Delay reseeding or winter overseeding of treated turfgrass for at least three (3) months following the last **Pendimethalin 0.86% + Fertilizer** application. Delay springing turfgrass for five (5) months after application.

WEEDS CONTROLLED

PENDIMETHALIN 0.86% + FERTILIZER provides preemergence control of weed species listed. Applications can be made around and over the top of ornamentals and to sites listed on this label.

| Grasses Controlled | | Broadleaf Weeds Controlled | |
|--------------------------|---------------------------------|----------------------------|---------------------------------|
| Common Name | Scientific Name | Common Name | Scientific Name |
| Barnyardgrass | <i>Echinochloa crus-galli</i> | Burweed, Lawn | <i>Soliva pterosperma</i> |
| Bluegrass, Annual | <i>Poa annua</i> | Carpetweed | <i>Mollugo verticillata</i> |
| Crabgrass | <i>Digitaria spp.</i> | Chickweed, Common | <i>Stellaria media</i> |
| Crowfootgrass | <i>Dactyloctenium aegyptium</i> | Chickweed, Mouseear | <i>Cerastium vulgatum</i> |
| Foxtail, Giant | <i>Setaria faberi</i> | Clover, Hop | <i>Trifolium procumbens</i> |
| Foxtail, Green | <i>Setaria viridis</i> | Cudweed | <i>Gnaphalium spp.</i> |
| Foxtail, Yellow | <i>Setaria glauca</i> | Evening primrose | <i>Oenothera biennis</i> |
| Goosegrass | <i>Eleusine indica</i> | Fiddleneck | <i>Amsinckia intermedia</i> |
| Itchgrass | <i>Rottboellia exaltata</i> | Filaree | <i>Erodium spp.</i> |
| Johnsongrass (from seed) | <i>Sorghum halepense</i> | Henbit | <i>Lamium amplexicaule</i> |
| Junglerice | <i>Echinochloa colona</i> | Knotweed, prostrate | <i>Polygonum aviculare</i> |
| Lovegrass (from seed) | <i>Eragrostis spp.</i> | Kochia | <i>Kochia scoparia</i> |
| Panicum, Browntop | <i>Panicum fasciculatum</i> | Lambsquarters | <i>Chenopodium album</i> |
| Panicum, Fall | <i>Panicum dichotomiflorum</i> | Pigweed | <i>Amaranthus spp.</i> |
| Panicum, Texas | <i>Panicum texanum</i> | Puncturevine | <i>Tribulus terrestris</i> |
| Sandbur, Field | <i>Cenchrus incertus</i> | Purslane | <i>Portulaca oleracea</i> |
| Signalgrass | <i>Bracharia platyphylla</i> | Pusley, Florida | <i>Richardia scabra</i> |
| Sprangletop, Mexican | <i>Leptochloa uninervia</i> | Rocket, London | <i>Sisymbrium irio</i> |
| Sprangletop, Red | <i>Leptochloa filiformis</i> | Shepherdspurge | <i>Capsella bursa-pastoris</i> |
| Witchgrass | <i>Panicum capillare</i> | Smartweed, Pennsylvania | <i>Polygonum pennsylvanicum</i> |
| Woolly Cupgrass | <i>Eriochloa villosa</i> | Speedwell, Corn | <i>Veronica arvensis</i> |
| | | Spurge, Annual | <i>Euphorbia spp.</i> |
| | | Spurge, Prostrate | <i>Chamaesyce humistrata</i> |
| | | Woodsorrel, Yellow | <i>Oxalis stricta</i> |
| | | Velvetleaf (Buttonweed) | <i>Abutilon theophrasti</i> |

INDUSTRIAL (UNIMPROVED) TURF

Industrial or unimproved turf areas often have a different spectrum of weeds to control than those found in fine turf. **PENDIMETHALIN 0.86% + FERTILIZER** will control annual grasses and broadleaf weeds listed for cool-season and warm-season grass, and the following weeds that might germinate in established grass in rights-of-way, roadsides, construction sites, parks, substations, lots, or similar areas:

crowfootgrass, itchgrass, Johnsongrass (from seed), Junglerice, lovegrass (from seed), browntop panicum, Texas panicum, field sandbur, signalgrass, Mexican sprangletop, red sprangletop, witchgrass, woolly cupgrass, carpetweed, fiddleneck, filaree, kochia, lambsquarters, pigweed, puncturevine, Florida pusley, London rocket, shepherdspurge, Pennsylvania smartweed, annual spurge, and velvetleaf.

Apply before weeds germinate. Use a postemergence herbicide such as 2,4-D, MCPA; MSMA; or similar product to control established weeds.

LANDSCAPE AND GROUNDS MAINTENANCE

PENDIMETHALIN 0.86% + FERTILIZER can be incorporated into landscape and grounds maintenance programs to provide extended preemergence control of most annual grasses and certain broadleaf weeds. Areas to be treated include but are not limited to established trees or ornamentals in the landscape or in mulch beds; or other commercially maintained landscape areas such as borders, around statuary or monuments in parks or in parking areas. If emerged weeds are present at time of application, either **shallow** cultivate to destroy existing weeds or, where practical, remove by hand or apply a postemergence product labeled for such use before an application of **PENDIMETHALIN 0.86% + FERTILIZER**. **PENDIMETHALIN 0.86% + FERTILIZER** may be used before or after application of herbicides registered for postemergence use (i.e. **Segment® herbicide**, **Roundup PRO® herbicide** or **Finale® herbicide**) for the control of established weeds.

DO NOT apply sprays containing **Roundup® herbicide** or **Finale** over the top of desirable plants, and follow label restrictions. A **PENDIMETHALIN 0.86% + FERTILIZER** treatment may be followed by any registered herbicide to control weeds not listed on the **PENDIMETHALIN 0.86% + FERTILIZER** label.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.
TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Pendulum and Segment are registered trademarks of BASF.

Acclaim Extra and Finale are registered trademarks of Bayer Environmental Sciences.

Trimec is a trademark of PBI Gordon Corporation.

Three-Way is a trademark of LESCO Inc.

Roundup and Roundup PRO are registered trademarks of Monsanto Company.

TCS GrowStar® is a registered trademark of Turf Care Supply Corp.

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website url: <http://www.aapfco.org/metals.htm>, or by calling 1-877-220-1014.

NVA 2013-04-123-0057

Net Weight: 50 lbs. (22.67 kg)

EPA REG. NO. 241-424-82757

EPA EST. NO. 82757-OH-001 (MF)

EPA EST. NO. 82757-MA-002 (H)

EPA EST. NO. 82757-FL-001 (S)

(First Letter of Lot # Indicates Manufacturing Site)

#900133

F1612

Distributed by:

Turf Care Supply Corp.

50 Pearl Road, Suite 200

Brunswick, OH 44212

