



US 20040186166A1

EXHIBIT  
117

(19) **United States**

(12) **Patent Application Publication**  
Burstein et al.

(10) **Pub. No.: US 2004/0186166 A1**

(43) **Pub. Date: Sep. 23, 2004**

(54) **CANNABINOID ANALOGS AS PEROXISOME  
PROLIFERATOR ACTIVATED NUCLEAR  
RECEPTOR GAMMA ACTIVATORS**

**Related U.S. Application Data**

(60) Provisional application No. 60/435,184, filed on Dec. 19, 2002.

(76) **Inventors: Sumner H. Burstein, Framingham, MA  
(US); J. Don Chen, New Brunswick,  
NJ (US); Robert B. Zurier, Princeton,  
MA (US)**

**Publication Classification**

(51) **Int. Cl.<sup>7</sup> ..... A61K 31/353**  
(52) **U.S. Cl. .... 514/454**

Correspondence Address:  
**FISH & RICHARDSON PC  
225 FRANKLIN ST  
BOSTON, MA 02110 (US)**

**ABSTRACT**

The present invention provides for methods and compositions for using cannabinoid compounds, such as ajulemic acid, for the treatment of disorders involving peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ). In particular, the invention provides for methods and compositions for treating disorders, such as autoimmune or inflammatory disorders, involving PPAR $\gamma$ , such as rheumatoid arthritis and diabetes.

(21) **Appl. No.: 10/741,219**

(22) **Filed: Dec. 19, 2003**

FILED IN OFFICE  
HART SUPERIOR COURT  
2019 OCT -2 AM 10:06  
*Frankie H. Gray, Clerk*  
FRANKIE H. GRAY, CLERK




FIG. 1

FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 3A




FIG. 3B




FIG. 3C





FIG. 3D




FIG. 3E




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 5A





FIG. 5B





Solvent FIG. 7A



AJA FIG. 7B



GW347845 FIG. 7C




FIG. 7D



**CANNABINOID ANALOGS AS PEROXISOME  
PROLIFERATOR ACTIVATED NUCLEAR  
RECEPTOR GAMMA ACTIVATORS**

**CROSS REFERENCE TO RELATED  
APPLICATIONS**

[0001] This application claims priority from U.S. Provisional Patent Application No. 60/435,184, filed on Dec. 19, 2002, which is incorporated herein by reference in its entirety.

**STATEMENT AS TO FEDERALLY SPONSORED  
RESEARCH**

[0002] This invention was made with support from grant numbers DA12178, AR38501, DK52542 and DK52888 from the National Institute of Health. The government has certain rights in the invention.

**FIELD OF THE INVENTION**

[0003] This application relates to autoimmune agents, and, in particular, to the use of cannabinoids for the treatment of disorders, such as autoimmune and inflammatory disorders, such as diabetes mellitus, and to medicinal preparations containing cannabinoids such as ajulemic acid.

**BACKGROUND OF THE INVENTION**

[0004] *Cannabis sativa* contains a group of biosynthetically related substances known collectively as cannabinoids. (F Grotenhermen and E Russo in "Cannabis and Cannabinoids", Haworth Press, New York (2002) pages xxvii-xxxi). The identification of an endogenous cannabinoid anandamide (Devane, *Trends Pharmacol Sci* 15: 40-1, 1994), a metabolite of the arachidonic acid that, among other actions, possesses anti-inflammatory activity, has generated a new wave of interest in this area. Tetrahydrocannabinol (THC), one of the major cannabinoids in Cannabis, is known to possess a number of activities such as analgesia and anti-inflammatory action. However, it also has certain undesirable effects on mood and behavior, the so-called psychotropic effects, which limits the clinical application of THC and Cannabis. Considerable effort has been expended toward the goal of discovering non-psychotropic cannabinoid analogs or derivatives that retain certain beneficial actions, but are free of psychotropic activity.

[0005] A useful template molecule is the cannabinoid, tetrahydrocannabinol-11-oic (THC-11-oic) acid, a metabolite of THC, because it does not show changes in mood or behavior in human or in animal models. However, it has only modest analgesic and anti-inflammatory activity (Burstein et al., *J Med Chem* 35:3135-41, 1992). It is known that manipulation of the side-chain structure of THC could produce significant modification of its activities. In particular, increasing the side-chain to seven carbon atoms and the introduction of methyl groups dramatically increases its potency. Based on the same analogy, manipulation of the side chain of THC-11-oic acid resulted in the production of ajulemic acid (AJA), a synthetic analog of THC-11-oic acid (FIG. 1; Burstein et al., *J Med Chem* 35:3135-41, 1992).

[0006] AJA has been shown to possess prolonged analgesic activity (Burstein et al., *J Med Chem* 35:3135-41, 1992; Dajani et al., *J Pharmacol Exp Ther* 291:31-8, 1999), and anti-inflammatory activity (Burstein et al., *J Med Chem* 35:3135-41, 1992; Zurier et al., *Arthritis Rheum* 41:163-70, 1998). In contrast to the non-steroidal anti-inflammatory drugs (NSAIDs), it is non-ulcerogenic at therapeutically relevant doses. Most importantly it shows none of the typical psychotropic actions of THC in animal models, as well as in humans.


[0007] Peroxisome proliferator-activated receptors (PPARs) are transducer proteins belonging to the steroid/thyroid/retinoid receptor superfamily. Three subtypes of PPAR have been identified, designated as PPAR-alpha (PPAR- $\alpha$ ), PPAR-beta (PPAR- $\beta$ , or PPAR-delta (PPAR- $\delta$ )), and PPAR-gamma (PPAR- $\gamma$ ). PPAR $\gamma$  plays a role in adipocyte differentiation, lipid metabolism, and glucose homeostasis, as well as in modulating anti-inflammatory (Jiang et al., *Nature* 391:82-6, 1998), and anti-tumor processes (Patel et al., *Curr Biol* 11:764-8, 2001). Activation of PPAR $\gamma$  can inhibit the expression of cytokines, such as interleukin-1 $\beta$  (IL-1 $\beta$ ), tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), nitric oxide (NO), at both protein and transcription levels (Jiang et al., *Nature* 391:82-86, 1998; Ricote et al., *Nature* 391:79-82, 1998). PPAR $\gamma$  agonist is expressed in adipose tissue, skeletal muscle, adrenal gland, colonic epithelium, heart, pancreas, and liver (Mukherjee et al., *J Biol Chem* 272:8071-6, 1997; Sarraf et al., *Nat Med* 4:1046-52, 1998). It is also expressed in immune system related cells such as splenocytes (Clark et al., *J Immunol* 164:1364-71, 2000; Kliewer et al., *Proc Natl Acad Sci USA* 91:7355-9, 1994), synoviocytes (Ji et al., *J Autoimmun* 17:215-21, 2001); Kawahito et al., *J Clin Invest* 106:189-97, 2000; Simonin et al., *Am J Physiol Cell Physiol* 282:C125-33, 2002), helper T cells (Clark et al., *J Immunol* 164:1364-71, 2000) and activated monocytes and macrophages (Jiang et al., *Nature* 391:82-6, 1998; Kawahito et al., *J Clin Invest* 106:189-97, 2000; Ricote et al., *Nature* 391:79-82, 1998).

**SUMMARY OF THE INVENTION**

[0008] The present invention relates to THC cannabinoid compounds, e.g., ajulemic acid (AJA, dimethylheptyl-THC-11-oic acid), pharmaceutical compositions containing them, and methods for using the cannabinoids, e.g., ajulemic acid, as medicaments. More specifically, cannabinoids, e.g., ajulemic acid, of the invention can be utilized in the modification, amelioration, reduction, or prevention of disorders associated with peroxisome proliferator-activated receptors (PPAR), in particular peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ), and are useful for the treatment of such disorders, including autoimmune, inflammatory, and other disorders, such as diabetes and rheumatoid arthritis.

[0009] The present invention is based, in part, on the discovery that autoimmune, inflammatory, and other disorders can be treated or ameliorated by using cannabinoids that effect PPAR activity, e.g., by using a cannabinoid with Formula I (see below) to bind to, and activate PPAR $\gamma$ . Accordingly, in one aspect, the invention pertains to a method for treating a disorder associated with peroxisome proliferator-activated receptor (PPAR) function by administering to a subject a pharmaceutically effective amount of a composition including a cannabinoid compound. The can-

nabinoid can be combined with an existing anti-inflammatory agent, or anti-diabetic agent. In one embodiment, the cannabinoid is that with "Formula I", e.g., ajulemic acid:



(Formula I)

[0010] in which R<sup>1</sup> can be a hydrogen atom, —COCH<sub>3</sub>, —COCH<sub>2</sub>CH<sub>3</sub>, or —CO(CH<sub>2</sub>)<sub>N</sub>CH<sub>3</sub> (where N=0-20) and R<sup>2</sup> can be a branched C<sub>5</sub>-C<sub>12</sub> alkyl. In certain embodiments, R<sup>1</sup> can be hydrogen and/or R<sup>2</sup> can be a C<sub>9</sub> alkyl, e.g., a branched alkyl, such as 1,1-dimethylheptyl. R<sub>3</sub> can be —OH, —OCH<sub>3</sub>, or —NHCH<sub>2</sub>COOH. Other cannabinoids include all of those listed in U.S. Pat. Nos. 4,847,290, 4,973,603, 5,338,753, 5,538,993, 5,635,530, 6,162,829, and 6,448,288, which are all hereby incorporated by reference in their entireties.

[0011] The cannabinoid can be used to ameliorate or treat a disorder associated with a PPAR $\gamma$  (e.g., an autoimmune disorder (e.g., diabetes mellitus, impaired glucose intolerance, diabetic retinopathy, obesity, systemic lupus erythematosus, rheumatoid arthritis, spondylo arthritis, asthma, inflammatory bowel disease, vasculitis, dermatomyositis, polymyositis, sjogren's syndrome, hyperthyroidism), ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis, osteoporosis, atherosclerosis, hypertension, hyperglycemia, coronary artery disease, dyslipidemia, exudative age-related macular degeneration, areolar age-related macular degeneration, atherosclerosis, an inflammatory disorder (e.g., rheumatoid arthritis, multiple sclerosis, myasthenia gravis, uveoretinitis, uveitis, iritis, cyclitis, chorioiditis, chorioretinitis, vitritis, keratitis, conjunctivitis, psoriasis, eczema, thyroiditis, a collagen vascular disorder (e.g., ankylosing spondylitis, rheumatoid arthritis, lupus erythematosus, Reiter syndrome, Behcet disease, ulcerative colitis, Crohn's disease, or Wegener's granulomatosis))). For example, the disorder can also be a non-autoimmune inflammatory disorder, such as osteoarthritis. In certain embodiments, for example, the disorder can be diabetes, which can be treated by administering to a subject with diabetes a pharmaceutically effective amount of cannabinoid of Formula I, e.g., ajulemic acid. As a further example, the disorder can be an inflammatory disease associated with PPAR $\gamma$ , such as rheumatoid arthritis, multiple sclerosis, myasthenia gravis, or uveoretinitis.

[0012] In another aspect, the invention features methods of modifying or treating an autoimmune disorder associated with peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) function, including administering to a subject in need thereof an amount of a PPAR $\gamma$  activator effective to modify or ameliorate an autoimmune disorder, wherein the PPAR $\gamma$  activator is a cannabinoid. In certain embodiments, the autoimmune disease is diabetes and can be treated in a subject sensitive to the cannabinoid of Formula I by identifying a subject having diabetes sensitive to Formula I, and administering to the subject an effective amount of a com-

pound of Formula I. Formula I can be administered in a composition orally, systemically, via an implant, for example an implant that provides slow release of the compound, or may also be administered intravenously. The compound can be administered at about 0.1 to 50 mg/kg body weight of the subject, e.g., a mammalian subject, and preferably about 0.2 to 2 mg/kg body weight of the mammalian subject.

[0013] In another aspect, the invention features methods of treating a subject (e.g., one sensitive to a cannabinoid) having a disorder (e.g., an autoimmune disorder (e.g., diabetes mellitus, impaired glucose intolerance, diabetic retinopathy, obesity, systemic lupus erythematosus, rheumatoid arthritis, spondylo arthritis, asthma, inflammatory bowel disease, vasculitis, dermatomyositis, polymyositis, sjogren's syndrome, hyperthyroidism), ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis, osteoporosis, atherosclerosis, hypertension, hyperglycemia, coronary artery disease, dyslipidemia, exudative age-related macular degeneration, areolar age-related macular degeneration, atherosclerosis, an inflammatory disorder (e.g., rheumatoid arthritis, multiple sclerosis, myasthenia gravis, uveoretinitis, uveitis, iritis, cyclitis, chorioiditis, chorioretinitis, vitritis, keratitis, conjunctivitis, psoriasis, eczema, thyroiditis, a collagen vascular disorder (e.g., ankylosing spondylitis, rheumatoid arthritis, lupus erythematosus, Reiter syndrome, Behcet disease, ulcerative colitis, Crohn's disease, or Wegener's granulomatosis))) associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, in which the method includes administering to the subject a pharmaceutically effective amount of a composition including a cannabinoid compound (e.g., a cannabinoid of Formula I (e.g., in which R<sup>2</sup> is a C<sub>9</sub> alkyl, a branched alkyl, 1,1-dimethylheptyl), ajulemic acid). In these methods, the dosage of the cannabinoid can be between about 0.1 and 50 mg/kg body weight of the subject, and the mode of administration of the cannabinoid can be oral, nasal, pulmonary, transdermal, or parenteral. The subject can be a human or an animal (e.g., a mammal (e.g., non-human primate, dog, cat, rodent, horse, cow, sheep, or goat)).

[0014] In another aspect, the invention features kits comprising a pharmaceutically effective amount of a composition including a cannabinoid compound (e.g., a cannabinoid of Formula I (e.g., in which R<sup>2</sup> is a C<sub>9</sub> alkyl, a branched alkyl, 1,1-dimethylheptyl), ajulemic acid), as well as instructions for use of the composition in treating a subject (e.g., one sensitive to a cannabinoid) having a disorder (e.g., an autoimmune disorder (e.g., diabetes mellitus, impaired glucose intolerance, diabetic retinopathy, obesity, systemic lupus erythematosus, rheumatoid arthritis, spondylo arthritis, asthma, inflammatory bowel disease, vasculitis, dermatomyositis, polymyositis, sjogren's syndrome, hyperthyroidism), ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis, osteoporosis, atherosclerosis, hypertension, hyperglycemia, coronary artery disease, dyslipidemia, exudative age-related macular degeneration, areolar age-related macular degeneration, atherosclerosis, an inflammatory disorder (e.g., rheumatoid arthritis, multiple sclerosis, myasthenia gravis, uveoretinitis, uveitis, iritis, cyclitis, chorioiditis, chorioretinitis, vitritis, keratitis, conjunctivitis, psoriasis, eczema, thyroiditis, a collagen vascular disorder (e.g., ankylosing spondylitis, rheumatoid arthritis, lupus erythematosus, Reiter syndrome, Behcet disease, ulcerative colitis,

Crohn's disease, or Wegener's granulomatosis))) associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function.

[0015] The invention also includes compositions for use as medicaments in treating a subject (e.g., one sensitive to a cannabinoid) having a disorder (e.g., an autoimmune disorder (e.g., diabetes mellitus, impaired glucose intolerance, diabetic retinopathy, obesity, systemic lupus erythematosus, rheumatoid arthritis, spondylo arthritis, asthma, inflammatory bowel disease, vasculitis, dermatomyositis, polymyositis, sjogren's syndrome, hyperthyroidism), ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis, osteoporosis, atherosclerosis, hypertension, hyperglycemia, coronary artery disease, dyslipidemia, exudative age-related macular degeneration, areolar age-related macular degeneration, atherosclerosis, an inflammatory disorder (e.g., rheumatoid arthritis, multiple sclerosis, myasthenia gravis, uveoretinitis, uveitis, iritis, cyclitis, choroiditis, chorioretinitis, vitritis, keratitis, conjunctivitis, psoriasis, eczema, thyroiditis, a collagen vascular disorder (e.g., ankylosing spondylitis, rheumatoid arthritis, lupus erythematosus, Reiter syndrome, Bechet disease, ulcerative colitis, Crohn's disease, or Wegener's granulomatosis))) associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, in which the compositions include a pharmaceutically effective amount of a cannabinoid compound or PPAR $\gamma$  activator (e.g., a cannabinoid of Formula I (e.g., in which R<sup>2</sup> is a C<sub>9</sub> alkyl, a branched alkyl, 1,1-dimethylheptyl), ajulemic acid). With respect to these compositions, the dosage of the cannabinoid can be between about 0.1 and 50 mg/kg body weight of the subject, and the mode of administration of the cannabinoid can be oral, nasal, pulmonary, transdermal, or parenteral. The subject can be a human or an animal (e.g., a mammal (e.g., non-human primate, dog, cat, rodent, horse, cow, sheep, or goat)).

[0016] In addition, the invention encompasses uses of compositions for the manufacture of a medicament for use in treating a subject (e.g., one sensitive to a cannabinoid) having a disorder (e.g., an autoimmune disorder (e.g., diabetes mellitus, impaired glucose intolerance, diabetic retinopathy, obesity, systemic lupus erythematosus, rheumatoid arthritis, spondylo arthritis, asthma, inflammatory bowel disease, vasculitis, dermatomyositis, polymyositis, sjogren's syndrome, hyperthyroidism), ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis, osteoporosis, atherosclerosis, hypertension, hyperglycemia, coronary artery disease, dyslipidemia, exudative age-related macular degeneration, areolar age-related macular degeneration, atherosclerosis, an inflammatory disorder (e.g., rheumatoid arthritis, multiple sclerosis, myasthenia gravis, uveoretinitis, uveitis, iritis, cyclitis, choroiditis, chorioretinitis, vitritis, keratitis, conjunctivitis, psoriasis, eczema, thyroiditis, a collagen vascular disorder (e.g., ankylosing spondylitis, rheumatoid arthritis, lupus erythematosus, Reiter syndrome, Bechet disease, ulcerative colitis, Crohn's disease, or Wegener's granulomatosis))) associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, in which the compositions include a pharmaceutically effective amount of a cannabinoid compound or PPAR $\gamma$  activator (e.g., a cannabinoid of Formula I (e.g., in which R<sup>2</sup> is a C<sub>9</sub> alkyl, a branched alkyl, 1,1-dimethylheptyl), ajulemic acid). In these uses, the dosage of the cannabinoid can be between about 0.1 and 50 mg/kg body weight of the subject, and the

mode of administration of the cannabinoid can be oral, nasal, pulmonary, transdermal, or parenteral. The subject can be a human or an animal (e.g., a mammal (e.g., non-human primate, dog, cat, rodent, horse, cow, sheep, or goat)).

[0017] The term "subject" as used herein refers to any living organism in which an immune response is elicited. The term subject includes, but is not limited to, humans, nonhuman primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs, and the like. The term does not denote a particular age or sex. Thus, adult and newborn subjects, as well as fetuses, whether male or female, are intended to be covered.

[0018] The term "sample" as used herein refers to a test item that has a component that is capable of interacting with a cannabinoid, e.g., a biological sample with PPAR $\gamma$  that can interact with AJA. The sample can be a liquid or fluid biological sample, or a solid biological sample. The biological sample can be a liquid sample e.g., blood, plasma, serum, cerebral spinal fluid, urine, amniotic fluid, interstitial fluid, and synovial fluid. The sample may be a solid, e.g., a tissue or cell matter. The term "sample" also refers to a non-biological sample such as a chemical solution, or synthetic composition. In one embodiment, the sample is blood. In another embodiment, the sample is plasma.

[0019] The term "peroxisome proliferator-activated receptor" and "PPAR" are art recognized term referring to a family of transducer proteins belonging to the steroid/thyroid/retinoid receptor superfamily. PPARs are described in a review article by Schoonjans (1996) *J. Lipid Res.* 37:907-925, and described in more detail below. Three subtypes of PPAR have been identified, and these are designated as alpha ( $\alpha$ ), beta ( $\beta$ ) (or delta ( $\delta$ )), and gamma ( $\gamma$ ). In a preferred embodiment, cannabinoids, such as AJA bind to, and activate PPAR $\gamma$ .

[0020] The term "PPAR $\gamma$ " as used herein refers to all isotypes of PPAR $\gamma$ . PPAR $\gamma$  exists as at least two isotypes, PPAR $\gamma$ 1 and PPAR $\gamma$ 2. The term "PPAR $\gamma$ " refers to any of these isotypes or combination thereof.

[0021] The phrase "PPAR $\gamma$  associated disorder" as used herein refers to a pathological condition in a subject that results at least, in part, from PPAR $\gamma$  function. The phrase "PPAR $\gamma$  associated disorder" is also intended to include any disorders in which the manifestation of the disorder is characterized by the disturbance in the regulation of mood, behavior, control of feeding behavior. Particularly preferred PPAR $\gamma$  associated disorder include autoimmune disorders that involve PPAR $\gamma$ . Examples of autoimmune diseases include, but are not limited to, rheumatoid arthritis, diabetes mellitus, glucose intolerance.

[0022] The term "modifies" or "modified" are used interchangeably herein and refer to the up-regulation, or down-regulation, or activation of the target PPAR, e.g., PPAR $\gamma$ . The term "modifies" or "modified" also refers to the increase, decrease, elevation, or depression of processes or signal transduction cascades involving a target PPAR, e.g., PPAR $\gamma$ . Modification to the PPAR $\gamma$  may occur when a cannabinoid, e.g., AJA, binds to the PPAR $\gamma$ . This modifica-

tion may directly affect the PPAR $\gamma$ , for example modifications that may result in an increase in PPAR $\gamma$  number. Alternatively, the modifications may occur as an indirect effect of binding to the PPAR $\gamma$ . For example, binding of AJA to the PPAR $\gamma$  can also lead to a change in downstream processes involving the PPAR $\gamma$ , such as the recruitment of co-activators by the PPAR $\gamma$ -AJA complex. In addition, antagonism of transcription factors can facilitate gene expression of mediators of inflammation, thereby reducing production of the mediators. The modifications can therefore be direct modifications of the PPAR $\gamma$ , or an indirect modification of a process or cascade involving the PPAR $\gamma$ . Non-limiting examples of modifications includes modifications of morphological and functional processes, under- or over production or expression of substances, induction of differentiation of cells, e.g., induction of adipocyte differentiation, and recruitment of co-activators, e.g., DRIP205.

[0023] The terms “modify” and “modified” also include treating a subject prophylactically to alter inflammation, apoptosis, proliferation, autoimmune function, and expression of oncogenes and other genes controlling cell metabolism. The present methods include both medical therapeutic and/or prophylactic treatment, as necessary.

[0024] The term “cannabinoid” as used herein refers to biosynthetically related compounds such as delta-8-tetrahydrocannabinol, delta-9-tetrahydrocannabinol, cannabidiol, olivetol, cannabinol, cannabigerol, nabilone, delta-9-tetrahydrocannabinol-11 oic acid. The non-psychoactive cannabinoid 3-dimethylheptyl 11 carboxylic acid homolog of, delta-8-tetrahydrocannabinol (Burstein et al. *J. Med. Chem.* (1992) 35: 3135). A useful cannabinoid is ajulemic acid (AJA; dimethylheptyl-THC-11-oic acid). The term “cannabinoid” also includes the cannabinoid compounds, their derivatives, analogs, tautomeric forms, stereoisomers, polymorphs, metabolic analogues, pharmaceutically acceptable salts, pharmaceutically acceptable solvates, and pharmaceutical compositions containing them. The compounds are useful in the treatment of diabetes mellitus (Type 1 and Type II), impaired glucose tolerance, insulin resistance, obesity, and other diseases. The AJA acting through PPAR $\gamma$  can also be used to treat or modify autoimmune diseases such as rheumatoid arthritis.

[0025] The term “activator” is used to denote any molecular species that results in activation of a PPAR, e.g., PPAR $\gamma$  regardless of whether the species itself binds to the receptor or a metabolite of the species binds to the receptor.

[0026] The term “PPAR $\gamma$  activator” as used herein refers to a molecular species that activates PPAR-gamma.

[0027] The term “alkyl” as used herein refers to a straight or branched hydrocarbon chain containing carbon atoms or cyclic hydrocarbon moieties. These alkyl groups may also contain one or more double bonds or triple bonds. By “substituted alkyl” is meant an alkyl in which an atom of the alkyl is substituted with an atom, e.g., a sulfur, oxygen, or halogen atom.

[0028] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods

and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

[0029] Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0030] FIG. 1 is a series of representations of the chemical structures of THC, THC-11-oic acid, and ajulemic acid.

[0031] FIG. 2A is a representation of a membrane showing partial protease digestion assays indicating that AJA binds to PPAR $\gamma$ . AJA induces PPAR $\gamma$  to a conformation that is more resistant to trypsin digestion. AJA treatment protects mPPAR $\gamma$  from trypsin digestion.

[0032] FIG. 2B is a representation of a membrane showing partial protease digestion assays indicating that AJA binds to PPAR $\gamma$ . The resistance to trypsin digestion was positively related to the concentration of AJA in a dose response manner.

[0033] FIG. 2C is a representation of a membrane showing partial protease digestion assays in which AJA provides concentration-dependent protection of mPPAR $\gamma$  against trypsin digestion.

[0034] FIG. 2D is a representation of a membrane showing partial protease digestion assays indicating that PPAR $\delta$  was digested evenly when treated with different concentrations of AJA.

[0035] FIG. 3A is a bar graph showing the activation of PPAR $\gamma$  by GW347845 (GW34) and AJA in a concentration-dependent manner.

[0036] FIG. 3B is a bar graph showing that PPAR $\alpha$  was not activated by 20  $\mu$ M of AJA in a reporter gene assay.

[0037] FIG. 3C is a bar graph showing that PPAR $\delta$  was not activated by 20  $\mu$ M of AJA in a reporter gene assay.

[0038] FIG. 3D is a bar graph showing that AJA does not activate RXR $\alpha$ . Only 9-cis RA activated RXR $\alpha$ .

[0039] FIG. 3E is a bar graph showing that the AF-2 helix truncated PPAR $\gamma$  cannot be activated by AJA or GW34. The activation of PPAR $\gamma$  requires AF-2 helix of the receptor.

[0040] FIG. 4A is a schematic diagram of the GAL4 based reporter system showing that AJA activates PPAR $\gamma$  in a heterologous reporter system. The PPAR $\gamma$  was expressed as a Gal4 DBD fusion protein, which binds to the UAS promoter containing 4 copies of the GAL4 binding sites upstream of the minimal TK promoter. Binding of PPAR $\gamma$  activators, such as AJA, activates the luciferase reporter gene expression.

[0041] FIG. 4B is a bar graph showing that PPAR $\gamma$  was significantly activated by 20  $\mu$ M of AJA or 1  $\mu$ M of GW34 that served as a positive control.

[0042] FIG. 4C is a graph showing the dose-dependent activation of PPAR $\gamma$  by AJA by measuring luciferase activity.

[0043] FIG. 4D is a graph showing the fold activation of human PPAR $\gamma$  by AJA, while AJA failed to activate PPAR $\alpha$  and PPAR $\delta$ , and PPAR $\gamma$ .

[0044] FIG. 5A is a photograph of a membrane showing the interaction of PPAR $\gamma$  with coactivator GST-DRIP205 (amino acids 527-970) in the presence of GW34 or AJA.

[0045] FIG. 5B is a photograph of a membrane showing the interaction of PPAR $\gamma$  with coactivator GST-RAC3 (amino acids 613-752) in the presence of GW34 or AJA.

[0046] FIG. 6A is a bar graph of PPAR $\gamma$  transfected cells in which AJA reduces the PMA-activated IL-8 promoter activity in a concentration (in  $\mu$ M)-dependent manner.

[0047] FIG. 6B is a bar graph showing that AJA had no effect on IL-8 promoter activity in the PPAR $\gamma$  $\Delta$ AF2 transfected cells.

[0048] FIG. 6C is a bar graph showing that GW347845 reduces the IL-8 promoter activity in cells transfected with wild type PPAR $\gamma$ .

[0049] FIG. 6D is a bar graph showing that GW347845 had no effect on IL-8 promoter activity in cells transfected with the PPAR $\gamma$  $\Delta$ AF2 mutant.

[0050] FIGS. 7A-C are representations of 3T3 L1 cells. FIG. 7A depicts treatment with 0.1% DMSO control (vehicle). FIG. 7B depicts treatment with 1  $\mu$ M of GW347845 in 0.1% DMSO (GW34). FIG. 7C depicts treatment with 20  $\mu$ M of AJA in 0.1% DMSO (AJA). The results demonstrate that AJA induces adipocyte differentiation in 3T3 L1 cells; and

[0051] FIG. 7D is a representation of a gel showing RT-PCR analysis of adipocyte specific genes.

#### DETAILED DESCRIPTION OF THE INVENTION

[0052] The practice of the present invention employs, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature (see, e.g., Sambrook, et al. *Molecular Cloning: A Laboratory Manual* (Current Edition); *DNA Cloning: A Practical Approach*, Vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., Current Edition); *Nucleic Acid Hybridization* (B. Hames & S. Higgins, eds., Current Edition); *Transcription and Translation* (B. Hames & S. Higgins, eds., Current Edition); *CRC Handbooks*).

[0053] The invention is based, in part, on the discovery that the cannabinoid with Formula I, e.g., ajulemic acid (AJA, dimethylheptyl-THC-11-oic acid), a synthetic analog of THC-11-oic acid, binds directly and specifically with peroxisome proliferator activated nuclear receptor gamma (PPAR $\gamma$ ), and modifies the effect of the receptor. In particular, AJA binding to PPAR $\gamma$  produces an AJA-PPAR $\gamma$  complex that effects down stream processes, for example, recruitment of co-activators such as DRIP205, and antagonism of transcription factors which facilitate gene expression of mediators of inflammation, thereby reducing production of the mediators.

[0054] The methods and compositions of the invention can be used to target PPAR $\gamma$  for therapeutic effects of AJA on autoimmune diseases or disorders that involve PPAR $\gamma$ ,

such as diabetes, insulin resistance, glucose intolerance, NIDDM, and lipid metabolism. The identification of PPAR $\gamma$  as a selective target for AJA allows development of more effective drugs for the treatment of diabetes and obesity, as well as other autoimmune diseases involving PPAR $\gamma$ .

[0055] The invention is described in more detail in the following subsections.

[0056] I. PPAR Family In one aspect, the invention features using THC cannabinoids of Formula I, such as ajulemic acid (AJA), as activators of a peroxisome proliferator-activated receptor (PPAR) family member, in particular, the activation of PPAR $\gamma$ . The PPARs are transducer proteins belonging to the steroid/thyroid/retinoid receptor superfamily. Three subtypes of PPAR have been identified, designated as PPAR-alpha (PPAR- $\alpha$ ), PPAR-beta (PPAR- $\beta$  or PPAR-delta (PPAR- $\delta$ )), and PPAR-gamma (PPAR- $\gamma$ ). These receptors function as activator-regulated transcription factors that control the expression of target genes by binding to their responsive DNA sequence as heterodimers with retinoid x receptor (RXR).

[0057] The PPAR $\alpha$  subtype has been cloned from *Xenopus*, humans, mouse, and rat; the PPAR $\beta$  (or PPAR $\delta$ ) subtype from *Xenopus*, humans, and mouse; and the PPAR $\gamma$  subtype from *Xenopus*, humans, and hamster. These subtypes are pharmacologically distinct and differentially activated by various agents (Yu et al., *Cell* 67:1251-1266, 1991). The following are corresponding Genbank accession numbers: PPAR $\alpha$ ( $\delta$ ) (AF246303), PPAR $\beta$  (AL022721), and PPAR $\gamma$  (AY157024). PPAR $\gamma$  exists as at least two isoforms, PPAR $\gamma$ 1 and PPAR $\gamma$ 2. PPAR $\gamma$ 2 is expressed selectively in adipose tissue, whereas PPAR $\gamma$ 1 is expressed at lower levels in a variety of other rodent and human tissues (Spiegelman, *Diabetes* 47:507-514, 1998).

[0058] PPAR $\gamma$  is a pharmacologically important member of the nuclear receptor superfamily (Houseknecht et al., *Domest Anim Endocrinol* 22:1-23, 2002). It plays important roles in a diverse array of biological processes including lipid metabolism, glucose homeostasis, and adipocyte differentiation. The crystal structure of the PPAR $\gamma$  ligand-binding domain reveals a large hydrophobic cavity for ligand binding (Uppenberg et al., *J Biol Chem* 273:31108-12, 1998; and Xu et al., *Proc Natl Acad Sci USA* 98:13919-24, 2001). Indeed, PPAR $\gamma$  binds to a wide range of synthetic and naturally occurring substances, including the antidiabetic drugs thiazolidinediones (Lehmann et al., *J Biol Chem* 270:12953-6, 1995; Willson et al., *J Med Chem* 39:665-8, 1996), the synthetic tyrosine analog GW347845 (Cobb et al., *J Med Chem* 41:5055-5069, 1998), polyunsaturated fatty acids (Kliwer et al., *Proc Natl Acad Sci USA* 94:4318-23, 1997), metabolites of arachidonic acid including 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (Forman et al., *Cell* 83:803-12, 1995; Kliwer et al., *Cell* 83:813-9, 1995), NSAIDs (Lehmann et al., *J Biol Chem* 272:3406-10, 1997), and compounds of oxidized low-density lipoprotein, such as 13-hydroxyoctadecadienoic acid (13-HODE) and 15-hydroxyeicosatetraenoic acid (15-HETE) (Nagy et al., *Cell* 93:229-40, 1998). Several of these PPAR $\gamma$  ligands exhibit antiinflammatory activity in vivo (Kawahito et al., *J Clin Invest* 106:189-97, 2000; Naito et al., *Aliment Pharmacol Ther* 15:865-73, 2001), and activation of PPAR $\gamma$  is directly linked to antiinflammatory (Jiang et al., *Nature* 391:82-6, 1998), and antitumor processes (Patel et al., *Curr Biol* 11:764-8, 2001). Accordingly,

activation of PPAR $\gamma$  inhibits the expression of cytokines, such as interleukin (IL)-1  $\beta$ , tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), and nitric oxide (NO) at both the protein and transcription levels (Jiang et al., *Nature* 391:82-6, 1998; Ricote et al., *Nature* 391:79-82, 1998). PPAR $\gamma$  is expressed in adipose tissue, skeletal muscle, adrenal gland, colonic epithelium, heart, pancreas, and liver (Mukherjee et al., *J Biol Chem* 272:8071-6, 1997; Sarraf et al., *Nat Med* 4:1046-52, 1998). It is also expressed in immune system related cells such as splenocytes (Clark et al., *J Immunol* 164:1364-71, 2000; Kliewer et al., *Proc Natl Acad Sci USA* 91:7355-9, 1994), synoviocytes (Ji et al., *J Autoimmun* 17:215-21, 2001; Kawahito et al., *J Clin Invest* 106:189-97, 2000; Simonin et al., *Am J Physiol Cell Physiol* 282:C125-33, 2002), helper T cells (Clark et al., *J Immunol* 164:1364-71, 2000), and activated monocytes and macrophages (Jiang et al., *Nature* 391:82-6, 1998; Kawahito et al., *J Clin Invest* 106:189-97, 2000; Ricote et al., *Nature* 391:79-82, 1998) suggesting that PPAR $\gamma$  has a direct role in modulating inflammation in addition to its role in lipid metabolism and glucose homeostasis.

[0059] PPAR $\gamma$  modulates the expression of genes involved in the regulation of growth and differentiation in a variety of cell types that express the receptor. PPAR $\gamma$  has been shown to be expressed in an adipose tissue-specific manner. Its expression is induced early during the course of differentiation of several preadipocyte cell lines. PPAR $\gamma$  plays a role in the adipogenic signaling cascade and also regulates the ob/leptin gene which is involved in regulating energy homeostasis.

[0060] It was determined in the present invention that AJA is an activator of PPAR $\gamma$  and binds to PPAR $\gamma$  directly (see Examples). The data demonstrates that activation of PPAR $\gamma$  by AJA may contribute to the analgesic and anti-inflammatory actions of AJA. These data also demonstrate that AJA may have other biological functions, since PPAR $\gamma$  regulates a wide range of cellular activities.

[0061] Treatment of peritoneal macrophages with 15d-PGJ2 or several synthetic PPAR $\gamma$  activators suppressed the expression of the inducible nitric oxide synthase (iNOS) and inhibited induction of gelatinase B and scavenger receptor, a gene induced by phorbol ester stimulation. The promoters of these genes were found to possess binding sites for activator protein-1 (AP-1), nuclear factor  $\kappa$ B (NF- $\kappa$ B), and signal transducer and activator of transcription (STAT). Furthermore, the inhibition of the inflammatory response in macrophages was found in part by antagonizing the activities of these transcription factors (Ricote et al., *Nature* 391, 1998).

[0062] Because PPAR $\gamma$  forms a permissive heterodimer with RXR, activators for both receptors were shown to inhibit the LPS-induced NO and TNF- $\alpha$  production and the combined treatment with the two activators resulted in synergistic inhibition. Activation of PPAR $\gamma$ /RXR did not affect the translocation of NF- $\kappa$ B to nucleus, NF- $\kappa$ B activation in (EMSA), or the phosphorylation of JNK on LPS stimulation, while in the meantime it suppressed LPS-activated NF- $\kappa$ B promoter activity. This indicates that the inhibition was achieved at the transcription level (Uchimura et al., *Hepatology* 33:91-9, 2001). Even though the promoter regions of the iNOS and TNF- $\alpha$  contain binding sites for NF- $\kappa$ B and AP-1, they do not contain consensus PPAR

response elements, indicative that it is unlikely that PPAR $\gamma$  binds to, or controls these promoters directly. The inhibition of inflammatory response by activation of PPAR $\gamma$  was confirmed by recent studies both in vitro (Ji et al., *J Autoimmun* 17:215-21, 2001) and in vivo (Dubuquoy et al., *Gastroenterol Clin Biol* 24:719-24, 2000; Kawahito et al., *J Clin Invest* 106:189-97, 2000; Naito et al., *Aliment Pharmacol Ther* 15:865-73, 2001).

[0063] In these studies, the anti-inflammatory effects of 15d-PGJ2 were shown to be more potent than any synthetic PPAR $\gamma$  activator. It has been demonstrated that 15d-PGJ2 and troglitazone modulated the expressions of LPS-induced iNOS, COX-2, and pro-inflammatory cytokines differently. 15d-PGJ2 suppressed the expressions of iNOS, COX-2, IL-1 $\beta$  and TNF $\alpha$ . Troglitazone only inhibited the iNOS, and TNF  $\alpha$  expressions, indicating that additional mechanisms may be involved in 15d-PGJ2-mediated anti-inflammatory effect (Simonin et al., *Am J Physiol Cell Physiol* 282:C125-33, 2002).

[0064] Recent studies demonstrated that 15d-PGJ2 also mediated anti-inflammatory action through PPAR $\gamma$ -independent manner. The 15d-PGJ2 suppressed the IL-1 $\beta$ -induced PGE $_2$  synthesis by inhibiting the expressions of COX-2 and cytosolic phospholipase A $_2$  (cPLA $_2$ ), which are critical enzymes during the synthesis of prostaglandins (Tsubouchi et al., *Biochem Biophys Res Commun* 283:750-5, 2001). In resting cells, NF- $\kappa$ B is sequestered in the cytoplasm by association with an inhibitory protein I $\kappa$ B. In response to the signaling from inflammatory cytokine, the I $\kappa$ B kinase (IKK) is activated and phosphorylates I $\kappa$ B, such that it is readily digested by proteasome, allowing NF- $\kappa$ B to migrate into the nucleus and activate gene expression (Ghosh et al., *Annu Rev Immunol* 16:225-60, 1998). 15d-PGJ2 was shown to inhibit IKK (Straus et al., *Proc Natl Acad Sci USA* 97:4844-9, 2000) and directly inhibit DNA binding of NF- $\kappa$ B (Straus et al., *Proc Natl Acad Sci USA* 97:4844-9, 2000; Simonin et al., *Am J Physiol Cell Physiol* 282:C125-33, 2002).

[0065] The activator binding domain of PPAR $\gamma$  is much larger than the other nuclear receptor so that PPAR $\gamma$  appears to bind a range of synthetic and naturally occurring substances, including antidiabetic drug thiazolidinediones (Lehmann et al., *J Biol Chem* 270:12953-6, 1995; Willson et al., *J Med Chem* 39:665-8, 1996), polyunsaturated fatty acid (Kliewer et al., *Proc Natl Acad Sci USA* 94:4318-23, 1997), 15-deoxy- $\Delta^{12,14}$ prostaglandin J $_2$  (Forman et al., *Cell* 83:803-12, 1995; Kliewer et al., *Cell* 83:813-9, 1995), NSAIDs (Lehmann et al., *J Biol Chem* 272:3406-10, 1997), and compounds of oxidized low-density lipoprotein, such as 13-hydroxyoctadecadienoic acid (13-HODE) and 15-hydroxyicosatetraenoic acid (15-HETE) (Nagy et al., *Cell* 93:229-40, 1998).

## [0066] II. Cannabinoids and Cannabinoid Receptors

[0067] In one aspect, the invention pertains to using THC cannabinoids, and derivatives of cannabinoids such as ajulemic acid, for modifying a disorder associated with PPAR $\gamma$ . The term "cannabinoids" refers to organic substances present in *Cannabis sativa*, having a variety of pharmacological properties.

[0068] Ajulemic acid (AJA, dimethylheptyl-THC-11-oic acid) is a non-psychoactive, synthetic cannabinoid based on the template of THC-11-oic acid. THC is biotransformed by

oxidation to its principal metabolite, THC-11-oic acid. AJA is synthesized by increasing the side chain of THC-11-oic acid from 5 carbon atoms (pentyl) to 7 atoms (heptyl) and by introducing 2 methyl groups.

[0069] AJA has been shown to produce potent and prolonged analgesic effects (Burststein et al., *Life Sci* 63:161-8, 1998; Dajani et al., *J Pharmacol Exp Ther* 291:31-8, 1999), and anti-inflammatory function (Burststein et al., *J Med Chem* 35:3135-41, 1992; Zurier et al., *Arthritis Rheum* 41:163-70, 1998). In the mouse tail clip assay for analgesia, AJA was shown to exert a potent analgesic function comparable with morphine and the effect lasted even longer than morphin (Dajani et al., *J Pharmacol Exp Ther* 291:31-8, 1999). AJA was also shown to suppress both acute inflammation induced by injection of IL-1 $\beta$  and TNF $\alpha$  into subcutaneous air pouches in mice, and chronic joint inflammation of adjuvant-induced poly-arthritis in rats (Zurier et al., *Arthritis Rheum* 41:163-70, 1998). In contrast to NSAIDs, it was totally non-ulcerogenic at therapeutically relevant doses and did not exhibit drug dependence or cause mutagenesis (Dajani et al., *J Pharmacol Exp Ther* 291:31-8, 1999; Burststein, *Curr Pharm Des* 6:1339-45, 2000). Most importantly, AJA does not show the typical psychoactive effect in animal models (Burststein et al., *J Med Chem* 35:3135-41, 1992). The mechanism of AJA function is still not fully understood. Its stereospecificity suggested that its functions might be receptor mediated. Two receptors for cannabinoids, cannabinoid receptor-1 (CB1) and CB2, have been identified and cloned (Pertwee, *Pharmacol Ther* 74:129-80, 1997).

[0070] A recent study demonstrated that AJA exhibits modest affinity to CB2, and was found to have CB2-mediated anti-tumor effects (Recht et al., *Biochem Pharmacol* 62:755-63, 2001). This finding is consistent with the lack of psychoactivity in AJA because, unlike the CB1 receptor, which is present in brain and peripheral tissue, the CB2 receptor is expressed mainly in extraneural, primarily immune tissue.

[0071] AJA was also found to suppress COX-2 activity at 1-10  $\mu$ M concentration range, while having no effect on COX-1 activity at concentrations between 0.25-25  $\mu$ M (Zurier et al., *Arthritis Rheum* 41:163-70, 1998). COX-1 is a constitutive enzyme thought to be responsible for production of eicosanoids that help maintain normal renal function, gastric mucosal integrity, and homeostasis (Griswold et al., *Med Res Rev* 16:181-206, 1996). COX-2 is expressed during inflammatory reactions such as those induced by cytokines, and is therefore thought to be one of the mediators of inflammation (Arias-Negrete et al., *Biochem Biophys Res Commun* 208:582-9, 1995). This probably can explain why AJA and cannabinoid acids have no ulcerogenicity, while indomethacin, which inhibits both COX-1 and COX-2, has such side effects (Griswold et al., *Med Res Rev* 16: 181-206, 1996).


[0072] The downregulation of COX-2 by AJA may cause a reduction of prostaglandin synthesis, and thus attenuate the cataleptic state induced by PGs (Burststein et al., *Experientia* 43:402-3, 1987). The properties of AJA, including anti-inflammation, anti-tumor effect, and in particular, the adipogenesis demonstrated recently (Recht et al., *Biochem Pharmacol* 62:755-63, 2001), raised the possibility that AJA may mediate its functions in part by activating PPAR $\gamma$ . In line with this hypothesis, some NSAIDs, the traditional

anti-inflammatory agents, have been recently demonstrated to be able to activate PPAR $\gamma$  (Lehmann et al., *J Biol Chem* 272:3406-10, 1997).

[0073] Ajulemic acid (AJA) is a non-psychoactive synthetic cannabinoid based on the template of THC-11-oic acid, which is the natural derivative of THC. THC is biotransformed by oxidation to its principal metabolite, THC-11-oic acid. AJA is synthesized by increasing the side chain of THC-11-oic acid from 5 carbon atoms (pentyl) to 7 atoms (heptyl) and by introducing 2 methyl groups.

### [0074] III. PPAR Activators

[0075] In one aspect, the invention pertains to using THC cannabinoids, e.g., AJA as activators of PPAR, e.g., PPAR $\gamma$ . In one embodiment, the cannabinoid has the following structural formula (Formula I):



[0076] where R<sup>1</sup> is a hydrogen atom, —COCH<sub>3</sub> or —COCH<sub>2</sub>CH<sub>3</sub>, and R<sup>2</sup> is a branched C<sub>5</sub>-C<sub>12</sub> alkyl. R<sup>1</sup> can be hydrogen, and R<sup>2</sup> can be a C<sub>9</sub> alkyl, which can be a branched alkyl such as 1,1-dimethylheptyl. R<sub>3</sub> is —OH, —OCH<sub>3</sub> or —NHCH<sub>2</sub>COOH.

[0077] The cannabinoids defined by Formula I have reduced or no psychoactivity and do not bind to the CB1 receptor. Such cannabinoids are known and can be synthesized (see, e.g., U.S. Pat. No. 5,338,753; Burststein et al., *J Medicinal Chem.* 35:3185-3141, 1992; Burststein, *Pharmacol. Ther.* 82:87-96, 1999).

[0078] Before administration to a subject, the cannabinoids can be tested for biological activity (i.e., ability to decrease cell proliferation) both in vitro or in vivo. In vitro testing can be performed as described in the examples section. In vivo animal models for diabetes are well known in the art, for example, obese-diabetic mice (ob/ob), and obese-diabetic (db/db) mice from the Jackson Laboratories (Bar Harbor, Me)(see, e.g., Collins et al., *J Biol Chem* 271:9437-9440, 1996; Darling, *Curr Opin Genet Dev* 6:289-294, 1996; Andersson, *Ann. Med.* 28:5-7, 1996; Van Heek et al., *J. Clin. Invest* 99:385-390, 1997). These animal models can be used to assess the effect of the cannabinoids on diabetes and obesity.

[0079] The cannabinoid, such as that in Formula I can be administered alone to activate PPAR $\gamma$ , or in addition with existing naturally occurring or synthetic activators to obtain a synergistic effect.

[0080] Examples of naturally occurring activators that modify the activity of PPAR $\gamma$  include, but are not limited to, arachidonic acid derivatives or metabolites such as eicosanoids (e.g., various isomeric forms of 8-hydroxytetraenoic acid) and cyclopentenone prostaglandins (e.g., prostaglandins in the J and A series and their metabolites), and polyunsaturated fatty acids.

[0081] Examples of synthetic activators that modify the activity of PPAR $\gamma$  include, but are not limited to, antidyslipidemic fibrates (e.g., clofibrate, fenofibrate, benzofibrate, ciprofibrate, gemfibrozil), thiazolidine derivatives (e.g., thiazolidinediones), oxazolidine derivatives (e.g., oxazolidinediones), alpha-alkylthio, alpha-alkoxy and carboxylic acid derivatives of thiazolidines and oxazolidines (Hulin et al., *J Med Chem* 39:3897-3907, 1996), N-2-L-tyrosine derivatives (Henke et al., *J Med Chem*; 41:5020-5036, 1998; Collins et al., *J Med Chem* 41:5037-5054, 1998; Cobb et al., *J Med Chem*, 41:5055-5069, 1998), phenyl acetic acid derivatives (Berger et al., *J Biol. Chem.* 274:6718-6725, 1999), and indole-thiazolidinedione derivatives (Lohray et al., *J Med Chem*, 41:1619-1630, 1998).

[0082] IV. Compositions and Formulations

[0083] In one aspect, this invention provides methods and compositions for preventing or inhibiting autoimmune diseases or disorders involving PPAR $\gamma$  by using a PPAR $\gamma$  cannabinoid activator, such as ajulemic acid. In general, the methods involve providing an amount of a PPAR $\gamma$  activator sufficient to modulate the expression of genes encoding proteins involved in autoimmune diseases, such as diabetes mellitus, inflammatory cytokines such as tumor necrosis factor- $\alpha$  (TNF  $\alpha$ ), or inhibition of the production of inflammatory cytokines such as IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-6, IL-8, and TNF- $\alpha$ , or the inhibition of their biological activity. While the PPAR $\gamma$  activator can be utilized alone, the therapy can also be utilized in combination with other therapeutics such as existing naturally occurring or synthetic activators, steroidal and non-steroidal anti-inflammatory agents, existing therapies for diabetes, and agents that modulate apoptosis in pathological cells.

[0084] In one embodiment of the invention, the cells to be treated are those involved in inflammatory disorders. These include inflammatory (immune system) cells (e.g., T lymphocytes and macrophages), PPAR $\gamma$  expressing cells and tissues involved in the pathogenesis of inflammatory diseases, including all forms of uveitis and uveoretinitis, iritis, cyclitis, choroiditis, chorioretinitis, vitritis, keratitis and conjunctivitis, systemic autoimmune disorders (e.g., type 1 diabetes mellitus, sjogren's syndrome and hyperthyroidism), and collagen vascular diseases (e.g., ankylosing spondylitis, rheumatoid arthritis, lupus erythematosus, Reiter syndrome, Bechet disease, ulcerative colitis, Crohn's disease, Wegener's granulomatosis).

[0085] In another embodiment, the cells to be treated are those involved in autoimmune disorders, such as the pancreatic B cells of a subject with diabetes.

[0086] Pharmaceutical compositions containing a compound of interest may be prepared by conventional techniques (e.g., as described in Remington: The Science and Practise of Pharmacy, 19th Ed., 1995). The compositions may appear in conventional forms, for example capsules, tablets, aerosols, solutions, suspensions or topical applications.

[0087] Typical compositions include the cannabinoid, e.g., AJA, or a derivative thereof, associated with a pharmaceutically acceptable excipient which may be a carrier or a diluent or be diluted by a carrier, or enclosed within a carrier which can be in the form of a capsule, sachet, paper or other container. In making the compositions, conventional tech-

niques for the preparation of pharmaceutical compositions may be used. For example, the compound of interest will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of an ampule, capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be solid, semi-solid, or liquid material which acts as a vehicle, excipient, or medium for the active compound. The compound of interest can be adsorbed on a granular solid container for example in a sachet. Some examples of suitable carriers are water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, peanut oil, olive oil, lactose, terra alba, sucrose, cyclodextrin, amylose, magnesium stearate, talc, gelatin, agar, pectin, acacia, stearic acid or lower alkyl ethers of cellulose, silicic acid, fatty acids, fatty acid amines, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, polyoxyethylene, hydroxymethylcellulose, and polyvinylpyrrolidone. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavouring agents. The formulations of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

[0088] The pharmaceutical compositions can be sterilized and mixed, if desired, with auxiliary agents, emulsifiers, salt for influencing osmotic pressure, buffers and/or coloring substances and the like, which do not deleteriously react with the active compounds.

[0089] The route of administration may be any route, which effectively transports the compound of interest to the appropriate or desired site of action, such as oral, nasal, pulmonary, transdermal or parenteral, e.g., rectal, depot, subcutaneous, intravenous, intraurethral, intramuscular, intranasal, ophthalmic solution or an ointment, the oral route being preferred.

[0090] For nasal administration, the preparation may contain the compound of interest dissolved or suspended in a liquid carrier, in particular an aqueous carrier, for aerosol application. The carrier may contain additives such as solubilizing agents, e.g., propylene glycol, surfactants, absorption enhancers such as lecithin (phosphatidylcholine) or cyclodextrin, or preservatives such as parabens.

[0091] To prepare topical formulations, the compound of interest is placed in a dermatological vehicle as is known in the art. The amount of the compound of interest to be administered and the compound's concentration in the topical formulations depend upon the vehicle, delivery system or device selected, the clinical condition of the patient, the side effects and the stability of the compound in the formulation. Thus, the physician employs the appropriate preparation containing the appropriate concentration of the compound of interest and selects the amount of formulation administered, depending upon clinical experience with the patient in question or with similar patients.

[0092] For ophthalmic applications, the compound of interest is formulated into solutions, suspensions, and ointments appropriate for use in the eye. The concentrations are usually as discussed above for local preparations. For oph-



thalmic formulations (see Mitra (ed.), *Ophthalmic Drug Delivery Systems*, Marcel Dekker, Inc., New York, N.Y. (1993) and also Havener, W. H., *Ocular Pharmacology*, C.V. Mosby Co., St. Louis (1983)).

[0093] For oral administration, either solid or fluid unit dosage forms can be prepared. For preparing solid compositions such as tablets, the compound of interest is mixed into formulations with conventional ingredients such as talc, magnesium stearate, dicalcium phosphate, magnesium aluminum silicate, calcium sulfate, starch, lactose, acacia, methylcellulose, and functionally similar materials as pharmaceutical diluents or carriers.

[0094] Capsules are prepared by mixing the compound of interest with an inert pharmaceutical diluent and filling the mixture into a hard gelatin capsule of appropriate size. Soft gelatin capsules are prepared by machine encapsulation of a slurry of the compound of interest with an acceptable vegetable oil, light liquid petrolatum or other inert oil. Fluid unit dosage forms for oral administration such as syrups, elixirs and suspensions can be prepared. The water soluble forms can be dissolved in an aqueous vehicle together with sugar, aromatic flavoring agents and preservatives to form a syrup. An elixir is prepared by using a hydroalcoholic (e.g., ethanol) vehicle with suitable sweeteners such as sugar and saccharin, together with an aromatic flavoring agent. Suspensions can be prepared with an aqueous vehicle with the aid of a suspending agent such as acacia, tragacanth, methylcellulose and the like.

[0095] Appropriate formulations for parenteral use are apparent to the practitioner of ordinary skill, such as the use of suitable injectable solutions or suspensions. Usually, the compound of interest is prepared in an aqueous solution in a concentration of from about 1 to about 100 mg/ml. More typically, the concentration is from about 10 to 60 mg/ml or about 20 mg/ml. Concentrations below 1 mg/ml may be necessary in some cases depending on the solubility and potency of the compound selected for use. The formulation, which is sterile, is suitable for various topical or parenteral routes including intra-dermal, intramuscular, intravascular, and subcutaneous.

[0096] In addition to the compound of interest, the compositions may include, depending on the formulation and mode of delivery desired, pharmaceutically-acceptable, non-toxic carriers or diluents, which include vehicles commonly used to form pharmaceutical compositions for animal or human administration. The diluent is selected so as not to unduly affect the biological activity of the combination. Examples of such diluents which are especially useful for injectable formulations are water, the various saline, organic or inorganic salt solutions, Ringer's solution, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may include additives such as other carriers; adjuvants; or non-toxic, non-therapeutic, non-immunogenic stabilizers and the like.

[0097] Furthermore, excipients can be included in the formulation. Examples include cosolvents, surfactants, oils, humectants, emollients, preservatives, stabilizers and antioxidants. Any pharmacologically acceptable buffer may be used, e.g., tris or phosphate buffers. Effective amounts of diluents, additives, and excipients are those amounts which are effective to obtain a pharmaceutically acceptable formulation in terms of solubility, biological activity, etc.

[0098] The compound of interest maybe incorporated into a microsphere. The compound of interest can be loaded into albumin microspheres, form which it is possible to recover such microspheres in a dry powder for nasal administration. Other materials suitable for the preparation of microspheres include agar, alginate, chitosan, starch, hydroxyethyl starch, ovalbumin, agarose, dextran, hyaluronic acid, gelatin, collagen, and casein. The microspheres can be produced by various processes known to the person skilled in the art such as a spray drying process or an emulsification process.

[0099] For example, albumin microspheres can be prepared by adding rabbit serum albumin in phosphate buffer to olive oil with stirring to produce a water in oil emulsion. Glutaraldehyde solution is then added to the emulsion and the emulsion stirred to cross-link the albumin. The microspheres can then be isolated by centrifugation, the oil removed and the spheres washed, e.g., with petroleum ether followed by ethanol. Finally, the microspheres can be sieved and collected and dried by filtration.

[0100] Starch microspheres can be prepared by adding a warm aqueous starch solution, e.g., of potato starch, to a heated solution of polyethylene glycol in water with stirring to form an emulsion. When the two-phase system has formed (with the starch solution as the inner phase) the mixture is then cooled to room temperature under continued stirring whereupon the inner phase is converted into gel particles. These particles are then filtered off at room temperature and slurried in a solvent such as ethanol, after which the particles are again filtered off and laid to dry in air.

[0101] The microspheres can be hardened by well known cross-linking procedures such as heat treatment or by using chemical cross-linking agents. Suitable agents include dialdehydes, including glyoxal, malondialdehyde, succinaldehyde, adipaldehyde, glutaraldehyde and phthalaldehyde, diketones such as butadione, epichlorohydrin, polyphosphate, and borate. Dialdehydes are used to cross-link proteins such as albumin by interaction with amino groups, and diketones form schiff bases with amino groups. Epichlorohydrin activates compounds with nucleophiles such as amino or hydroxyl to an epoxide derivative.

[0102] The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for subjects, e.g., mammalian subjects, e.g., humans, dogs, cats, and rodents, each unit containing a predetermined quantity of active material calculated to produce the desired pharmaceutical effect in association with the required pharmaceutical diluent, carrier or vehicle. The specifications for the unit dosage forms of this invention are dictated by and dependent on (a) the unique characteristics of the active material and the particular effect to be achieved and (b) the limitations inherent in the art of compounding such an active material for use in humans and animals. Examples of unit dosage forms are tablets, capsules, pills, powder packets, wafers, suppositories, granules, cachets, teaspoonfuls, tablespoonfuls, dropperfuls, ampoules, vials, aerosols with metered discharges, segregated multiples of any of the foregoing, and other forms as herein described. The compositions can be included in kits, which can contain one or more unit dosage forms of the composition and instructions for use to treat one or more of the disorders described herein.

[0103] Slow or extended-release delivery systems, including any of a number of biopolymers (biological-based systems), systems employing liposomes, colloids, resins, and other polymeric delivery systems or compartmentalized reservoirs, can be utilized with the compositions described

herein to provide a continuous or long term source of therapeutic compound. Such slow release systems are applicable to formulations for delivery via topical, intraocular, oral, and parenteral routes.

**[0104]** An effective quantity of the compound of interest is employed in treatment. The dosage of compounds used in accordance with the invention varies depending on the compound and the condition being treated. For example, the age, weight, and clinical condition of the recipient patient. Other factors include: the route of administration, the patient, the patient's medical history, the severity of the disease process, and the potency of the particular compound. The dose should be sufficient to ameliorate symptoms or signs of the disease treated without producing unacceptable toxicity to the patient. In general, an effective amount of the compound is that which provides either subjective relief of symptoms or an objectively identifiable improvement as noted by the clinician or other qualified observer.

**[0105]** Broadly, for a PPAR $\gamma$  activator, the oral dose can be determined from the following formula: oral dose (in mg)= $(k_1)(EC_{50})(k_2)$  (LBW) (MW), wherein  $k_1$  is a constant of 5 to 100,  $EC_{50}$  is the concentration (amount) of compound required to activate or bind to 50% of PPAR $\gamma$  in the sample or patient and is in mole/L units,  $k_2$  is the fractional water content of the lean body weight (LBW) of the patient=0.72 L/kg (see Geigy Scientific Tables Vol. 1, Lentner (ed.), page 217, Giba-Geigy Ltd., Basle, Switzerland (1981)), and MW is the molecular weight of the compound in g/mole.

**[0106]** The compounds in this invention can also be given orally in combination with natural or synthetic compounds that bind to or modify the activity of the vitamin D receptor or in combination with compounds that bind to or modify the activity of the retinoid X receptor to provide for a synergistic effect in the treatment or prevention of the disorders. Examples of such compounds that provide for synergistic effect when given in combination with the drugs encompassed by the current invention include vitamin D analogs, various retinoic acid derivatives, and other activators for retinoid X receptors or retinoic acid receptors including but not limited to compounds such as LG100268, tazarotene, TTNPB, AGN 190121, adapalene, or LGD1069 (Targretin).

**[0107]** Each dosage unit for oral administration contains suitably from 0.1 mg to 500 mg/kg, e.g., from 0.1 mg to 50 mg/kg, e.g., from 0.2 mg to 20 mg/kg, and, e.g., from 0.2 mg to 2 mg/kg, and each dosage unit for parenteral administration can contain from 0.1 mg to 100 mg/kg of a compound of the cannabinoid or derivative thereof calculated as the free base. Each dosage unit for intranasal administration can contain 1-400 mg, e.g., 10 to 200 mg per person.

**[0108]** V. Uses of Cannabinoids as PPAR Activators

**[0109]** In one aspect, the invention pertains to using cannabinoids, e.g., AJA for modifications of disorders associated with PPAR, e.g., PPAR $\gamma$ . The present invention relates to compounds, pharmaceutical compositions containing them, methods for preparing the compounds, and their use as medicaments. More specifically, compounds of the invention can be utilized in the treatment of conditions mediated by nuclear receptors, in particular the PPAR. The present compounds reduce blood glucose and triglyceride levels and are accordingly useful for the treatment of ailments and disorders such as diabetes and obesity.

**[0110]** (i) Use of AJA in Diabetes

**[0111]** In one embodiment, the invention pertains to using AJA as an activator of PPAR $\gamma$  as a method for modifying an autoimmune disorder such as diabetes. Diabetes is a multifactorial disease that occurs through the failure and/or destruction of the pancreatic  $\beta$ -cell. There is a large body of evidence in support of the idea that inflammatory cytokines have cytotoxic effects on islet  $\beta$ -cells (Rabinovitch, *J Clin Endocrinol Metab.* 71:152-6, 1990) and this cytotoxicity plays a part in  $\beta$ -cell destruction in insulin-dependent diabetes mellitus (IDDM).

**[0112]** Obesity-linked non-insulin-dependent diabetes mellitus (NIDDM) is preceded by years of insulin resistance, during which normal blood glucose levels are maintained through effective compensation by pancreatic  $\beta$ -cells. In approximately 20% of obese individuals, the compensation wanes, hyperglycemia appears, and overt NIDDM is diagnosed. The depressed  $\beta$ -cell function is thought to be due to excess free fatty acids released from adipocytes in obesity (Campbell et al., *Am J Physiol.* 266: E600-5., 1994; DeFronzo, *Diabetes Metab Rev.* 4:727-47, 1988) acting to initially stimulate, but ultimately impair, the function of  $\beta$ -cells, and thus limit their compensatory capability. Thus, impaired  $\beta$ -cell function is a characteristic of both IDDM and NIDDM.

**[0113]** The compounds are useful for the modification, treatment, and/or prophylaxis of insulin resistance (Type II diabetes), impaired glucose tolerance, dyslipidemia, disorders related to Syndrome X such as hypertension, obesity, insulin resistance, hyperglycemia, atherosclerosis, hyperlipidemia, coronary artery disease and other cardiovascular disorders associated with diabetes. The disorders also include those associated with diabetes, e.g., the anti-apoptotic effect of PPAR $\gamma$  activators that serves to protect cells from premature death and promote their survival, as in degenerative and dystrophic diseases, e.g., retinal neural and glial cells in diabetic retinopathy and both "wet" (exudative) and "dry" (acreolar) age-related macular degeneration.

**[0114]** (ii) Use of AJA in Lipid Metabolism and Glucose Homeostasis

**[0115]** In another embodiment, the invention pertains to using AJA as an activator of PPAR $\gamma$ , for the treatment or modification of disorders related to lipid metabolism and energy homeostasis. Also, compounds that block PPAR $\gamma$  activity would be useful for interfering with the maturation of preadipocytes into adipocytes and thus would be useful for the treatment of obesity and related disorders associated with undesirable adipocyte maturation.

**[0116]** Adipose tissue plays a central role in lipid homeostasis and the maintenance of energy balance in vertebrates. Adipocytes store energy in the form of triglycerides during periods of nutritional affluence and release it in the form of free fatty acids at times of nutritional deprivation. The development of white adipose tissue is the result of a continuous differentiation process throughout life. Much evidence points to the central role of PPAR $\gamma$  activation in initiating and regulating this cell differentiation. Several highly specialized proteins are induced during adipocyte differentiation, most of them being involved in lipid storage and metabolism. The exact link from activation of PPAR $\gamma$  to changes in glucose metabolism, most notably a decrease in

insulin resistance in muscle, has not yet been clarified. A possible link is via free fatty acids such that activation of PPAR $\gamma$  induces Lipoprotein Lipase (LPL), Fatty Acid Transport Protein (FATP) and Acyl-CoA Synthetase (ACS) in adipose tissue but not in muscle tissue. This, in turn, reduces the concentration of free fatty acids in plasma dramatically, and due to substrate competition at the cellular level, skeletal muscle and other tissues with high metabolic rates eventually switch from fatty acid oxidation to glucose oxidation with decreased insulin resistance as a consequence. PPAR $\gamma$  is involved in stimulating  $\beta$ -oxidation of fatty acids.

[0117] (iii) Use of AJA in Inflammation Involving PPAR $\gamma$

[0118] In another embodiment, the invention pertains to using AJA as an activator of PPAR $\gamma$ , for the treatment or modification of disorders related to those inflammatory responses involving PPAR $\gamma$  activation. The cannabinoid compounds, e.g., AJA, can be used for the treatment of diseases including, but not limited to, immunologically-mediated inflammatory diseases such as rheumatoid arthritis, systemic lupus erythematosus, psoriasis, eczema, multiple sclerosis, diabetes and thyroiditis. In addition, the present compounds can modulate bone formation/resorption and are useful in the treatment of conditions including but not limited to ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis and osteoporosis.

[0119] The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

## EXAMPLES

### Example 1

#### Materials and Methods

[0120] (i) Reagents

[0121] The PPAR $\gamma$ -specific ligand, GW347845, was obtained from GlaxoSmithKline. The 9-cis retinoid acid (9c-RA) was obtained from Activator Pharmaceuticals. Ajulemic acid was obtained from Organix Inc (Woburn, Mass.). All other chemicals were purchased from commercial sources (e.g., Sigma).

[0122] (ii) Plasmids

[0123] The pGEX-DRIP205 (527-970) was obtained from the Sloan-Kettering Cancer Center. The Gal4-hPPAR LBD constructs were obtained from GlaxoSmithKline, Research Triangle Park, NC. The IL8-Luciferase reporter was obtained from Celgene, San Diego, Calif. The plasmids mPPAR $\alpha$ , mPPAR $\gamma$ 1, mPPAR $\delta$ , and hRXR $\alpha$  were in the pCMX vector (Umesono et al., *Cell* 65:1255-1266, 1991).

[0124] The pCMX-Gal4-mPPAR $\gamma$ 1 was constructed by fusing the mPPAR $\gamma$ 1 coding sequence to the yeast GAL4 DNA binding domain (amino acid 1-147) in the pCMX vector. The AF-2 helix-truncated mutant of mPPAR $\gamma$ 1 (PPAR $\gamma$  $\Delta$ AF2) was generated by PCR amplification-to introduce a stop codon and a NheI restriction site after amino acid 489, and then-subcloned into the pCMX vector. The PPRE-TK-LUC, MH100-TK-LUC-and DR1-TK-LUC reporters have been described previously (Forman et al., *Cell* 83:803-812, 1995; Umesono et al., *Cell* 65:1255-1266, 1991).

[0125] (iii) Cell Culture and Transient Transfection

[0126] For PPAR $\gamma$  activation assay, HEK293 cells were plated in 12-well cell culture plates and maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) at 37° C. under 5% CO<sub>2</sub>. Cells were changed to phenol red-free DMEM supplemented with 10% charcoal-stripped fetal bovine serum 3 hours before transfection by a standard calcium-phosphate precipitation method. Twelve hours after transfection, the cells were washed with phosphate-buffered saline (PBS) and fed again with fresh medium containing the indicated concentrations of compounds. After 36 hours, cells were harvested for  $\beta$ -galactosidase and luciferase activities as described previously (Li et al., *Proc Natl Acad Sci USA* 94:8479-84, 1997). The average normalized luciferase activity was determined via triplicate experiments. For IL-8 promoter assay, HeLa cells were maintained and transfected in the same way as described above. After transfection, cells were recovered for 4 hours before treatment with AJA, GW347845, or solvent. After 3-4 hours, cells were treated with or without 25 nM of phorbol myristate acetate (PMA) for 24 hours.

[0127] (iv) Partial Protease Digestion Assay

[0128] Partial protease digestion was carried out as described by Leng et. al. (*J Steroid Biochem Mol Biol* 46:643-61, 1993). PPAR proteins were made by in vitro transcription/translation reactions in reticulocyte lysate according to the manufacture's instructions (Promega). AJA, GW347845, or vehicle alone was incubated with the 35S-labeled PPARs at room temperature for one hour before trypsin digestion. Reactions were stopped by boiling in SDS-containing sample buffer, and lysates were subjected to SDS-PAGE and autoradiography.

[0129] (v) Expression and Purification of GST Fusion Protein

[0130] The glutathione S-transferase (GST)-DRIP205 (amino acid 527-970) and GST-RAC3 RID (amino acid 613-752) fusion proteins were expressed in BL21 cells by induction with 0.1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) at room temperature for 4 hours. The bacterial pellet was resuspended in STE buffer (150 mM NaCl, 10 mM Tris pH 8.0, 1 mM EDTA), containing 5 mM dithiothreitol (DTT), 1 mM phenylmethylsulfonyl fluoride (PMSF) and 1.5% Sarkosyl. The cell suspension was sonicated and centrifuged at 7400 rpm at 4° C. for 30 minutes. The supernatant was isolated and 0.02% Triton-X 100 was added, and the mixture was then incubated with 1 ml 50% slurry of glutathione-Sepharose beads on a rotator at 4° C. for 30 minutes. The beads were spun down at 3000 rpm for 10 minutes, the supernatant was removed, and the beads were suspended in 1 ml cold PBS.

[0131] (vi) GST Pull-Down Assay

[0132] Approximately 5  $\mu$ g of GST-DRIP205 or GST-RAC3 bound on glutathione-Sepharose beads and 4  $\mu$ l <sup>35</sup>S-methionine-labeled mPPAR $\gamma$ 1 were incubated with the indicated concentrations of AJA, GW347845, or vehicle alone in H buffer (20 mM Hepes [pH 7.7], 75 mM KCl, 0.1 mM EDTA, 2.5 mM MgCl<sub>2</sub>, 0.05% NP40, 0.1 mM methionine, 1 mM DTT) containing 1 mg/ml bovine serum albumin (BSA) on a rotator at 4° C. overnight. After three washes with cold PBS, the bound protein was eluted in SDS sample

buffer and boiled for 10 min before SDS-PAGE and autoradiography. To ensure that equal amounts of GST fusion proteins were recovered in the pull-down assay, the gel was stained with Coomassie blue before autoradiography.

**[0133]** (vii) Adipocyte Differentiation Assay and RT-PCR

**[0134]** The adipocyte differentiation assay was performed as described by Mukherjee et. al. (*Mol Endocrinol* 14: 1425-33, 2000). The 3T3 L1 cells (American Type Culture Collection) were cultured in DMEM media supplemented with 10% calf serum. Two days after reaching confluence, cells were treated with AJA, GW347845, or vehicle (0.1% DMSO) in the presence of 10  $\mu\text{g}/\text{ml}$  insulin every other day. After 10 days of treatment with AJA or 7 days with GW347845 at confluence, cells were fixed and stained with Oil Red O (Sigma). For RT-PCR, total cellular RNAs were isolated by TRIzol (Gibco). Reverse transcriptase polymerase chain reaction (RT-PCR) was performed using the superscript first-strand synthesis kit (Invitrogen). After first-strand cDNA was synthesized by use of oligo (dT), cDNA was amplified by PCR. The forward primers and reverse primers used in the amplifications were 5'-GCT GTT ATG GGT GAA ACT CTG GGA G-3' (SEQ ID NO: 1), and 5'-CTT CAT GAG GCC TGT TGT AGA GC-3' (SEQ ID NO:2), for PPAR $\gamma$ 2, 5'-GAG CAA ATG GAG TTC CCA GAT G-3' (SEQ ID NO: 3), and 5'-GCA AAC AAT GGG AAT AGT TCA CAG TAG-3' (SEQ ID NO: 4), for aP2, and 5'-GAC CAC AGT CCA TGC CAT CAC-3' (SEQ ID NO: 5), and 5'-CAT ACC AGG AAA TGA GCT GAC-3' (SEQ ID NO: 6), for GAPDH. PCR amplifications were performed in 50  $\mu\text{L}$  volume with Taq DNA polymerase for 30 cycles.

#### Example 2

##### Binding of Ajulemic Acid to PPAR $\gamma$

**[0135]** To determine if AJA can bind to PPAR $\gamma$ , in vitro translated  $^{35}\text{S}$ -labeled mPPAR $\gamma$ 1 was incubated with AJA or a carrier vehicle before trypsin digestion. To test if AJA binds to PPARs, partial proteinase digestion assays were conducted as described previously (Leng et al., *J Steroid Biochem Mol Biol* 46:643-61, 1993). Ligand binding is known to cause conformational changes of nuclear receptors, which is an essential process for transcriptional activation by the receptors (Allen et al., *J Biol Chem* 267:19513-20, 1992). Such conformational change can be detected by limited proteinase digestion based on alteration of the accessibility of proteolytic sites on the surface of the receptors.

**[0136]** The PPAR proteins were synthesized and labeled with  $^{35}\text{S}$ -methionine in reticulocyte lysate and incubated with AJA, GW347845, or vehicle alone at room temperature for one hour before trypsin digestion. The reactions were terminated by boiling in SDS-containing protein sample buffer. The proteolytic patterns of PPAR proteins were then analyzed by SDS-PAGE and autoradiography, and compared with vehicle-treated control samples.

**[0137]** In the experiment illustrated in FIG. 2A, the final concentrations of ligands were 20  $\mu\text{M}$  AJA and 1  $\mu\text{M}$  GW347845 in 0.2% DMSO (vehicle). Partial proteinase digestions were conducted with 30  $\mu\text{g}/\text{ml}$  of trypsin for 30 or 60 minutes as indicated. FIG. 2A shows a clear difference in the proteolytic profiles between AJA and vehicle-treated PPAR $\gamma$  proteins was observed. Two prominent trypsin-resistant fragments of 30 and 24 kDa, respectively, were

detected after incubation with 20  $\mu\text{M}$  AJA when compared with the control sample, in which the 30-kDa band disappeared completely after 60 min of digestion and two smaller fragments emerged after 30 min of digestion. As a positive control, the potent PPAR $\gamma$  agonist GW347845 (Cobb et al., *J Med Chem*; 41:5055-5069, 1998; Suh et al., *Cancer Res* 59:5671-3, 1999), also produced a proteinase digestion pattern similar to the one seen in AJA-treated sample. These findings show that AJA treatment can cause a conformational change in PPAR $\gamma$ , demonstrating that AJA may bind directly to PPAR $\gamma$ .

**[0138]** The AJA concentrations required to protect PPAR $\gamma$  from trypsin digestion were further measured. The two proteinase-resistant PPAR $\gamma$  fragments were observed at 2, 20 and 100  $\mu\text{M}$  AJA concentrations, with a slight dose-dependent increase between 2 and 20  $\mu\text{M}$  (FIG. 2B). In contrast, AJA at all of these concentrations did not affect the proteinase sensitivity of PPAR $\alpha$  (FIG. 2C) or PPAR $\delta$  (FIG. 2D). These data indicate that AJA binds specifically to PPAR $\gamma$ , but it does not interact with PPAR $\alpha$  or PPAR $\delta$ , suggesting that AJA is a selective ligand for PPAR $\gamma$ .

#### Example 3

##### Ajulemic Acid Activates PPAR $\gamma$

**[0139]** The observation that AJA induces conformational change of PPAR $\gamma$  prompted the investigation of whether AJA was a PPAR $\gamma$  agonist, and could activate PPAR $\gamma$  transcriptional activity. Therefore, transient transfections were performed to investigate this possibility. The PPRE-TK-LUC reporter (Kliewer et al., *Proc Natl Acad Sci USA* 91:7355-9, 1994) was cotransfected with the PPAR $\gamma$  expression vector into the human kidney HEK293T cells followed by AJA treatment of the cells. The average luciferase activities were normalized with the cotransfected  $\beta$ -galactosidase in triplicate experiments. After transfections, cells were treated with indicated concentrations of ligands or vehicle alone for 36 hours before harvesting for  $\beta$ -galactosidase and luciferase assays. In this functional assay, 1  $\mu\text{M}$  GW347845 stimulated the transcriptional activity of PPAR $\gamma$  4 fold. AJA also stimulated PPAR $\gamma$  transcriptional activity 2 to 3 fold at 1 to 10  $\mu\text{M}$  concentrations in a dose-dependent manner (FIG. 3A). These results suggest that AJA indeed can activate the transcriptional activity of PPAR $\gamma$ , consistent with its ability to bind PPAR $\gamma$  and induce an active conformation.

**[0140]** In an effort to confirm that AJA activates PPAR $\gamma$  selectively, the ability of AJA to activate PPAR $\alpha$  and PPAR $\delta$  was analyzed in a similar transient transfection assay. PPAR $\alpha$  had a 3-fold higher basal transcriptional activity than PPAR $\gamma$  in the absence of ligand; however, no further activation was observed after treatment with 1 to 20  $\mu\text{M}$  AJA concentrations (FIG. 3B and data not shown). Similarly, neither did AJA activate PPAR $\delta$  in this assay (FIG. 3C). These data indicate that AJA activates PPAR $\gamma$  selectively, consistent with its ligand-binding specificity.

**[0141]** PPAR $\gamma$  forms a permissive heterodimeric complex with the retinoid X receptor (RXR) and ligands for either PPAR $\gamma$  or RXR can both activate the receptor heterodimer (for a review, see reference Leblanc et al., *Genes Dev* 9:1811-6, 1995). In order to rule out the possibility that AJA might act through RXR, the ability of AJA to activate RXR $\alpha$

on a DR1-LUC reporter was analyzed, where RXR is known to form a homodimer that can be activated only by RXR-specific ligands. The hRXR $\alpha$  and a DR1-TK-LUC reporter were cotransfected into HEK293 cells, followed by 9-cis RA or AJA treatments. In this experiment, it was found that the RXR-specific ligand 9-cis RA activated the reporter gene expression 3 fold, but AJA failed to activate the reporter at saturating concentrations (**FIG. 3D**), suggesting that AJA does not bind or activate RXR $\alpha$ . These data indicate that the transcriptional activation of PPAR $\gamma$  by AJA in vivo is not mediated through its heterodimeric partner RXR $\alpha$ .

[0142] PPAR $\gamma$  contains two transcriptional activation domains: a constitutive N-terminal AF-1 domain and a ligand-dependent C-terminal AF-2 domain. The AF-2 function depends on the presence of an AF-2 helix (helix 12) located at the extreme C-terminal end of LBD. To examine whether activation of PPAR $\gamma$  by AJA is mediated through the ligand-dependent AF-2 function, the AF-2 helix of PPAR $\gamma$  was deleted to create a PPAR $\gamma$  $\Delta$ AF2 mutant, and then tested as to whether AJA could still activate the PPRE-Luc reporter through the  $\Delta$ AF2 mutant. AJA and GW347845 both failed to activate expression of the reporter gene (**FIG. 3E**), suggesting that activation of PPAR $\gamma$  by AJA is mediated specifically through the AF-2 function. This is also consistent with the idea that AJA is an activating ligand for PPAR $\gamma$ .

[0143] To further confirm activation of PPAR $\gamma$  by AJA, and also to determine PPAR $\gamma$  species specificity, the ability of AJA to activate the chimeric Gal4-DBD/PPAR fusion proteins on a Gal4-dependent MH100-Luc reporter was assessed (**FIG. 4A**). The PPARs were expressed as Gal4 DBD fusion proteins, which bind to the UAS promoter containing 4 copies of a synthetic Gal4 binding site upstream of the minimal thymidine kinase (TK) promoter. Transient transfection was conducted in HEK293 cells with the mPPAR $\gamma$  expression vector and the MH100-tk-Luc reporter. The Gal4 DBD/PPAR fusion protein will activate reporter gene expression in response to agonist binding to PPAR. In this system, activation of the reporter will be mediated exclusively through the chimeric Gal4-DBD fusion protein, thus eliminating potential interference from endogenous receptors. In this assay, both AJA and GW347845 activated the reporter gene expression significantly (**FIG. 4B**). Transfected cells were treated with increasing concentrations of AJA from 10-60  $\mu$ M and the luciferase activities were determined in triplicate experiments. The estimated EC-50 for activation of PPAR $\gamma$  by AJA is 13  $\mu$ M in this assay. The activation of Gal4-DBD/mPPAR $\gamma$  by AJA was concentration-dependent, with an estimated EC-50 of approximately 13  $\mu$ M (**FIG. 4C**). The efficacy of AJA to activate human PPAR $\gamma$  in this assay was also tested, and confirmed that AJA also activates human PPAR $\gamma$  as efficiently as mouse PPAR $\gamma$  (**FIG. 4D**). Consistently, AJA did not activate human PPAR $\alpha$  or PPAR $\delta$ . Taken together, these data strongly suggest that AJA activates both mouse and human PPAR $\gamma$  specifically at pharmacologically relevant concentrations.

#### Example 4

##### AJA Stimulates Coactivator Binding to PPAR $\gamma$

[0144] This example demonstrates that AJA-bound PPAR $\gamma$  can recruit coactivators. Nuclear receptor coactivators are known to interact with ligand-activated receptors to

enhance transcriptional activation by recruiting chromatin modifying enzymes and RNA polymerase. The receptor-associated coactivator-3 (RAC3) of the p160/SRC family, and the DRIP205 subunit of the DRIP coactivator complex are known PPAR $\gamma$  coactivators (Yang et al., *Mol Cell Biol* 20:8008-17, 2000).

[0145] To evaluate further the mechanism whereby AJA influences PPAR $\gamma$  transcriptional activity, the ability of PPAR $\gamma$  to interact with DRIP205 and/or RAC3 in response to AJA treatment by GST-pull down assay was examined. The GST-DRIP205 (amino acids 527-970) and the GST-RAC3 (amino acids 613-752) fusion proteins were purified and used in binding reactions containing in vitro translated <sup>35</sup>S-labeled PPAR $\gamma$  in the presence of vehicle (0.1% DMSO), GW347845 (GW34, 1  $\mu$ M), or AJA (20  $\mu$ M). GST alone was used as a negative control in the pull down reaction. In this experiment, the <sup>35</sup>S-methionine-labeled PPAR $\gamma$  showed negligible binding to GST alone, GST-DRIP205 or GST-RAC3 in the absence of ligand (**FIGS. 5A and 5B**). In contrast, AJA and GW347845 pretreatment of PPAR $\gamma$  caused increased binding of PPAR $\gamma$  to the GST-DRIP205 and GST-RAC3. The interaction between PPAR $\gamma$  and DRIP205 appeared to be stronger than the interaction between PPAR $\gamma$  and RAC3, consistent with the fact that DRIP205 is a more potent coactivator than RAC3 for PPAR $\gamma$  (Yang et al., *Mol Cell Biol* 20:8008-17, 2000; Zhu et al., *J Biol Chem* 272:25500-6, 1997). These data indicate that AJA treatment can promote the interaction of PPAR $\gamma$  with nuclear receptor coactivators, corroborating further the hypothesis that AJA binds directly to PPAR $\gamma$  and induces a transcriptionally active conformation of the receptor.

#### Example 5

##### Reduction of IL-8 Promoter Activity by Ajulemic Acid Through PPAR $\gamma$

[0146] This example demonstrates the link between activation of PPAR $\gamma$  by AJA to the antiinflammatory activity of AJA. The IL-8 luciferase reporter controlled by the human IL-8 promoter (-97/-69) was cotransfected with mPPAR $\gamma$  or mPPAR $\gamma$  $\Delta$ AF2 mutant into HeLa cells. After transfection, cells were treated with AJA, GW347845 or solvent after PMA treatment for 24 hours. This was performed by determining the effect of AJA on IL-8 promoter activity and the role of PPAR $\gamma$  in this process. IL-8 is a biomarker for inflammation, and reduction of IL-8 levels correlates with a decrease in inflammation. The involvement of PPAR $\gamma$  in regulating IL-8 promoter activity was determined by comparing the IL-8 promoter activity in the presence of wild type PPAR $\gamma$  or its  $\Delta$ AF2 mutant. In the experiment illustrated in **FIG. 6A**, PMA stimulated IL-8 promoter activity 3 fold, which was set as 100% promoter activity. AJA treatment reduced the IL-8 promoter activity by about 40% at 20  $\mu$ M concentration in cells transfected with the wild type PPAR $\gamma$ . This effect appears concentration-dependent, and the reduction of IL-8 promoter activity is statistically significant at 10 and 20  $\mu$ M AJA concentrations. AJA had no effect on IL-8 promoter activity in cells transfected with the PPAR $\gamma$  $\Delta$ AF2 mutant (**FIG. 6B**), suggesting that the reduction of IL-8 promoter activity is mediated through activation of PPAR $\gamma$  by AJA. As a control, GW347845 dramatically reduced IL-8 promoter activity in cells cotransfected with the wild type PPAR $\gamma$ , but not with the  $\Delta$ AF2 mutant (**FIGS.**

6C and D). These data indicate that activation of PPAR $\gamma$  by AJA is involved in the inhibition of IL-8 promoter activity, suggesting a putative mechanism for the antiinflammatory action of AJA.

#### Example 6

##### Induction of Adipocyte Differentiation by Ajulemic Acid

[0147] To provide biological evidence that AJA is a bona fide activator for PPAR $\gamma$ , the ability of AJA to induce differentiation of 3T3 L1 fibroblasts into adipocytes was assessed. Activation of PPAR $\gamma$  by its ligands is an essential process for the onset of adipocyte differentiation, which is characterized by morphological changes, droplet formation, and induction of adipocyte-specific genes such as PPAR $\gamma$ 2 and aP2 (Barak et al., *Mol Cell* 4:585-95, 1999; Rosen et al., *Genes Dev* 16:22-6, 2002; Tontonoz et al., *Cell* 79:1147-56, 1994). PPAR $\gamma$  ligands can substitute for the adipogenic hormones during differentiation of preadipocytes into adipocytes (Chawla et al., *Proc Natl Acad Sci USA* 91:1786-90, 1994), and ectopically expressed PPAR $\gamma$  is able to transdifferentiate myoblasts into adipocytes (Hu et al., *Proc Natl Acad Sci USA* 92:9856-60, 1995).

[0148] 3T3 L1 fibroblasts cells were cultured in DMEM supplemented with 10% calf serum. Two days after confluence, cells were treated with 0.1% DMSO, 20  $\mu$ M AJA, or 1  $\mu$ M GW347845 in the presence of 10  $\mu$ g/ml of insulin. The cells were treated for 10 days with AJA, or 7 days with GW347845, and lipid accumulation in cells was assessed by Oil Red O staining. A dramatic increase in lipid droplet staining in the cytoplasm was observed after treatment with AJA or GW347845 (FIG. 7A), suggesting that both AJA and GW347845 can induce differentiation of 3T3 L1 fibroblasts into adipocytes.

[0149] To confirm that AJA and GW347845 induce 3T3 L1 cell differentiation into adipocytes, the expression levels of the adipocyte-specific genes PPAR $\gamma$ 2 and aP2 were measured by RT-PCR. Total RNA was isolated after AJA or GW347845 treatment, and compared with vehicle control. Reverse transcription was conducted and the relative amounts of PPAR $\gamma$ 2 and aP2 transcripts were measured by PCR reactions using primer sets specific to each gene. PCR products were analyzed on 1% agarose gel and stained with Ethidium Bromide. Both AJA and GW347845 enhanced PPAR $\gamma$ 2 and aP2 gene expression significantly in comparison with vehicle control treatment (FIG. 7B). As a control for the induction specificity and the PCR amplification reaction, the expression levels of GADPH were measured in all samples, and found not affected by any treatment. These data indicate that AJA can induce differentiation of 3T3 L1 fibroblasts into adipocytes, further suggesting that AJA is an active ligand for PPAR $\gamma$ .

[0150] The molecular basis for the therapeutic action of ajulemic acid (AJA), were investigated. The data upon which the invention is based, in part, demonstrate that AJA can bind selectively to PPAR $\gamma$  in vitro and that AJA activates the transcriptional activity of PPAR $\gamma$  in vivo. Activation of PPAR $\gamma$  by AJA depends on the presence of the AF-2 helix in the receptor. AJA binding enables PPAR $\gamma$  to recruit nuclear receptor coactivators. In addition, AJA inhibits IL-8 promoter activity in a PPAR $\gamma$ -dependent manner, and induces

differentiation of 3T3 L1 fibroblasts into adipocytes. The data suggest that AJA exerts its therapeutic actions, at least in part, through activation of PPAR $\gamma$ .

[0151] Based on the structural and functional similarity of AJA and several known PPAR $\gamma$  ligands, it was demonstrated that AJA binds to PPAR $\gamma$ . In a partial proteinase digestion assay, AJA effectively protects PPAR $\gamma$  from proteinase digestion (FIGS. 2A-D), reflecting direct binding of AJA to PPAR $\gamma$ . The two trypsin resistant fragments likely contain the ligand-binding domain (LBD) of PPAR $\gamma$ , because ligand binding induces a compact conformation in the LBD (Xu et al., *Proc Natl Acad Sci USA* 98:13919-24, 2001), which is expected to be more resistant to proteinase digestion when compared with the unliganded receptor. Because AJA causes proteinase resistance only to PPAR $\gamma$ , but not PPAR $\alpha$  or PPAR $\delta$ , it is clear that AJA binds selectively to PPAR $\gamma$ . Indeed, the three PPARs have distinct ligand binding specificities (Xu et al., *Proc Natl Acad Sci USA* 98:13919-24, 2001), and physiological functions (Berger et al., *Annu Rev Med* 53:409-35, 2002). The selective binding of AJA to PPAR $\gamma$  suggests that PPAR $\gamma$  may mediate the therapeutic activity of AJA.

[0152] Consistent with the binding of AJA to PPAR $\gamma$ , the transient reporter gene assay further demonstrates that AJA activates the target promoter through PPAR $\gamma$ , but not PPAR $\alpha$  or PPAR $\delta$  (FIGS. 3A-E). These data indicate that AJA is a PPAR $\gamma$ -specific agonist, a conclusion that is further supported by the inability of AJA to activate the PPAR $\gamma$  $\Delta$ AF2 mutant, which is defective in ligand-dependent transcriptional activation (FIG. 3E). The observation that PPAR $\gamma$  activation by AJA requires the AF-2 function is consistent with the hypothesis that AJA binds to PPAR $\gamma$  LBD and activates its ligand-dependent transcriptional function. By using the Gal4-DBD fusion protein system, the EC-50 of AJA for PPAR $\gamma$  activation was measured to be 13  $\mu$ M (FIGS. 4A-D), a concentration that is within the pharmacologically effective doses of AJA (Burststein, *Curr Pharm Des* 6:1339-45, 2000). Furthermore, AJA also activates human PPAR $\gamma$  equally well compared with mouse PPAR $\gamma$ , implying that the PPAR $\gamma$ -dependent therapeutic activity of AJA observed in mice might be applicable to humans.

[0153] As an agonist for PPAR $\gamma$ , AJA's binding enables PPAR $\gamma$  to recruit nuclear receptor coactivators (FIGS. 5A and 5B). Nuclear receptor coactivators are known to interact with liganded receptors to facilitate transcriptional activation of a target promoter by recruiting histone acetyltransferase activity to the receptor (Leo and Chen 2000). The interaction of PPAR $\gamma$  with DRIP205 in response to AJA treatment appears to be more prominent than the interaction with RAC3. This is consistent with a report showing that DRIP205 is a potent coactivator for PPAR $\gamma$  (Yang et al., *Mol Cell Biol* 20:8008-17, 2000). AJA may induce formation of a coactivator-binding surface to allow docking of the coactivator LXXLL motif. Furthermore, addition of AJA causes differentiation of 3T3 L1 fibroblasts into adipocytes (FIGS. 7A-D), a process that is known to be mediated by PPAR $\gamma$ . Together, these data provide strong evidence that AJA is an agonist of PPAR $\gamma$ . These data also demonstrate the involvement of PPAR $\gamma$  in the signaling of this cannabinoid class of analgesic and antiinflammatory drugs.

[0154] Activation of PPAR $\gamma$  is, at least partly, responsible for the antiinflammatory action of AJA, and other cannabinoids as well. In fact, several natural cannabinoids, such as THC and cannabidiol (CBD), also activate PPAR $\gamma$ . The involvement of PPAR $\gamma$  in AJA-mediated antiinflammatory

activity is further supported by the observation that AJA inhibits IL-8 promoter activity in a PPAR $\gamma$ -dependent manner (FIGS. 6A-D). This inhibition occurs only in the presence of wild type PPAR $\gamma$ , but not the  $\Delta$ AF-2 mutant, suggesting that transcriptional activation by PPAR $\gamma$  is required for the repression of IL-8 promoter activity by AJA. It is not clear whether inhibition of cytokine promoter activity by PPAR $\gamma$  is due to direct binding of PPAR $\gamma$  to the promoter. Since there is no evidence for direct binding of PPAR $\gamma$  to the cytokine promoter, it is possible that this inhibition is indirect.

[0155] Treatment of peritoneal macrophages with several PPAR $\gamma$  ligands such as 15d-PGJ<sub>2</sub> also suppresses expression of the inducible nitric oxide synthase (iNOS), gelatinase B, and scavenger receptor A in response to phorbol ester stimulation. The promoters of these genes were found to possess binding sites for activator protein-1 (AP-1), nuclear factor  $\kappa$ B (NF- $\kappa$ B), and the signal transducer and activator of transcription (STAT). Furthermore, the inhibition of inflammatory response by PPAR $\gamma$  ligands in macrophages was produced in part by antagonizing the activities of these transcription factors (Ricote et al., *Nature* 391:79-82, 1998). Although 15d-PGJ<sub>2</sub> was shown in some studies to be able to mediate an anti-inflammatory action in a PPAR $\gamma$ -independent manner (Straus et al., *Proc Natl Acad Sci USA* 97:4844-9, 2000; Tsubouchi et al., *Biochem Biophys Res Commun* 283:750-5, 2001), the inhibition of inflammatory responses by activation of PPAR $\gamma$  was confirmed by several recent studies both in vitro (Ji et al., *J Autoimmun* 17:215-21, 2001) and in vivo (Dubuquoy et al., *Gastroenterol Clin Biol* 24:719-24, 2000; Kawahito et al., *J Clin Invest* 106:189-97, 2000; Naito et al., *Aliment Pharmacol Ther* 15:865-73, 2001).

[0156] It is interesting to note that, like the activation of PPAR $\gamma$ , the immunosuppressive functions of some cannabinoids were also reported to be exerted by the inhibition of NF- $\kappa$ B, STAT, and AP-1 (Faubert et al., *J Leukoc Biol* 67:259-66, 2000; Jeon et al., *Mol Pharmacol* 50:334-41, 1996; Zheng et al., *Biochem Pharmacol* 51:967-73, 1996). The complete mechanism of the antiinflammatory action of AJA is likely to involve a complex signaling network. PPAR $\gamma$  is highly expressed in several immune cells, such as macrophages and monocytes. Therefore, it is possible that AJA may activate PPAR $\gamma$  in these cells to modulate immune inflammatory responses. Because PPAR $\gamma$  is involved in several physiological processes including lipid metabolism, glucose homeostasis, and adipocyte differentiation, the discovery that AJA is an active ligand for PPAR $\gamma$  led to the discovery that AJA may have a broader range of pharmacological activities than previously expected. For example, AJA is a useful agent for treatment of diabetes mellitus.

#### Example 7

##### Treatment of a Patient with Diabetes Mellitus

[0157] A 53 year old male patient who weighs 105 kg is diagnosed by his physician as having diabetes mellitus. The patient is mildly obese, has high blood sugar, and exhibits early signs of atherosclerosis. The physician prescribes treatment with ajulemic acid.

[0158] Accordingly, under physician supervision the patient takes an oral dose of 10 mg of ajulemic acid (in a pharmaceutically and palatably acceptable carrier) once every two weeks for three months. The physician expects this course of treatment to cause the patient to lose weight

at a healthy rate, lower the patient's blood sugar back into the normal healthy range, and slow progression of the patient's atherosclerosis.

#### Example 8

##### Treatment of a Patient with Rheumatoid Arthritis

[0159] A 62 year old female patient who weighs 70 kg is diagnosed by her rheumatologist as having severe rheumatoid arthritis. The patient feels stiffness and prolonged soreness and aching in her joints, particularly in her elbow, knee, and shoulder joints, which has limited her movement and mobility. She feels general fatigue. Her rheumatoid factor is also consistently measured to be outside the normal range. The rheumatologist prescribes treatment with ajulemic acid.


[0160] Once per month the patient visits her rheumatologist for an injection into her arm of 15 mg of ajulemic acid (delivered in a pharmaceutically acceptable carrier). This course of treatment continues indefinitely. The rheumatologist expects treatment with ajulemic acid to reduce inflammation in the patient's joints, thus alleviating the stiffness, soreness, and aching, and allowing enhanced mobility.

#### Other Embodiments

[0161] It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

What is claimed is:

1. A method of treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, the method comprising administering to the subject a pharmaceutically effective amount of a composition comprising a cannabinoid compound.
2. The method of claim 1, wherein the cannabinoid has the general formula:



wherein R<sup>1</sup> is a hydrogen atom, —COCH<sub>3</sub> or —COCH<sub>2</sub>CH<sub>3</sub>, R<sup>2</sup> is a branched C<sub>5</sub>-C<sub>12</sub> alkyl, and R<sup>3</sup> is OH, OCH<sub>3</sub>, or NHCH<sub>2</sub>COOH.

3. The method of claim 2, wherein R<sup>2</sup> is a C<sub>9</sub> alkyl.
4. The method of claim 2, wherein R<sup>2</sup> is a branched alkyl.
5. The method of claim 2, wherein R<sup>2</sup> is 1,1-dimethylheptyl.
6. The method of claim 1, wherein the cannabinoid is ajulemic acid.
7. The method of claim 1, wherein the subject is sensitive to the cannabinoid.

8. The method of claim 1, wherein the disorder is an autoimmune disorder.

9. The method of claim 8, wherein the autoimmune disorder is diabetes mellitus, impaired glucose intolerance, diabetic retinopathy, obesity, systemic lupus erythematosus, rheumatoid arthritis, spondylo arthritis, asthma, inflammatory bowel disease, vasculitis, dermatomyositis, polymyositis, sjogren's syndrome, or hyperthyroidism.

10. The method of claim 1, wherein the disorder is ankylosing spondylitis, gout, arthritis associated with gout, osteoarthritis, osteoporosis, atherosclerosis, hypertension, hyperglycemia, coronary artery disease, or dyslipidemia.

11. The method of claim 1, wherein the disorder is exudative age-related macular degeneration or aereolar age-related macular degeneration.

12. The method of claim 1, wherein the disorder is an inflammatory disorder.

13. The method of claim 12, wherein the inflammatory disorder is rheumatoid arthritis, multiple sclerosis, myasthenia gravis, uveoretinitis, uveitis, iritis, cyclitis, choroiditis, chorioretinitis, vitritis, keratitis, conjunctivitis, psoriasis, eczema, thyroiditis, or a collagen vascular disorder.

14. The method of claim 13, wherein the collagen vascular disease is ankylosing spondylitis, lupus erythematosus, Reiter syndrome, Bechet disease, ulcerative colitis, Crohn's disease, or Wegener's granulomatosis.

15. The method of claim 1, wherein the mode of administration of the composition is oral, nasal, pulmonary, transdermal, or parenteral.


16. The method of claim 1, wherein the subject is a mammal.

17. The method of claim 16, wherein the mammal is a human, non-human primate, dog, cat, rodent, horse, cow, sheep, or goat.

18. A method of treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, the method comprising administering to the subject a pharmaceutically effective amount of a PPAR $\gamma$  activator.

19. The method of claim 18, wherein the PPAR $\gamma$  activator is a composition comprising a cannabinoid compound.

20. The method of claim 19, wherein the cannabinoid has the general formula:




wherein R<sup>1</sup> is a hydrogen atom, —COCH<sub>3</sub> or —COCH<sub>2</sub>CH<sub>3</sub>, R<sup>2</sup> is a branched C<sub>5</sub>-C<sub>12</sub> alkyl, and R<sup>3</sup> is OH, OCH<sub>3</sub>, or NHCH<sub>2</sub>COOH.

21. The method of claim 19, wherein the cannabinoid is ajulemic acid.

22. A kit comprising a pharmaceutically effective amount of a composition comprising a cannabinoid compound and comprising instructions for use of the composition in treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function.

23. The kit of claim 22, wherein the cannabinoid has the general formula:



wherein R<sup>1</sup> is a hydrogen atom, —COCH<sub>3</sub> or —COCH<sub>2</sub>CH<sub>3</sub>, R<sup>2</sup> is a branched C<sub>5</sub>-C<sub>12</sub> alkyl, and R<sup>3</sup> is OH, OCH<sub>3</sub>, or NHCH<sub>2</sub>COOH.

24. The kit of claim 23, wherein R<sup>2</sup> is a C<sub>9</sub> alkyl.

25. The kit of claim 23, wherein R<sup>2</sup> is a branched alkyl.

26. The kit of claim 23, wherein R<sup>2</sup> is 1,1-dimethylheptyl.

27. The kit of claim 22 wherein the cannabinoid is ajulemic acid.

28. A composition for use as a medicament in treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, wherein the composition comprises a pharmaceutically effective amount of a cannabinoid compound.

29. Use of a composition for the manufacture of a medicament for use in treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, wherein the composition comprises a pharmaceutically effective amount of a cannabinoid compound.

30. A composition for use as a medicament in treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, wherein the composition comprises a pharmaceutically effective amount of a PPAR $\gamma$  activator.

31. Use of a composition for the manufacture of a medicament for use in treating a subject having a disorder associated with peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ) function, wherein the composition comprises a pharmaceutically effective amount of a PPAR $\gamma$  activator.

\* \* \* \* \*



**ACTUAL PATENT URL LINK:**

**[https://patentimages.storage.googleapis.com/  
03/5f/af/95650b42a64f76/US20040186166A1.p  
df](https://patentimages.storage.googleapis.com/03/5f/af/95650b42a64f76/US20040186166A1.pdf)**


**PATENT CURRENT ASSIGNEE URL  
LINK:**

**[https://patents.google.com/patent/US2004018  
6166A1/en?q=us2004%2f0186166](https://patents.google.com/patent/US20040186166A1/en?q=us2004%2f0186166)**

### Abstract

The present invention provides for methods and compositions for using cannabinoid compounds, such as ajulemic acid, for the treatment of disorders involving peroxisome proliferator-activated receptor-gamma (PPAR $\gamma$ ). In particular, the invention provides for methods and compositions for treating disorders, such as autoimmune or inflammatory disorders, involving PPAR $\gamma$ , such as rheumatoid arthritis and diabetes.

### Images (7)



### Classifications

A61K31/415 1,2-Diazoles

**US20040186166A1**  
United States

Download PDF Find Prior Art Similar

**Inventor:** Sumner Burstein, J. Chen, Robert Zurier  
**Current Assignee:** University of Massachusetts UMass

**Worldwide applications**  
2003 - [WO](#) [AU](#) [US](#)

- Application US10/741,219 events**
- 2002-12-19 • Priority to US43518402P
  - 2003-12-19 • Application filed by University of Massachusetts UMass
  - 2003-12-19 • Priority to US10/741,219
  - 2004-06-08 • Assigned to MASSACHUSETTS, UNIVERSITY OF
  - 2004-09-23 • Publication of US20040186166A1