



National Registry of Emergency Medical Technicians
Advanced Level Psychomotor Examination

2020-Q1

SUPRAGLOTTIC AIRWAY DEVICE

SGA

Candidate: _____ Examiner: _____
Date: _____ Signature: _____
Device: _____

NOTE: If candidate elects to initially ventilate with BVM attached to reservoir and oxygen, full credit must be awarded for steps denoted by “***” so long as first ventilation is delivered within 30 seconds.

Actual Time Started: _____	Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions	1	
Opens the airway manually	1	
Elevates tongue, inserts simple adjunct [oropharyngeal or nasopharyngeal airway]	1	
NOTE: Examiner now informs candidate no gag reflex is present and patient accepts adjunct		
**Ventilates patient immediately with bag-valve-mask device unattached to oxygen	1	
**Ventilates patient with room air	1	
NOTE: Examiner now informs candidate that ventilation is being performed without difficulty and that pulse oximetry indicates the patient's blood oxygen saturation is 85%		
Attaches oxygen reservoir to bag-valve-mask device and connects to high-flow oxygen regulator [12 – 15 L/minute]	1	
Ventilates patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) with appropriate volumes	1	
NOTE: After 30 seconds, examiner auscultates and reports breath sounds are present and equal bilaterally and medical direction has ordered insertion of a supraglottic airway. The examiner must now take over ventilation.		
Checks/prepares supraglottic airway device	1	
Lubricates distal tip of the device [may be verbalized]	1	
NOTE: Examiner to remove OPA and move out of the way when candidate is prepared to insert device.		
Positions head properly	1	
Performs a tongue-jaw lift	1	
Inserts device to proper depth	1	
Secures device in patient [inflates cuffs with proper volumes and immediately removes syringe or secures strap]	1	
Ventilates patient and confirms proper ventilation [correct lumen and proper insertion depth] by auscultation bilaterally over lungs and over the epigastrium	1	
Adjusts ventilation as necessary [ventilates through additional lumen or slightly withdraws tube until ventilation is optimized]	1	
Verifies proper tube placement by secondary confirmation such as capnography, capnometry, EDD or colorimetric device	1	
NOTE: The examiner must now ask the candidate, “How would you know if you are delivering appropriate volumes with each ventilation?”		
Secures device or confirms that the device remains properly secured	1	
Ventilates patient at proper rate and volume while observing capnography/capnometry and pulse oximeter	1	
Actual Time Ended: _____	TOTAL	18

Critical Criteria

- ___ Failure to initiate ventilations within 30 seconds after taking body substance isolation precautions or interrupts ventilations for greater than 30 seconds at any time
- ___ Failure to take or verbalize appropriate PPE precautions
- ___ Failure to voice and ultimately provide high oxygen concentration [at least 85%]
- ___ Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds)
- ___ Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible]
- ___ Failure to pre-oxygenate patient prior to insertion of the supraglottic airway device
- ___ Failure to insert the supraglottic airway device at a proper depth or location within 3 attempts
- ___ Failure to inflate cuffs properly and immediately remove the syringe
- ___ Failure to secure the strap (if present) prior to cuff inflation
- ___ Failure to confirm that patient is being ventilated properly (correct lumen and proper insertion depth) by auscultation bilaterally over lungs and over the epigastrium
- ___ Insertion or use of any adjunct in a manner dangerous to the patient
- ___ Failure to manage the patient as a competent EMT
- ___ Exhibits unacceptable affect with patient or other personnel
- ___ Uses or orders a dangerous or inappropriate intervention

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.



National Registry of Emergency Medical Technicians®
Emergency Medical Technician Psychomotor Examination

2020-Q1
OB Emergency

PATIENT ASSESSMENT/MANAGEMENT – MEDICAL

Candidate: _____

Examiner: _____

Date: _____

Signature: _____

Scenario # Field Birth

Actual Time Started: _____

	Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions	1	
SCENE SIZE-UP		
Determines the scene/situation is safe	1	
Determines the mechanism of injury/nature of illness	1	
Determines the number of patients	1	
Requests additional EMS assistance if necessary	1	
Considers stabilization of the spine	1	
PRIMARY SURVEY/RESUSCITATION		
Verbalizes the general impression of the patient	1	
Determines responsiveness/level of consciousness (AVPU)	1	
Determines chief complaint/apparent life-threats	1	
Assesses airway and breathing -Assessment (1 point) -Assures adequate ventilation (1 point) -Initiates appropriate oxygen therapy (1 point)	3	
Assesses circulation -Assesses/controls major bleeding (1 point) -Checks pulse (1 point) -Assesses skin [either skin color, temperature or condition] (1 point)	3	
Identifies patient priority and makes treatment/transport decision	1	
HISTORY TAKING		
History of the present illness -Onset (1 point) -Quality (1 point) -Severity (1 point) -Provocation (1 point) -Radiation (1 point) -Time (1 point) -Clarifying questions of associated signs and symptoms related to OPQRST (2 points)	8	
Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) -Medications (1 point) -Last oral intake (1 point)	5	
SECONDARY ASSESSMENT		
Assesses affected body part/system -Cardiovascular -Neurological -Integumentary -Reproductive -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social	5	
VITAL SIGNS		
-Blood pressure (1 point) -Pulse (1 point) -Respiratory rate and quality (1 point each)	4	
States field impression of patient	1	
Interventions [verbalizes proper interventions/treatment]	1	
REASSESSMENT		
Demonstrates how and when to reassess the patient to determine changes in condition	1	
Provides accurate verbal report to arriving EMS unit	1	
Actual Time Ended: _____	TOTAL	42

CRITICAL CRITERIA

- ___ Failure to initiate or call for transport of the patient within 15 minute time limit
- ___ Failure to take or verbalize appropriate PPE precautions
- ___ Failure to determine scene safety before approaching patient
- ___ Failure to voice and ultimately provide appropriate oxygen therapy
- ___ Failure to assess/provide adequate ventilation
- ___ Failure to find or appropriately manage problems associated with airway, breathing, hemorrhage or shock
- ___ Failure to differentiate patient's need for immediate transportation versus continued assessment or treatment at the scene
- ___ Performs secondary examination before assessing and treating threats to airway, breathing and circulation
- ___ Orders a dangerous or inappropriate intervention
- ___ Failure to provide accurate report to arriving EMS unit
- ___ Failure to manage the patient as a competent EMT
- ___ Exhibits unacceptable affect with patient or other personnel
- ___ Uses or orders a dangerous or inappropriate intervention

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.



National Registry of Emergency Medical Technicians®
Emergency Medical Technician Psychomotor Examination

2020-Q1
CPR

CARDIAC ARREST MANAGEMENT / AED

Candidate: _____ Examiner: _____
Date: _____ Signature: _____

Actual Time Started: _____	Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions	1	
Determines the scene/situation is safe	1	
Checks patient responsiveness	1	
Direct assistant to retrieve AED	1	
Requests additional EMS assistance	1	
Checks breathing and pulse simultaneously	1	
NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and pulseless."		
Immediately begins chest compressions [adequate depth and rate; allows the chest to recoil completely]	1	
Performs 2 minutes of high-quality, 1-rescuer adult CPR -Adequate depth and rate (1 point) -Correct compression-to-ventilation ratio (1 point) -Allows the chest to recoil completely (1 point) -Adequate volumes for each breath (1 point) -Minimal interruptions of no more than 10 seconds throughout (1 point)	5	
NOTE: After 2 minutes (5 cycles), candidate assesses patient and second rescuer resumes compressions while candidate operates AED.		
Turns on power to AED	1	
Follows prompts and correctly attaches AED to patient	1	
Stops CPR and ensures all individuals are clear of the patient during rhythm analysis	1	
Ensures that all individuals are clear of the patient and delivers shock from AED	1	
Immediately directs rescuer to resume chest compressions	1	
Actual Time Ended: _____	TOTAL	17

Critical Criteria

- ___ Failure to take or verbalize appropriate PPE precautions
- ___ Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds
- ___ Failure to immediately begin chest compressions as soon as pulselessness is confirmed
- ___ Failure to demonstrate acceptable high-quality, 1-rescuer adult CPR
- ___ Interrupts CPR for more than 10 seconds at any point
- ___ Failure to correctly attach the AED to the patient
- ___ Failure to operate the AED properly
- ___ Failure to deliver shock in a timely manner
- ___ Failure to ensure that all individuals are clear of patient during rhythm analysis and before delivering shock [verbalizes "All clear" and observes]
- ___ Failure to immediately resume compressions after shock delivered
- ___ Failure to manage the patient as a competent EMT
- ___ Exhibits unacceptable affect with patient or other personnel
- ___ Uses or orders a dangerous or inappropriate intervention

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.