



Advanced Level Psychomotor Examination

PATIENT ASSESSMENT - MEDICAL

2025-Q1: Patient Assessment

Candidate: _____

Examiner: _____

Date: _____

Signature: _____

Scenario: _____

Actual Time Started: _____

Possible Points	Points Awarded
-----------------	----------------

Takes or verbalizes appropriate PPE precautions	1	
SCENE SIZE-UP		
Determines the scene/situation is safe	1	
Determines the mechanism of injury/nature of illness	1	
Determines the number of patients	1	
Requests additional help if necessary	1	
Considers stabilization of spine	1	
PRIMARY SURVEY		
Verbalizes general impression of the patient	1	
Determines responsiveness/level of consciousness	1	
Determines chief complaint/apparent life-threats	1	
Assesses airway and breathing -Assessment (1 point) -Assures adequate ventilation (1 point) -Initiates appropriate oxygen therapy (1 point)	3	
Assesses circulation -Assesses/controls major bleeding (1 point) -Assesses skin [either skin color, temperature, or condition] (1 point) -Assesses pulse (1 point)	3	
Identifies priority patients/makes transport decision	1	
HISTORY TAKING AND SECONDARY ASSESSMENT		
History of present illness -Onset (1 point) -Severity (1 point) -Provocation (1 point) -Time (1 point) -Quality (1 point) -Clarifying questions of associated signs and symptoms as related to OPQRST (2 points) -Radiation (1 point)	8	
Past medical history -Allergies (1 point) -Past pertinent history (1 point) -Events leading to present illness (1 point) -Medications (1 point) -Last oral intake (1 point)	5	
Performs secondary assessment [assess affected body part/system or, if indicated, completes rapid assessment] -Cardiovascular -Neurological -Integumentary -Reproductive -Pulmonary -Musculoskeletal -GI/GU -Psychological/Social	5	
Vital signs -Pulse (1 point) -Respiratory rate and quality (1 point each) -Blood pressure (1 point) -AVPU (1 point)	5	
Diagnostics [blood glucose test, pulse oximetry, etc] as indicated	2	
States field impression of patient	1	
Verbalizes treatment plan for patient and calls for appropriate intervention(s) (Intranasal Narcan Administrations)	1	
Transport decision re-evaluated	1	
REASSESSMENT		
Repeats primary survey	1	
Repeats vital signs	1	
Evaluates response to treatments	1	
Repeats secondary assessment regarding patient complaint or injuries	1	
Actual Time Ended: _____		
CRITICAL CRITERIA	TOTAL	48

- _____ Failure to initiate or call for transport of the patient within 15 minute time limit
- _____ Failure to take or verbalize appropriate PPE precautions
- _____ Failure to determine scene safety before approaching patient
- _____ Failure to voice and ultimately provide appropriate oxygen therapy
- _____ Failure to assess/provide adequate ventilation
- _____ Failure to find or appropriately manage problems associated with airway, breathing, hemorrhage or shock [hypoperfusion]
- _____ Failure to differentiate patient's need for immediate transportation versus continued assessment and treatment at the scene
- _____ Does other detailed history or physical examination before assessing and treating threats to airway, breathing, and circulation
- _____ Failure to determine the patient's primary problem
- _____ Orders a dangerous or inappropriate intervention
- _____ Failure to provide for spinal protection when indicated



Medical Responder Psychomotor Examination
CARDIAC ARREST MANAGEMENT / AED

2025-Q1: CPR

Candidate: _____

Examiner: _____

Date: _____

Signature: _____

Actual Time Started: _____

	Possible Points	Points Awarded
--	-----------------	----------------

Takes or verbalizes appropriate PPE precautions	1	
Determines the scene/situation is safe	1	
Checks patient responsiveness	1	
Direct assistant to retrieve AED	1	
Requests additional EMS assistance	1	
Checks breathing and pulse simultaneously	1	
NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and pulseless."		
Immediately begins chest compressions [adequate depth and rate; allows the chest to recoil completely]	1	
Performs 2 minutes of high-quality, 1-rescuer adult CPR <ul style="list-style-type: none">-Adequate depth and rate (1 point)-Correct compression-to-ventilation ratio (1 point)-Allows the chest to recoil completely (1 point)-Adequate volumes for each breath (1 point)-Minimal interruptions of no more than 10 seconds throughout (1 point)	5	
NOTE: After 2 minutes (5 cycles), candidate assesses patient and second rescuer resumes compressions while candidate operates AED.		
Turns on power to AED	1	
Follows prompts and correctly attaches AED to patient	1	
Stops CPR and ensures all individuals are clear of the patient during rhythm analysis	1	
Ensures that all individuals are clear of the patient and delivers shock from AED	1	
Immediately directs rescuer to resume chest compressions	1	
Actual Time Ended: _____	TOTAL	17

Critical Criteria

- ___ Failure to take or verbalize appropriate PPE precautions
- ___ Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds
- ___ Failure to immediately begin chest compressions as soon as pulselessness is confirmed
- ___ Failure to demonstrate acceptable high-quality, 1-rescuer adult CPR
- ___ Interrupts CPR for more than 10 seconds at any point
- ___ Failure to correctly attach the AED to the patient
- ___ Failure to operate the AED properly
- ___ Failure to deliver shock in a timely manner
- ___ Failure to ensure that all individuals are clear of patient during rhythm analysis and before delivering shock [verbalizes "All clear" and observes]
- ___ Failure to immediately resume compressions after shock delivered
- ___ Failure to manage the patient as a competent EMR
- ___ Exhibits unacceptable affect with patient or other personnel
- ___ Uses or orders a dangerous or inappropriate intervention

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.



SUPRAGLOTTIC AIRWAY DEVICE

2025-Q1: SGA

Candidate: _____ Examiner: _____
Date: _____ Signature: _____
Device: _____

NOTE: If candidate elects to initially ventilate with BVM attached to reservoir and oxygen, full credit must be awarded for steps denoted by “**” so long as first ventilation is delivered within 30 seconds.

Actual Time Started: _____	Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions	1	
Opens the airway manually	1	
Elevates tongue, inserts simple adjunct [oropharyngeal or nasopharyngeal airway]	1	
NOTE: Examiner now informs candidate no gag reflex is present and patient accepts adjunct		
**Ventilates patient immediately with bag-valve-mask device unattached to oxygen	1	
**Ventilates patient with room air	1	
NOTE: Examiner now informs candidate that ventilation is being performed without difficulty and that pulse oximetry indicates the patient's blood oxygen saturation is 85%		
Attaches oxygen reservoir to bag-valve-mask device and connects to high-flow oxygen regulator [12 – 15 L/minute]	1	
Ventilates patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) with appropriate volumes	1	
NOTE: After 30 seconds, examiner auscultates and reports breath sounds are present and equal bilaterally and medical direction has ordered insertion of a supraglottic airway. The examiner must now take over ventilation.		
Checks/prepares supraglottic airway device	1	
Lubricates distal tip of the device [may be verbalized]	1	
NOTE: Examiner to remove OPA and move out of the way when candidate is prepared to insert device.		
Positions head properly	1	
Performs a tongue-jaw lift	1	
Inserts device to proper depth	1	
Properly secures device per manufacturer's instructions	1	
Ventilates patient and confirms proper ventilation by auscultation bilaterally over lungs and over the epigastrium	1	
Adjusts ventilation as necessary	1	
Verifies proper tube placement by secondary confirmation such as capnography, capnometry, EDD or colorimetric device	1	
NOTE: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?"		
Secures device or confirms that the device remains properly secured	1	
Ventilates patient at proper rate and volume while observing capnography/capnometry and pulse oximeter	1	
Actual Time Ended: _____	TOTAL	18

Critical Criteria

- ___ Failure to initiate ventilations within 30 seconds after using appropriate PPE or interrupts ventilations for greater than 30 seconds at any time
- ___ Failure to take or verbalize appropriate PPE precautions
- ___ Failure to voice and ultimately provide high oxygen concentration [at least 85%]
- ___ Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds)
- ___ Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible]
- ___ Failure to pre-oxygenate patient prior to insertion of the supraglottic airway device
- ___ Failure to insert the supraglottic airway device at a proper depth or location within 3 attempts
- ___ Failure to properly secure device per manufacturer's instructions
- ___ Failure to confirm that patient is being ventilated properly by auscultation bilaterally over lungs and over the epigastrium
- ___ Insertion or use of any adjunct in a manner dangerous to the patient
- ___ Failure to manage the patient as a competent EMT
- ___ Exhibits unacceptable affect with patient or other personnel
- ___ Uses or orders a dangerous or inappropriate intervention

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.



Cardiac Arrest Management / AED

Candidate Name _____ Date _____

Scenario	Start Time: _____	Points Possible	Points Awarded
*Takes or verbalizes appropriate standard precautions		1	
*Determines the scene/situation is safe		1	
*Determines unresponsiveness		1	
NOTE: The examiner must now inform the candidate, "The patient is unresponsive."			
* Assesses respiratory status (observes the patient and determines the absence of breathing or abnormal breathing - gasping or agonal respirations)		1	
NOTE: The examiner must now inform the candidate, "The patient is apneic," or "The patient has gasping, agonal respirations."			
*Assesses pulse status (no more than 10 seconds)		1	
NOTE: The examiner must now Inform the candidate, "The patient is pulse /e ss."			
*Immediately begins chest compressions		1	
Requests additional EMS response		1	
Performs 2 minutes of high quality, adult CPR			
-*Adequate depth and rate of compressions		1	
-*Correct compression-to ventilation ratio		1	
-*Allows the chest to recoil completely		1	
-*Adequate volumes for each breath		1	
-*Minimize interruptions - no longer than 10 seconds throughout		1	
NOTE: After 2 minutes (5 cycles), patient is assessed. The examiner tells the candidate that the second rescuer w/11 resume compressions while candidate operates the AED.			
Turns on power to AED		1	
* Attaches AED to patient properly		1	
*Initiates analysis of rhythm (stops CPR and ensures individuals are clear of patient)		1	
*Ensures all individuals are standing clear of patient prior to delivering each shock		1	
*Delivers shock per national guidelines		1	
*Directs immediate resumption of CPR		1	
Gathers additional information on event from bystanders (POLST, DNR, Lividity, etc.)		1	
Delivers verbal report to medical control to determine need for transport		1	
Properly packages patient for transport		1	
Passing score is 17 (at least 80%)		Total:	21

Stop Time: _ _ _ _ _ Elapsed Time: _ _ Max Time Allowed: 10 Pass / Fail

Critical Criteria

- Failure to take or verbalize appropriate standard precautions
- Failure to determine the scene/situation is safe
- Failure to determine unresponsiveness
- Failure to assess respiratory status
- Failure to assess pulse status (no more than 10 seconds)
- Failure to immediately begin chest compressions after initially finding the patient has no pulse
- Failure to perform CPR with the adequate depth and rate of compressions
- Failure to perform CPR with the correct compression-to ventilation ratio
- Failure to perform CPR properly to allow the chest to recoil completely
- Failure to perform CPR with the adequate volumes for each breath
- Failure to perform minimize interruptions during CPR (no longer than 10 seconds throughout)
- Failure to properly attach the AED to patient
- Failure to properly initiate analysis of rhythm (stops CPR and ensures individuals are clear of patient)
- Failure to ensure all individuals were standing clear of the patient prior to delivering each shock
- Failure to deliver shock per national standards
- Failure to direct immediate resumption of CPR
- Uses or orders a dangerous or inappropriate intervention
- Failure to manage the patient as a competent EMT
- Exhibits unacceptable affect with patient or other personnel

You must factually document your rationale for checking any of the above critical items on the form

Asterisk denotes Critical Criteria

Failure to perform any Critical Criteria constitutes a failure of this skills station

Evaluator Name: _ _ _ _ _
Printed

Signature