



## Advanced Level Psychomotor Examination

### PATIENT ASSESSMENT - MEDICAL

#### 2026-Q1: Patient Assessment

**Candidate:** \_\_\_\_\_

**Date:** \_\_\_\_\_

**Scenario:** \_\_\_\_\_

**Actual Time Started:** \_\_\_\_\_

**Examiner:** \_\_\_\_\_  
**Signature:** \_\_\_\_\_

**Possible  
Points** \_\_\_\_\_      **Points  
Awarded** \_\_\_\_\_

Takes or verbalizes appropriate PPE precautions	1	
<b>SCENE SIZE-UP</b>		
Determines the scene/situation is safe	1	
Determines the mechanism of injury/nature of illness	1	
Determines the number of patients	1	
Requests additional help if necessary	1	
Considers stabilization of spine	1	
<b>PRIMARY SURVEY</b>		
Verbalizes general impression of the patient	1	
Determines responsiveness/level of consciousness	1	
Determines chief complaint/apparent life-threats	1	
Assesses airway and breathing -Assessment (1 point) -Assures adequate ventilation (1 point) -Initiates appropriate oxygen therapy (1 point)	3	
Assesses circulation -Assesses/controls major bleeding (1 point)   -Assesses skin [either skin color, temperature, or condition] (1 point) -Assesses pulse (1 point)	3	
Identifies priority patients/makes transport decision	1	
<b>HISTORY TAKING AND SECONDARY ASSESSMENT</b>		
History of present illness -Onset (1 point)   -Severity (1 point) -Provocation (1 point)   -Time (1 point) -Quality (1 point)   -Clarifying questions of associated signs and symptoms as related to OPQRST (2 points) -Radiation (1 point)	8	
Past medical history -Allergies (1 point)   -Past pertinent history (1 point)   -Events leading to present illness (1 point) -Medications (1 point)   -Last oral intake (1 point)	5	
Performs secondary assessment [assess affected body part/system or, if indicated, completes rapid assessment] -Cardiovascular   -Neurological   -Integumentary   -Reproductive -Pulmonary   -Musculoskeletal   -GI/GU   -Psychological/Social	5	
Vital signs -Pulse (1 point)   -Respiratory rate and quality (1 point each) -Blood pressure (1 point)   -AVPU (1 point)	5	
Diagnostics [blood glucose test, pulse oximetry, etc] as indicated	2	
States field impression of patient	1	
Verbalizes treatment plan for patient and calls for appropriate intervention(s) (Internasal Narcan Administrations)	1	
Transport decision re-evaluated	1	
<b>REASSESSMENT</b>		
Repeats primary survey	1	
Repeats vital signs	1	
Evaluates response to treatments	1	
Repeats secondary assessment regarding patient complaint or injuries	1	
<b>Actual Time Ended:</b> _____		
<b>CRITICAL CRITERIA</b>	<b>TOTAL</b>	48

- Failure to initiate or call for transport of the patient within 15 minute time limit
- Failure to take or verbalize appropriate PPE precautions
- Failure to determine scene safety before approaching patient
- Failure to voice and ultimately provide appropriate oxygen therapy
- Failure to assess/provide adequate ventilation
- Failure to find or appropriately manage problems associated with airway, breathing, hemorrhage or shock [hypoperfusion]
- Failure to differentiate patient's need for immediate transportation versus continued assessment and treatment at the scene
- Does other detailed history or physical examination before assessing and treating threats to airway, breathing, and circulation
- Failure to determine the patient's primary problem
- Orders a dangerous or inappropriate intervention
- Failure to provide for spinal protection when indicated



## Patient Assessment / Management - Trauma

Candidate Name \_\_\_\_\_ Date \_\_\_\_\_

Scenario \_\_\_\_\_ Start Time: \_\_\_\_\_ Points Possible \_\_\_\_\_ Points Awarded \_\_\_\_\_

*Takes or verbalizes appropriate standard precautions		1	
<b>SCENE SIZE-UP</b>			
*Determines the scene situation is safe (environment, hazards, violence, etc.)		1	
Determines mechanism of injury(MOI)/nature of illness (NOI)		1	
Determines the number of patients		1	
Requests additional help if necessary		1	
*Assess for and provide spinal stabilization when indicated		1	
<b>*PRIMARY ASSESSMENT/ RESUSCITATION (Must be completed prior to the secondary assessment)</b>			
Verbalizes general impression of patient		1	
Determines responsiveness/level of consciousness (AVPU)		1	
*Determines chief complaint/apparent life threats		1	
<b>*Airway</b>	Assesses for problems		1
	Assures patent airway		1
<b>*Breathing</b>	Assesses breathing rate, rhythm, quality, depth		1
	Assures adequate ventilation		1
	Initiates appropriate oxygen therapy		1
	Manages any injury which may compromise breathing/ventilation		1
<b>*Circulation</b>	Assesses pulse rate, rhythm, quality		1
	Assesses/controls major bleeding		1
	Assesses skin (color, temp, moisture, cap refill)		1
	Initiates shock management (positions patient properly, conserves body heat)		1
<b>Vital Signs</b>	-Pulse (1 point)	-Respirations (1 point)	-Blood Pressure (1 point)
* Identifies patient priority and makes treatment/transport decision		1	
Considers notifying hospital for Trauma Alert/Activation per Trauma Triage Tool		1	
<b>HISTORY TAKING</b>			
Obtains S.A.M.P. L.E. history if possible		1	
<b>SECONDARY ASSESSMENT</b>			
Assesses the head	Inspects and palpates the scalp and ears		1
	Assesses the eyes		1
	Assesses the facial area including oral & nasal area		1
Assesses the neck	Inspects and palpates the cervical spine		1
	Assesses for JVD		1
	Assesses for tracheal deviation		1
Assesses the chest	Expose patient in order to: Inspect the chest		1
	Palpate the chest		1
	Auscultate the chest		1
Assesses the abdomen/pelvis	Inspects and palpates the abdomen		1
	Assesses and palpates the pelvic girdle		1
	Verbalizes assessment of genitalia/perineum as needed		1
Assesses the extremities	(1 point for each extremity)		4
	Expose patient in order to: Inspect, palpate, and assess distal pulses, capillary refill, sensation and movement		
Assesses the posterior	Inspects and palpates thorax/thoracic vertebrae		1
	Inspects and palpates lower back/lumbar vertebrae		1
	Inspects and palpates flank areas		1

	Points Possible	Points Awarded
Manages secondary injuries and wounds appropriately	1	
<b>REASSESSMENT</b>		
Demonstrates reassessment of the patient appropriate to patient condition	1	
*Provides accurate verbal report to arriving EMS unit or receiving facility	1	
<b>Passing score is 38 (at least 80%)</b>	<b>TOTAL:</b>	<b>47</b>

**Stop Time:** \_\_\_\_\_ **Elapsed Time:** \_\_\_\_\_ **Max Time Allowed: 10** **Pass/ Fail**

**Critical Criteria**

- Failure to take or verbalize proper standard precautions
- Failure to determine scene/situation safety before approaching patient
- Failure to assess for and provide spinal stabilization when indicated
- Failure to complete primary assessment prior to the secondary assessment
- Failure to determine chief complain apparent life threats
- Failure to find problems associated with airway, breathing, hemorrhage or shock
- Failure to appropriately manage problems associated with airway, breathing, hemorrhage or shock
- Failure to identify patient priority and make appropriate treatment transport decision
- Failure to initiate or call for transport of the patient within 10 minute time limit
- Uses or orders a dangerous or inappropriate intervention
- Failure to provide an accurate report to arriving EMS unit or receiving facility
- Failure to manage the patient as a competent EMT
- Exhibits unacceptable affect with patient or other personnel

**You must factually document your rationale for checking any of the above critical items on the form**

**Asterisk denotes Critical Criteria**

**Failure to perform any Critical Criteria constitutes a failure of this skill station**

**Evaluator Name:** \_\_\_\_\_

Printed

\_\_\_\_\_

Signature



**Medical Responder Psychomotor Examination  
CARDIAC ARREST MANAGEMENT / AED**

**2026-Q1: CPR**

Candidate: \_\_\_\_\_

Examiner: \_\_\_\_\_

Date: \_\_\_\_\_

Signature: \_\_\_\_\_

Actual Time Started: \_\_\_\_\_

**Possible  
Points**      **Points  
Awarded**

Takes or verbalizes appropriate PPE precautions	1	
Determines the scene/situation is safe	1	
Checks patient responsiveness	1	
Direct assistant to retrieve AED	1	
Requests additional EMS assistance	1	
Checks breathing and pulse simultaneously	1	
<b>NOTE: After checking responsiveness, then checking breathing and pulse for no more than 10 seconds, examiner informs candidate, "The patient is unresponsive, apneic and pulseless."</b>		
Immediately begins chest compressions [adequate depth and rate; allows the chest to recoil completely]	1	
Performs 2 minutes of high-quality, 1-rescuer adult CPR -Adequate depth and rate (1 point) -Correct compression-to-ventilation ratio (1 point) -Allows the chest to recoil completely (1 point) -Adequate volumes for each breath (1 point) -Minimal interruptions of no more than 10 seconds throughout (1 point)	5	
<b>NOTE: After 2 minutes (5 cycles), candidate assesses patient and second rescuer resumes compressions while candidate operates AED.</b>		
Turns on power to AED	1	
Follows prompts and correctly attaches AED to patient	1	
Stops CPR and ensures all individuals are clear of the patient during rhythm analysis	1	
Ensures that all individuals are clear of the patient and delivers shock from AED	1	
Immediately directs rescuer to resume chest compressions	1	

Actual Time Ended: \_\_\_\_\_

**TOTAL**      **17**

**Critical Criteria**

- Failure to take or verbalize appropriate PPE precautions
- Failure to check responsiveness, then check breathing and pulse simultaneously for no more than 10 seconds
- Failure to immediately begin chest compressions as soon as pulselessness is confirmed
- Failure to demonstrate acceptable high-quality, 1-rescuer adult CPR
- Interrupts CPR for more than 10 seconds at any point
- Failure to correctly attach the AED to the patient
- Failure to operate the AED properly
- Failure to deliver shock in a timely manner
- Failure to ensure that all individuals are clear of patient during rhythm analysis and before delivering shock [verbalizes "All clear" and observes]
- Failure to immediately resume compressions after shock delivered
- Failure to manage the patient as a competent EMR
- Exhibits unacceptable affect with patient or other personnel
- Uses or orders a dangerous or inappropriate intervention

**You must factually document your rationale for checking any of the above critical items on the reverse side of this form.**



## SUPRAGLOTTIC AIRWAY DEVICE

### 2026-Q1: SGA

Candidate: \_\_\_\_\_ Examiner: \_\_\_\_\_  
 Date: \_\_\_\_\_ Signature: \_\_\_\_\_  
 Device: \_\_\_\_\_

NOTE: If candidate elects to initially ventilate with BVM attached to reservoir and oxygen, full credit must be awarded for steps denoted by \*\*\* so long as first ventilation is delivered within 30 seconds.

Actual Time Started: \_\_\_\_\_

Possible Points \_\_\_\_\_  
 Points Awarded \_\_\_\_\_

Takes or verbalizes appropriate PPE precautions	1	
Opens the airway manually	1	
Elevates tongue, inserts simple adjunct [oropharyngeal or nasopharyngeal airway]	1	
<b>NOTE: Examiner now informs candidate no gag reflex is present and patient accepts adjunct</b>		
**Ventilates patient immediately with bag-valve-mask device unattached to oxygen	1	
**Ventilates patient with room air	1	
<b>NOTE: Examiner now informs candidate that ventilation is being performed without difficulty and that pulse oximetry indicates the patient's blood oxygen saturation is 85%</b>		
Attaches oxygen reservoir to bag-valve-mask device and connects to high-flow oxygen regulator [12 – 15 L/minute]	1	
Ventilates patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) with appropriate volumes	1	
<b>NOTE: After 30 seconds, examiner auscultates and reports breath sounds are present and equal bilaterally and medical direction has ordered insertion of a supraglottic airway. The examiner must now take over ventilation.</b>		
Checks/prepares supraglottic airway device	1	
Lubricates distal tip of the device [may be verbalized]	1	
<b>NOTE: Examiner to remove OPA and move out of the way when candidate is prepared to insert device.</b>		
Positions head properly	1	
Performs a tongue-jaw lift	1	
Inserts device to proper depth	1	
Properly secures device per manufacturer's instructions	1	
Ventilates patient and confirms proper ventilation by auscultation bilaterally over lungs and over the epigastrium	1	
Adjusts ventilation as necessary	1	
Verifies proper tube placement by secondary confirmation such as capnography, capnometry, EDD or colorimetric device	1	
<b>NOTE: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?"</b>		
Secures device or confirms that the device remains properly secured	1	
Ventilates patient at proper rate and volume while observing capnography/capnometry and pulse oximeter	1	
Actual Time Ended: _____	<b>TOTAL</b>	<b>18</b>

#### Critical Criteria

- Failure to initiate ventilations within 30 seconds after using appropriate PPE or interrupts ventilations for greater than 30 seconds at any time
- Failure to take or verbalize appropriate PPE precautions
- Failure to voice and ultimately provide high oxygen concentration [at least 85%]
- Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds)
- Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible]
- Failure to pre-oxygenate patient prior to insertion of the supraglottic airway device
- Failure to insert the supraglottic airway device at a proper depth or location within 3 attempts
- Failure to properly secure device per manufacturer's instructions
- Failure to confirm that patient is being ventilated properly by auscultation bilaterally over lungs and over the epigastrium
- Insertion or use of any adjunct in a manner dangerous to the patient
- Failure to manage the patient as a competent EMT
- Exhibits unacceptable affect with patient or other personnel
- Uses or orders a dangerous or inappropriate intervention

**You must factually document your rationale for checking any of the above critical items on the reverse side of this form.**