



Failure to provide for spinal protection when indicated

National Registry of Emergency Medical Technicians Advanced Level Psychomotor Examination

PATIENT ASSESSMENT - MEDICAL

2024-Q3: Patient Assess	sment	Examiner:		
Candidate:		0:		
Date:			Possible	Points
			Points	Awarded
Scenario: Actual Time Started:				
Actual Time Started.	_			
Takes or verbalizes appropriate P	PE precautions		1	T
SCENE SIZE-UP				
Determines the scene/situation is			1	
Determines the mechanism of inju	,		1	
Determines the number of patient			1	
Requests additional help if necess	sary		1	
Considers stabilization of spine			1	
PRIMARY SURVEY				
Verbalizes general impression of			1	
Determines responsiveness/level			1	1
Determines chief complaint/appar	ent life-threats		1	
Assesses airway and breathing				
-Assessment (1 point)			3	
-Assures adequate ventil				
-Initiates appropriate oxyg	gen therapy (1 point)			
Assesses circulation		11 5 74 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		
_		es skin [either skin color, temperature, or condition] (1 point)) 3	
-Assesses pulse (1 point)				
Identifies priority patients/makes t			1 1	
HISTORY TAKING AND SECONI	JARY ASSESSMENT			
History of present illness	0			
-Onset (1 point)	-Severity (1 point)			
-Provocation (1 point)	-Time (1 point)		8	
-Quality (1 point)	-Clarifying questions of ass	sociated signs and symptoms as related to OPQRST (2 points)	its)	
-Radiation (1 point)				-
Past medical history	Doot worth out history (4 m	Secretaria di ante agreca d'ille es d'anciett	_	
-Allergies (1 point)	-Past pertinent history (1 po	oint) -Events leading to present illness (1 point)	5	
-Medications (1 point)	-Last oral intake (1 point)	return on if in directed accomplates would accompand		
-Cardiovascular		/stem or, if indicated, completes rapid assessment] -Integumentary -Reproductive	E	
	-Neurological -Musculoskeletal	•	5	
-Pulmonary	-iviusculoskeletai	-GI/GU -Psychological/Social		1
Vital signs	Despiratory rate on	d quality (1 point apply)	E	
-Pulse (1 point)		d quality (1 point each)	5	
-Blood pressure (1 point)	· ' '	d	2	+
Diagnostics [blood glucose test, p States field impression of patient	uise oximetry, etcj as indicate	eu .	1	-
	ent and calls for appropriate i	ntervention(s) (Internasal Narcan Administrations)	1	+
Transport decision re-evaluated	and cans for appropriate in	intervention(s) (internasai Narcan Administrations)	1	-
REASSESSMENT				
Repeats primary survey			1 1	1
Repeats vital signs			1	-
Evaluates response to treatments			1	-
Repeats secondary assessment r		iniuries	1	
Actual Time Ended:	_	njunes		
CRITICAL CRITERIA		то	TAL 48	
Failure to initiate or call for	transport of the patient within	n 15 minute time limit		on
	appropriate PPE precautions			
Failure to determine scene	safety before approaching pa	atient		
	tely provide appropriate oxyge	en therapy		
Failure to assess/provide a				
		ated with airway, breathing, hemorrhage or shock [hypoperf		
		sportation versus continued assessment and treatment at the		
-		re assessing and treating threats to airway, breathing, and o	rculation	
Failure to determine the pa				
Orders a dangerous or inar	opropriate intervention			



National Registry of Emergency Medical Technicians® Emergency Medical Responder Psychomotor Examination CARDIAC ARREST MANAGEMENT / AED

2024-Q3: CPR

Candidate:	Examiner:		
Date:	Signature:		
Actual Time Started:		Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions		1	
Determines the scene/situation is safe		1	
Checks patient responsiveness		1	
Direct assistant to retrieve AED		1	
Requests additional EMS assistance		1	
Checks breathing and pulse simultaneously		1	
NOTE: After checking responsiveness, then checking breathing a examiner informs candidate, "The patient is unresponsive, apneic		ds,	
Immediately begins chest compressions [adequate depth and rate; allow	s the chest to recoil completely]	1	
Performs 2 minutes of high-quality, 1-rescuer adult CPR -Adequate depth and rate (1 point) -Correct compression-to-ventilation ratio (1 point) -Allows the chest to recoil completely (1 point) -Adequate volumes for each breath (1 point) -Minimal interruptions of no more than 10 seconds throughout (1 point)	nt)	5	
NOTE: After 2 minutes (5 cycles), candidate assesses patient and candidate operates AED.	second rescuer resumes compressi	ions while	
Turns on power to AED		1	
Follows prompts and correctly attaches AED to patient		1	
Stops CPR and ensures all individuals are clear of the patient during rhy		1	
Ensures that all individuals are clear of the patient and delivers shock from	om AED	1	
Immediately directs rescuer to resume chest compressions		1	
Actual Time Ended:	TOTAL	17	
Critical Criteria Failure to take or verbalize appropriate PPE precautions Failure to check responsiveness, then check breathing and pulse selection immediately begin chest compressions as soon as pulse Failure to demonstrate acceptable high-quality, 1-rescuer adult CPI Interrupts CPR for more than 10 seconds at any point Failure to correctly attach the AED to the patient Failure to operate the AED properly Failure to deliver shock in a timely manner Failure to ensure that all individuals are clear of patient during rhyth [verbalizes "All clear" and observes] Failure to immediately resume compressions after shock delivered Failure to manage the patient as a competent EMR Exhibits unacceptable affect with patient or other personnel Uses or orders a dangerous or inappropriate intervention	lessness is confirmed R	nds	

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.



National Registry of Emergency Medical Technicians Advanced Level Psychomotor Examination

SUPRAGLOTTIC AIRWAY DEVICE

2024-Q3:SGA

Candidate:	Examiner:			
Date:				
Device:	<u></u>			
NOTE: If candidate elects to initially ventilate with BVM attached to re "**" so long as first ventilation is delivered within 30 seconds		ed for steps o	denoted by	
so long as first vertiliation is delivered within 30 seconds		Possible	Points	
Actual Time Started:		Points	Awarded	
Takes or verbalizes appropriate PPE precautions		1		
Opens the airway manually		1		
Elevates tongue, inserts simple adjunct [oropharyngeal or nasopharyngeal or nasophar	yngeal airway]	1		
NOTE: Examiner now informs candidate no gag reflex is present				
**Ventilates patient immediately with bag-valve-mask device unattact	ched to oxygen	1		
**Ventilates patient with room air		1		
NOTE: Examiner now informs candidate that ventilation is being the patient's blood oxygen saturation is 85%	g performed without difficulty and that pul	se oximetry	indicates	
Attaches oxygen reservoir to bag-valve-mask device and connects	o high-flow oxygen regulator		I	
[12 – 15 L/minute]	3 1 1 7 3 1 1 1 3 1 1 1 1	1		
Ventilates patient at a rate of 10 – 12/minute (1 ventilation every 5 –	6 seconds) with appropriate volumes	1		
NOTE: After 30 seconds, examiner auscultates and reports brea		and medical	direction	
has ordered insertion of a supraglottic airway. The examiner m	ust now take over ventilation.			
Checks/prepares supraglottic airway device		1		
Lubricates distal tip of the device [may be verbalized]		1		
NOTE: Examiner to remove OPA and move out of the way when	candidate is prepared to insert device.			
Positions head properly		1		
Performs a tongue-jaw lift		1		
Inserts device to proper depth		1		
Properly secures device per manufacturer's instructions		1		
Ventilates patient and confirms proper ventilation by auscultation bil	aterally over lungs and over the	4		
epigastrium		1		
Adjusts ventilation as necessary		1		
Verifies proper tube placement by secondary confirmation such as of	apnography, capnometry, EDD or	1		
colorimetric device		1		
NOTE: The examiner must now ask the candidate, "How would each ventilation?"	you know if you are delivering appropriate	volumes wi	ith	
Secures device or confirms that the device remains properly secure	d	1		
Ventilates patient at proper rate and volume while observing capnot		1		
Actual Time Ended:	TOTAL	18		
Critical Criteria	The PDF and the second and the secon	. 00		
 Failure to initiate ventilations within 30 seconds after using appropriance. Failure to take or verbalize appropriate PPE precautions 	ate PPE or interrupts ventilations for greater tha	n 30 seconds	at any time	
Failure to voice and ultimately provide high oxygen concentration [a	t least 85%]			
Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilate				
Failure to provide adequate volumes per breath [maximum 2 errors/				
Failure to pre-oxygenate patient prior to insertion of the supraglottic				
Failure to insert the supraglottic airway device at a proper depth or I Failure to properly secure device per manufacturer's instructions	ocation within 3 attempts			
Failure to confirm that patient is being ventilated properly by auscult	ation bilaterally over lungs and over the epigastri	ium		
Insertion or use of any adjunct in a manner dangerous to the patien				
Failure to manage the patient as a competent EMT				
Exhibits unacceptable affect with patient or other personnel				
Uses or orders a dangerous or inappropriate intervention				

You must factually document your rationale for checking any of the above critical items on the reverse side of this form.





Patient Assessment / Management - Trauma

Candidate Name	Date		
ScenarioStart Time:		Points Possible	Points Awarded
*Takes or verbalizes app	ropriate standard precautions	1	
SCENE SIZE-UP			•
*Determines the scene s	tuation is safe (environment, hazards, violence, etc.)	1	
Determines mechanism	of injury(MOI)/nature of illness (NOi)	1	
Determines the number	of patients	1	
Requests additional help	if necessary	1	
	spinal stabilization when indicated NT/ RESUSCITATION (Must be completed prior to the secondary a	1	
Verbalizes general impre		1	
		1	
	ess/level of consciousness (AVPU)	1	
*Determines chief compla	Assesses for problems	1	
*Airway	Assures patent airway	1	
	Assesses breathing rate, rhythm, quality, depth	1	
*Breathing	Assures adequate ventilation	1	
	Initiates appropriate oxygen therapy Manages any injury which may compromise breathing/ventilation	1 1	
	Assesses pulse rate, rhythm, quality	1	
*Circulation	Assesses/controls major bleeding Assesses skin (color, temp, moisture, cap refill)	1	
Circulation	Initiates shock management (positions patient properly, conserves body heat)	1	
Vital Signs -Puls	se (1 point) -Respirations (1 point) -Blood Pressure (1 point)	3	
* Identifies patient priority	and makes treatment/transport decision	1	
Considers notifying hospi	tal for Trauma Alert/Activation per Trauma Triage Tool	1	
HISTORY TAKING		·	
Obtains S A.M.P. L.E. his	story if possible	1	
SECONDARY ASSESS	MENT	l	
Assesses the head	Inspects and palpates the scalp and ears	1	
	Assesses the eyes	1	
	Assesses the facial area including oral & nasal area	1	
Assesses the neck	Inspects and palpates the cervical spine Assesses for JVD	1	
	Assesses for tracheal deviation	1 1	
Assesses the chest	Exposes patient in order to: Inspect the chest	1	
	Palpate the chest	1	
	Auscultate the chest	1	
Assesses the	Inspects and palpates the abdomen	1	
abdomen/pelvis	Assesses and palpates the pelvic girdle Verbalizes assessment of genitalia/perineum as needed	1	
Aggregate -		1	
Assesses the extremities	(1 point for each extremity) Exposes patient in order to: Inspect, palpate, and assess distal	4	
	pulses, capillary refill, sensation and movement		
Assesses the posterior	Inspects and palpates thorax/thoracic vertebrae	1	
	Inspects and palpates lower back/lumbar vertebrae Inspects and palpates flank areas	1	
	moposto ana parpatoo nam aroas	1	

DOH 530-093 January 2013 Page 19

	Points Possible	Points Awarded
Manages secondary injuries and wounds appropriately	1	
REASSESSMENT		
Demonstrates reassessment of the patient appropriate to patient condition	1	
*Provides accurate verbal report to arriving EMS unit or receiving facility	1	
Passing score is 38 (at least 80%) TOTAL:	47	

Stop Time:	Elapsed Time:	_	Max Time Allowed: 10	Pass/ Fail
Critical Criteria				
Failure to determine so Failure to assess for ar Failure to complete print Failure to determine che Failure to find problems Failure to appropriately Failure to identify patient Failure to initiate or call Uses or orders a danger Failure to provide an action of Failure to manage the provide and failure to manag	• .	e approation whee the second threats breathin stated without within ervention EMS united the second to the second the sec	en indicated ondary assessment g, hemorrhage or shock th airway, breathing, hemorreatment transport decision 10 minute time limit tor receiving facility	•
You must factually document Asterisk denotes Critical Crit Failure to perform any Critica	eria			ns on the form
Evaluator Name:				

Printed

Signature