



PANLEUKOPENIA

FELINE DISTEMPER

HOME DISINFECTION GUIDE

STEP 5: AIR OUT AND QUARANTINE

- Allow the space to dry completely and air out for 24 hours.
- Quarantine recovered or new cats for at least 2-3 weeks.
- Avoid bringing unvaccinated kittens or cats into the area during this period.

STEP 6: VACCINATION AND PREVENTION

- Ensure all cats and kittens are fully vaccinated with FVRCP before entering the cleaned area.
- Boosters should be up to date, especially for fosters or new intakes.

BIOSECURITY PROTOCOLS

Maintaining proper biosecurity is essential to prevent the spread of infectious diseases within the environment. These protocols outline how to safely manage cleaning and disinfection procedures, especially following exposure to highly contagious illnesses such as feline panleukopenia.

GENERAL BIOSECURITY PRINCIPLES

- Always wear appropriate personal protective equipment (PPE) such as gloves, gowns, and shoe covers when cleaning.
- Handwashing is required before and after handling each animal, cleaning an area, or touching contaminated items.
- Clean from low-risk areas to high-risk areas (healthy kittens → isolation → intake → quarantine).
- Use separate cleaning supplies (mops, buckets, rags) for each section to avoid cross-contamination.