

2020 HIV DRUG CHART

This quick-reference chart compares antiretroviral (ARV) options for the treatment of HIV, including adult dosing and dietary restrictions. Visit poz.com/drugchart for more info.

*Generic version available in the U.S. (Pills not shown actual size)

Complete Regimens



ATRIPLA
efavirenz + tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 600 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take on an empty stomach. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



BIKTARVY
bictegravir + tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 50 mg bictegravir + 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with or without food.



CABENUVA
cabotegravir + rilpivirine
A long-acting injectable regimen administered as two intramuscular injections every four weeks after a one-month lead-in period with oral Vocabria (cabotegravir) + Edurant (rilpivirine). Take with food.



COMPLERA
rilpivirine + tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 25 mg rilpivirine + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with a meal.



DELSTRIGO
doravirine + tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 100 mg doravirine + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



DOVATO
dolutegravir + lamivudine
One tablet once a day. Each tablet contains 50 mg dolutegravir + 300 mg lamivudine. Take with or without food.



GENVOYA
elvitegravir + cobicistat + tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 150 mg elvitegravir + 150 mg cobicistat + 10 mg tenofovir alafenamide + 200 mg emtricitabine. Take with food.

Nucleoside/Nucleotide Reverse Transcriptase Inhibitors (NRTIs, or nukes)



CIMDUO
tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



COMBIVIR *
zidovudine + lamivudine
One tablet twice a day. Each tablet contains 300 mg zidovudine + 150 mg lamivudine. Take with or without food.



DESCOXY
tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with or without food.



EMTRIVA
emtricitabine (also known as FTC)
One 200 mg capsule once a day. Take with or without food.



EPIVIR *
lamivudine (also known as 3TC)
One 300 mg tablet once a day, or one 150 mg tablet twice a day. Take with or without food. Also approved for the treatment of hepatitis B virus but at a lower dose. People living with both viruses should use the HIV dose.



EPZICOM *
abacavir + lamivudine
One tablet once a day. Each tablet contains 600 mg abacavir + 300 mg lamivudine. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.



RETROVIR *
zidovudine (also known as AZT)
One 300 mg tablet twice a day. Take with or without food.



TEMIXYS
tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



TRUVADA
tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with or without food.



VIREAD *
tenofovir disoproxil fumarate
One 300 mg tablet once a day. Take with or without food.

Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs, or non-nukes)



EDURANT
rilpivirine
One 25 mg tablet once a day. Take with food.



INTELENCE
etravirine
One 200 mg tablet twice a day. Take with food.



PIFELTRO
doravirine
One 100 mg tablet once a day. Take with or without food.



SUSTIVA *
efavirenz
One 600 mg tablet once a day, or three 200 mg capsules once a day. Take on an empty stomach or with a low-fat snack. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



VIRAMUNE *
nevirapine
One 200 mg Viramune immediate release (IR) tablet once a day for the first 14 days, then one 400 mg Viramune extended release (XR) tablet once a day. Take with or without food.

Protease Inhibitors (PIs)



EVOTAZ
atazanavir + cobicistat
One tablet once a day. Each tablet contains 300 mg atazanavir + 150 mg cobicistat. Take with food.



KALETRA
lopinavir + ritonavir
Two tablets twice a day, or four tablets once a day, depending on HIV drug resistance. Each tablet contains 200 mg lopinavir + 50 mg ritonavir. Take with or without food.



PREZCOBIX
darunavir + cobicistat
One tablet once a day. Each tablet contains 800 mg darunavir + 150 mg cobicistat. Take with food.



PREZISTA
darunavir
One 800 mg tablet (or two 400 mg tablets) plus one 100 mg Norvir tablet, or one 600 mg tablet plus one 100 mg Norvir tablet twice a day, depending on drug resistance. Take with food.



REYATAZ *
atazanavir
Two 200 mg capsules once a day, or one 300 mg capsule plus one 100 mg Norvir tablet once a day. Take with food.

**JULUCA****dolutegravir + rilpivirine**

One tablet once a day. Each tablet contains 50 mg dolutegravir + 25 mg rilpivirine. Take with a meal.

**ODEFSEY****rilpivirine + tenofovir alafenamide + emtricitabine**

One tablet once a day. Each tablet contains 25 mg rilpivirine + 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with a meal.

**STRIBILD****elvitegravir + cobicistat + tenofovir disoproxil fumarate + emtricitabine**

One tablet once a day. Each tablet contains 150 mg elvitegravir + 150 mg cobicistat + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with food.

**SYMFI AND SYMFI LO****efavirenz + tenofovir disoproxil fumarate + lamivudine**

One tablet of either Symfi or Symfi Lo once a day. Each tablet of Symfi contains 600 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Each tablet of Symfi Lo (above) contains 400 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take on an empty stomach. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.

**SYM TUZA****darunavir + cobicistat + tenofovir alafenamide + emtricitabine**

One tablet once a day. Each tablet contains 800 mg darunavir + 150 mg cobicistat + 10 mg tenofovir alafenamide + 200 mg emtricitabine. Take with food.

**TRIUMEQ****dolutegravir + abacavir + lamivudine**

One tablet once a day. Each tablet contains 50 mg dolutegravir + 600 mg abacavir + 300 mg lamivudine. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.

**ZIAGEN *****abacavir**

One 300 mg tablet twice a day, or two 300 mg tablets once a day. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.

**RUKOBIA****fostemsavir**

One 600 mg tablet twice a day for people with HIV treatment experience. Take with or without food.

**SELZENTRY****maraviroc**

One 150 mg, 300 mg or 600 mg tablet twice a day, depending on other meds used, for people with HIV treatment experience. Take with or without food.

**TROGARZO****ibalizumab**

A long-acting injectable administered intravenously as a single loading dose of 2,000 mg followed by maintenance dose of 800 mg every two weeks for people with HIV treatment experience.

**NORVIR *****ritonavir**

Six 100 mg tablets twice a day. The full dose of Norvir is rarely used. It is most often used at lower doses to boost the levels of other ARVs in the blood. Take with food.

**TYBOST****cobicistat**

One 150 mg tablet once a day in combination with ARVs that require boosting. Used only to boost other drugs. Take with food.

**ISENTRESS****raltegravir**

Two 600 mg Isentress HD tablets (above) once a day for those who are treatment naive or whose virus has been suppressed on an initial regimen of Isentress. One 400 mg Isentress tablet twice daily for people with HIV treatment experience. Take with or without food.


**TIVICAY****dolutegravir**

One 50 mg tablet once a day for those first starting ARV therapy or for those who have not used an integrase inhibitor in the past. One 50 mg tablet twice a day for treatment-experienced individuals who have HIV that is resistant to other integrase inhibitors and when taken with certain ARVs. Take with or without food.

**VOCABRIA****cabotegravir**

One 30mg tablet taken once a day with once-daily Edurant for a month as a lead-in regimen before switching to Cabenuva injections or for short-term treatment. Take with food.

These antiretroviral medications are rarely prescribed and no longer recommended:

**APTIVUS**
tipranavir**CRIVAN**
indinavir**FUZEON**
enfuvirtide**INVIRASE**
saquinavir**LEXIVA**
fosamprenavir**TRIZIVIR ***
abacavir + zidovudine + lamivudine**VIRACEPT**
nelfinavir**ZERIT**
stavudine



A HEALTHIER LIFE CAN START WITH HIV TREATMENT.



Starting HIV treatment as soon as possible helps stop the damage HIV causes to your body. Plus, doctors and scientists have found that it can help reduce the risk of some infections, certain cancers, and even AIDS.

TREATMENT HELPS PREVENT THE SPREAD OF HIV.

Starting and sticking to HIV treatment can lower the amount of virus in the body so much, it can't be measured by a test. It's called being undetectable.


According to current research, *sticking to daily treatment and staying undetectable means there's basically no risk of spreading HIV through sex.*

HIV is still in the body, and being undetectable doesn't prevent other STIs. So use condoms and practice safer sex.



TALK TO YOUR HEALTHCARE PROVIDER.

Have an open conversation. There's no cure for HIV, but when you work together it helps your healthcare provider find the treatment that's right for you.



Watch HIV: "Treat 2 Prevent"

See how staying on treatment can help protect you and the people you care about.

[YouTube.com/HelpStopTheVirus](https://www.youtube.com/HelpStopTheVirus)

STOPPING THE VIRUS CAN START WITH YOU.

Here are two resources that can help.

STOP THE VIRUS.

Watch videos, share information, and see how we can all help stop the virus.

HelpStopTheVirus.com

[YouTube.com/HelpStopTheVirus](https://www.youtube.com/HelpStopTheVirus)

HelpStopTheVirus.Tumblr.com



Get the answers you need,
privately, on your phone.

HIVanswers.com/app



HIV ANSWERS, the HIV ANSWERS Logo, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc. All other marks are the property of their respective owners.
© 2018 Gilead Sciences, Inc. All rights reserved. UNBC5702 06/18

START HIV TREATMENT. HELP PROTECT YOUR HEALTH.



There is no cure for HIV, but find out how treatment helps make it possible to live a healthier life.

See
Inside

