

Author: [REDACTED] MD

Service: Pulmonology

Author Type: Physician

Filed: 7/5/2016 4:17 PM

Date of Service: 7/5/2016 4:12 PM

Status: Addendum

Editor: [REDACTED] MD (Physician)

Related Notes: Original Note by [REDACTED] MD (Physician) filed at 7/5/2016 4:14 PM

Pre-procedure Diagnoses:

1. Recurrent left pleural effusion [J90]

Post-procedure Diagnoses:

1. Recurrent left pleural effusion [J90]

Procedures:

1. THORACENTESIS [PRO306 (Custom)]

[REDACTED]
[REDACTED] HOSPITAL

Interventional Pulmonology

[REDACTED], MD, MEd, Director

Thoracentesis Procedure Note

Date of procedure: 7/5/2016

I was asked to perform a left thoracentesis. Informed consent was obtained prior to the start of the procedure. A time out for patient safety was performed. Ultrasound (US) was then performed to identify the pleural effusion and to distinguish it from the surrounding structures, including the lung, effusion, diaphragm and subdiaphragmatic structures. A mark was placed in the most appropriate position for safely guiding the catheter into the fluid based on the ultrasound images.

Thereafter, the area was prepped in sterile fashion. 1% lidocaine was used to anesthetize the skin and pleura. A small incision was made and subsequently a 6 French catheter was placed into the pleural space using the Safe-T-Centesis kit from Carefusion. The effusion was drained as much as possible. A band aide was placed over the thoracentesis site at the conclusion of the procedure.

The amount of fluid removed was 1200 ml

The color of the fluid was yellow

Specimens were not sent for analysis.

Additional participants in the procedure included: [REDACTED] MD

Thank you for allowing interventional pulmonology to participate in this patient's care. Please feel free to call the IP pager [REDACTED] with any questions or concerns.

[REDACTED] MD, MEd

Director, Interventional Pulmonology

Division of Pulmonary, Critical Care and Sleep Medicine

[REDACTED] School of Medicine

[REDACTED] Hospital

Office: [REDACTED]