

WARNING

KEEP OUT OF REACH OF CHILDREN

DO NOT SWALLOW. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Users are encouraged to seek further advice.

STORAGE:

DO NOT STORE product in direct sunlight.

DO NOT STORE product in temperature below 5°C



Bang P™

NPKS (13-19-0-0) + Organic Catalysts



Scan for
product SDS

$N-P_2O_5-K_2O$ 13-43-0

Analysis:	w/v	(weight/volume)
Product:	pH	6.1-7.0
Specific Gravity:		1.34

GUARANTEED ANALYSIS

Nitrogen	N	as ammonium	13.0%	130 g/L
Phosphorus	P	as polyphosphate	19.0%	190 g/L
		as P_2O_5	43.6%	436 g/L
Catalysts	OC	as proprietary	1%	10 g/L

Distributed by Advanced Nutrients Pty Ltd
Address: 27 Westbourne Drive, Wights Mountain, QLD 4520
Phone: 1800 244 009
Email: info@advancednutrients.com.au
Web: www.advancednutrients.com.au

This label complies with the National Code of Practice for fertiliser description and labelling.

BROADACRE	RATE L/Ha	WATER	COMMENTS
Cotton - Injection	10-30	1:10-20	Apply at plant with water injection
Cotton - Foliar	5-15	1:20	Apply as foliar at 6-8 leaf stage
Cereals - Injection	10-30	1:2-4	Apply at plant for P-retentive soils
Cereals - Foliar	5-15	1:10-20	Apply as foliar at early tillering
Legumes	5-15	1:10-20	Apply 3-4 weeks after emergence
Canola	5-15	1:10-20	Apply at 4-6 leaf to early cabbage
Maize / Sorghum	5-15	1:10-20	Apply 4-6 leaf up to flowering
Pasture	10-15	1:10-20	Apply at Spring and Autumn flush
SUGAR	RATE L/Ha	WATER	COMMENTS
Plant Cane	15	Min 100 L/Ha	Apply with liquid fungicide at planting
Ratoon Cane	10-15	Min 100 L/Ha	At foliar application stage
TURF	RATE L/Ha	WATER	COMMENTS
Greens and Tees	0.2-0.4 L/100m ²	6-10 L/100m ²	Early morning or afternoon
Fairways	20-40 L/Ha	600-1000 L/ha	Early morning or afternoon
VITICULTURE	RATE L/Ha	WATER	COMMENTS
Wine / Table Grapes	5-10	1:200	From bud burst to flowering
Fertigation	5-10	*	Apply 2-3 applications from bud burst to flowering
TREE CROPS	RATE L/Ha	WATER	COMMENTS
Citrus	10-15	1:200	Apply during flush
Avocado	7-15	1:200	Apply during flush to flowering
Olives	7-15	1:200	Apply prior to flowering
Fertigation	10-20	*	Apply 2-3 applications prior to flowering
HORTICULTURE	RATE L/Ha	WATER	COMMENTS
Beans	5-7	1:100	Apply at 4-6 true leaf stage
Brassica/Lettuce/Strawberry	5-7	1:100	Apply at 4-6 leaf stage
Carrots	5-7	1:100	Apply at seedling and vegetative stage
Potatoes/Tomatoes/Cucurbits	5-7	1:100	Apply at seedling and prior to flowering

*Application rate depends on application process

Bang P™

NPKS (13-19-0-0) +
Organic Catalysts

APPLICATION GUIDE

Apply through boom spray, aerial, band spray, fertigation, or overhead irrigation. DO NOT apply when conditions are extremely hot. Best applied in temperatures in the range of 5 - 25°C.

MIXING INSTRUCTIONS

1. Half fill tank to the required volume with water.
2. Add the required amount of Bang P™.
3. Add appropriate amount of wetting agent as per manufacturer's instructions.
4. Fill the remainder of tank with water and maintain agitation.

PACKAGE WEIGHTS

VOLUME	NET WEIGHT	GROSS WEIGHT
1000 Litres	1340 kg	1440 kg
200 Litres	268 kg	280 kg
20 Litres	26.8 kg	28.8 kg

COMPATIBILITY

Product may be compatible with herbicides and insecticides, please check with the manufacturer. Please be aware that physical compatibility does not guarantee product efficacy.

CONDITIONS OF SALE

These goods are manufactured and tested to specific standards and are sold in good faith, however, no guarantee is given as conditions and methods under which they are used are beyond the control of Advanced Nutrients Pty Ltd. Before using, user should determine the suitability of the product for his or her intended use and user assumes the risk and liability in connection therewith.

Office: 27 Westbourne Drive, Wights Mountain, QLD 4520

Phone: 1800 244 009

Email: info@advancednutrients.com.au

Web: www.advancednutrients.com.au