

Teodoro P. Nissen, M.D., Q.M.E. Fellowship Trained Board Certified Joseph M. Centeno, M.D. Fellowship Trained Board Certified

Phone: (707) 645-7210

Fax: (707) 645-7249

ARTHROSCOPIC ACROMIOPLASTY PROTOCOL

Includes Treatments for:

Arthroscopic Acromio-Clavicular Excision

Arthroscopic Partial Thickness Rotator Cuff Debridement

Post-op

0-1 Week:

Sling worn for comfort

- 1. Pendulum exercises 3x/day minimum
- 3. Elbow and wrist AROM 3x/day minimum
- 4. Cryocuff/ ice: days 1-2 as much as possible, then post activity

for pain

1-2 Weeks:

Avoid active elevation (anterior deltoid) until 6 weeks

- 1. PROM/AAROM (wand, pulley) within pain tolerance
- 2. Sub-maximal isometrics for shoulder musculature
- 3. Manual resistance for scapular motions
- 4. Postural awareness education

- 5. Gentle soft tissue mobilization
- 6. Modalities as indicated for pain or inflammation
- 7. Therapist supervised UBE for motion, avoiding substitution or Impingement

2-6 Weeks:

- 1. Progress PROM and AROM as tolerated
- Joint mobilization for scapula and glenohumeral mobility (posterior

capsule stretching)

3. Strengthening exercises for scapular stabilizers and rotator cuff.

Emphasis of strengthening on high reps and low weight, with

postural

awareness. May include PRE's, theratube/band, PNF and weight

equipment

Goal: Involved shoulder AROM grossly WNL without shoulder hiking

by 6 weeks

Phone: (707) 645-7210

Fax: (707) 645-7249