



BAY AREA ORTHOPEDIC

SURGERY & SPORTS MEDICINE

100 Hospital Drive. Suite 303 Vallejo CA 94589

Teodoro P. Nissen, M.D., Q.M.E.
Fellowship Trained
Board Certified

Joseph M. Centeno, M.D.
Fellowship Trained
Board Certified

DISCHARGE INSTRUCTIONS FOR TOTAL SHOULDER REPLACEMENT SURGERY

ANESTHESIA

- You may feel dizzy, light-headed, or sleepy for the first 12-24 hours after your operation. You cannot drive, operate any mechanical or electrical devices, or drink alcohol. Do not make any important decisions for 24 hours, and as long as you are taking prescription narcotic medication.
- If you have had general anesthesia, it is normal for you to feel generalized aching, sore muscles, and have a strange taste in your mouth or possibly a sore throat. This is normal and will dissipate in 24-48 hours.
- Resume your diet gradually as tolerated.

ACTIVITY

- Rest on the day of surgery.
- Start pendulums and passive forward flexion exercises. These are all you need to do prior to starting formal physical therapy.

ICE

- Ice (or cryocuff) the operative shoulder as tolerated for the first 2-3 days is necessary to reduce swelling and pain. Always place a dry towel as a barrier between the skin and ice. Take extra precaution if you have had a nerve block and are unable to feel the arm.

PAIN MEDICATION

- Discomfort from the surgical site is normal and expected. Medications prescribed during your pre-operative appointment are to be taken as directed. You should have medication available to you when you leave the surgery center or hospital.

CARE OF DRESSING/SHOWERING

- **DO NOT GET YOUR BANDAGE WET UNDER ANY CIRCUMSTANCES.** Wet bandages increase the risk for infection. Keep dressing clean and dry until follow up appointment.
- It is not uncommon to have small spots of bleeding show up on the outer aspect of the dressing. If this occurs, do not become alarmed. This type of bleeding typically stops overnight. If it seems to persist greater than 24 hours, please call my office. You may have small pieces of paper tape called steri-strips (or a Zipline) over the incision – **DO NOT REMOVE THE STERI-STRIPS OR ZIPLINE.**
- Report to our office any fever/chills, excessive swelling, redness to skin, discolored drainage, uncontrolled pain, persistent nausea/vomiting or any other concerns. Our office number is (707) 645-7210. Phones are attended to 24 hours/day.

POST-OP VISIT

- Please call the office the next working day after surgery to let us know how you are doing and to confirm your follow-up appointment.

Patient: _____ RN: _____