CALGARY SKIN CANCER CENTRE

#316 – 3320 17TH AVE SW CALGARY, AB T3E 0B4 P: (403) 700-0110 F: (403) 700-0271

Patient Post-Operative Aftercare Instructions – Biopsy/Wound care/Liquid Nitrogen

Instructions:
Site:
 Remove the initial pressure dressing after 72 hours (if applicable) Wash your hands thoroughly before removing the dressing You are now ok to shower as normal Cleanse the wound with a mild, antibacterial soap (ie. Dial Antibacterial Foaming Soap) Apply a thin layer of to the wound Cover the wound with a non-stick dressing and medical tape, or a band-aid type dressing Repeat this process once daily until healed.
DISCOMFORT If you have discomfort following surgery, take 1000mg Acetaminophen (two Extra Strength Tylenol®) every 6 hours (do not exceed 4000mg in 24 hours) OR a prescription pain medication if one has been prescribed to you. If this is not sufficient to control the pain, please call the office. AVOID Aspirin for the first week after surgery, and avoid medications that contain ibuprofen, naproxen and Celebrex (e.g. Aleve®, Excedrin®, Advil®, Motrin®) for 72 hours, as these products increase the risk of bleeding.
ACTIVITY Please avoid strenuous activity/exercise and heavy lifting with your full body for three (3) day to reduce your risk of bleeding and wound complications that will delay healing. This includes bending, lifting anything over 5 pounds, straining, going to the gym, golf, walking for exercise, yard wor housework, etc. or doing any type of activity that will increase your heart rate/blood pressure or cause stretching in the area the surgery was done. Other activity restrictions:

THINGS TO WATCH FOR:

BLEEDING

Your incision MAY ooze or bleed lightly within the first 48 hours after your procedure, especially if you take blood thinners. Your pressure dressing will help reduce this. However, if you have mild bleeding, it is not a cause for concern. **DO NOT REMOVE YOUR PRESSURE DRESSING**. You can control the bleeding with firm pressure for 30 minutes over top of the dressing. If you are experiencing excessive bleeding that is coming through your dressing and you are not able to get it to stop with firm pressure, please call the **24-hour emergency number at (403) 700-2840** and continue to apply pressure while proceeding to your nearest ER or Urgent Care Centre. If you have been hospitalized for <u>any reason</u> within the 10 days after your procedure, please notify our staff as soon as possible at (403) 700-0110, extension 4.

SWELLING

Swelling is common after surgery. Apply an ice pack around the surgical site and dressing – for 10 minutes every 1 hour, until you go to bed. You may continue this, if necessary, as long as the swelling persists. This will help with swelling, pain, and bruising. Sleep with your head elevated by using extra pillows or a recliner chair for 3 days.

INFECTION

Please notify us immediately if you see any signs of infection (excessive redness, swelling, yellow drainage, if the wound site becomes warm), or if the pain starts to get worse instead of better (after the first 48 hours). 403-700-0110, extension 4

FOLLOW-UP		
		aling process, please contact our Nursing Triage line are checked throughout the day between surgeries.
☐ Follow-up at Calgary	Skin Cancer Centre in	weeks
	pearance of your scar, recove	nely required. Should you have any concerns about ery, or any other concerns, please contact us to
☐ Contact your referrir	g physician,	to schedule your next skin check.
<u>RECOMMENDATI</u>	<u>ONS</u>	
Sun/Radiation Pro	tection:	
Non-sunscreen	options:	
• Sunscreen:		
Scar Care:		
• Product: Apply	Stratadermx/day onc	e
	in weeks. Massage for surgical site 5 times per da	5 minutes in a firm, circular motion all along ay for 3 months.
Skin Care:		
• AM:		
• PM•		