# CALGARY SKIN CANCER CENTRE

#316 – 3320 17<sup>TH</sup> AVE SW CALGARY, AB T3E 0B4 P: (403) 700-0110 F: (403) 700-0271

### **Patient Post-Operative Aftercare Instructions – SUTURES**

#### **INSTRUCTIONS:**

•	Your white pressure dressing(s) can be removed after HOURS.			
•	At this time, you can <b>shower as normal</b> .			
•	Cleanse the wound with a mild antibacterial soap (ie. Dial Antibacterial Foaming Soap)			
	☐ Apply a thin layer of ointment to the incision. Cover with a non-stick dressing and tape or band-aid.			
	☐ Apply ointment 2-3 x per day to keep incision moist.			
•	Repeat daily until sutures are removed.			

## **DISCOMFORT**

• Suture removal in \_\_\_\_ days.

If you have discomfort following surgery, take 1000mg Acetaminophen (two Extra Strength Tylenol®) every 6 hours (do not exceed 4000mg in 24 hours) OR a prescription pain medication if one has been prescribed to you. If this is not sufficient to control the pain, please call the office. AVOID Aspirin for the first week after surgery, and avoid medications that contain ibuprofen, naproxen and Celebrex (e.g. Aleve®, Excedrin®, Advil®, Motrin®) for 72 hours, as these products increase the risk of bleeding.

#### **ACTIVITY**

Please avoid strenuous activity/exercise and heavy lifting with your full body for the next three (3) days to reduce your risk of bleeding and wound complications that will delay healing. This includes bending, lifting anything over 5 pounds, straining, going to the gym, golf, walking for exercise, yard work, housework, etc. or doing any type of activity that will increase your heart rate/blood pressure or cause stretching in the area the surgery was done.

Other activit	y restrictions:	

#### **SWELLING**

Swelling is common after surgery. Apply an ice pack around the surgical site and dressing – for 10 minutes every 1 hour, until you go to bed for 5 days. You may continue this, if necessary, as long as the swelling persists. This will help with swelling, pain, and bruising.

#### **DRIVING**

DO NOT drive a motor vehicle for the remainder of the day after your surgery.

#### **THINGS TO WATCH FOR:**

#### **BLEEDING**

Your incision MAY ooze or bleed lightly within the first 48 hours after your procedure, especially if you take blood thinners. Your pressure dressing will help reduce this. However, if you have mild bleeding, it is not a cause for concern. **DO NOT REMOVE YOUR PRESSURE DRESSING**. You can control the bleeding with firm pressure for 30 minutes over top of the dressing. If you are experiencing excessive bleeding that is coming through your dressing and you are not able to get it to stop with firm pressure, please call the **24-hour emergency number at (403) 700-2840** and continue to apply pressure while proceeding to your nearest ER or Urgent Care Centre. If you have been hospitalized for <u>any reason</u> within the 10 days after your procedure, please notify our staff as soon as possible at (403) 700-0110, extension 4.

## **INFECTION**

Please notify us immediately if you see any signs of infection (excessive redness, swelling, yellow drainage, if the wound site becomes warm), or if the pain starts to get worse instead of better (after the first 48 hours). 403-700-0110, extension 4

FOLLOW-UP If you have any questions or concerns during the healing process at 403-700-0110, extension 4 *Please note messages are checken	
☐ Follow-up at Calgary Skin Cancer Centre in weeks	3
*Follow-up at Calgary Skin Cancer Centre is not routinely require your wound healing, appearance of your scar, recovery, or any copyour concerns (403-700-0110 ext. 4).	,
□ Contact your referring physician,	to schedule your next skin check.
RECOMMENDATIONS	
Sun/Radiation Protection:	
Non-sunscreen options:	
Sunscreen:	
<ul> <li>Scar Care:</li> <li>Product: Apply Stratadermx/day once</li> <li>Massage: Start in weeks. Massage for 5 minutes in a</li> </ul>	
the surgical site 5 times per day for 3 months.	mm, circular motion an along and around
Skin Care:	
• AM:	

• PM: \_\_\_\_\_