








How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
Do NOT use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months	_____	_____	_____	_____
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL 	2.5 mL* 	_____	_____
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL 	4 mL* 	_____	_____
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL 	5 mL* → 	1 tablet 	_____
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL 	 ← 7.5 mL*	1 ½ tablets 	_____
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL 	10 mL* → 	2 tablets 	1 tablet 
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years	_____	 ← 12.5 mL*	2 ½ tablets 	1 tablet 
72 to 95 pounds (about 33 to 43 kilograms)	11 years	_____	15 mL* → 	3 tablets 	1 ½ tablets 
96 pounds or more (44 kilograms or more)	12 years or older	_____	10 mL + 10 mL = 20 mL* 10 mL* →  and 10 mL* → 	4 tablets 	2 tablets 

***Note:** This dosage is for children's liquid ibuprofen products given by dropper. See the previous column for correct dose-by-weight for infant ibuprofen drops, which are more concentrated.



healthychildren.org

Powered by pediatricians. Trusted by parents.
 from the American Academy of Pediatrics