



124779/33

124779/33

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

FEEDING AND MANAGEMENT INSTRUCTIONS

When used as directed, **Mol-Mix Classic 28/2 WITH ALTOSID® FLY CONTROL** will provide sufficient (S)-methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season.

DIRECTIONS FOR USE

Each kilogram of Mol-Mix Classic 28/2 with Altosid® Fly Control contains 16.71 mg of S-Methoprene.

Feed beef cattle according to the table below to provide 7.5 to 15.1 mg (S-Methoprene) per 454 kilograms (1000 lb animal) of body weight per day. The average targeted rate for S-Methoprene is to provide 11.3 mg per 454 kg animal per day.

Animal Wt, kgs	Target Consumption, kgs/hd/d	Low Consumption, kgs/hd/d	High Consumption, kgs/hd/d
150	0.22	0.15	0.30
200	0.30	0.20	0.40
250	0.37	0.25	0.50
300	0.45	0.30	0.60
350	0.52	0.35	0.70
400	0.60	0.40	0.80
450	0.67	0.45	0.90
500	0.75	0.50	1.00
550	0.82	0.55	1.10
600	0.90	0.60	1.20

PREVENTS ADULT HORN FLY EMERGENCE FROM MANURE OF TREATED CATTLE.

Liquid feed supplement containing S-Methoprene* Insect Growth Regulator for continuous feeding to cattle through the Horn Fly season.

Active Ingredient

(S)-Methoprene (CAS #65733-16-6)..... 0.00167 %

Other Ingredients 99.99833 %

TOTAL 100.00000 %

*As the Insect Growth Regulator ALTOSID®

This feed contains added selenium at 4.54 mg/kg.

Guaranteed Analysis

Crude Protein	Minimum	28.0 %
(This includes a maximum of 25.0% equivalent crude protein from non-protein sources.)		
Crude Fat.....	Minimum	0.01 %
Crude Fiber.....	Maximum	0.1 %
Calcium (Ca).....	Actual	2.0 %
Phosphorus (P)	Actual	0.5 %
Salt (NaCl).....	Actual	5.6 %
Sodium (Na)	Actual	2.4 %
Magnesium (Mg)	Actual	0.1 %
Potassium (K)	Actual	1.6 %
Sulfur (S)	Actual	0.4 %
Cobalt (Co)	Actual	3.4 mg/kg
Fluorine (F)	Maximum	55.0 mg/kg
Copper (Cu).....	Actual	205 mg/kg
Iron (Fe).....	Actual	455 mg/kg
Manganese (Mn)	Actual	725 mg/kg
Zinc (Zn)	Actual	650 mg/kg
Iodine (I)	Actual	13.0 mg/kg
Vitamin A	Minimum	150,000 IU/kg
Vitamin D ₃	Minimum	40,000 IU/kg
Vitamin E	Minimum	100 IU/kg
TSI (Total Sugars as Invert)	Minimum	21.0 %
Dry Matter.....	Minimum	62.0 %
Moisture.....	Maximum	38.0 %

Ingredients

A list of the ingredients used in this feed may be obtained from the manufacturer or registrant.

CAUTION: USE ONLY AS DIRECTED.

KEEP OUT OF REACH OF CHILDREN

CAUTION: See side panel for additional Precautionary Statements and Directions

WARNING

THIS SUPPLEMENT SHOULD BE USED ONLY IN ACCORDANCE WITH THE DIRECTIONS ON THIS TAG. THIS PRODUCT CONTAINS SUPPLEMENTAL SELENIUM (SE) INTAKE OF SUPPLEMENTAL SELENIUM SHOULD NOT EXCEED 0.3 PPM ON A DRY BASIS.

Registered by:



Westway
FEED PRODUCTS

Where solutions flow.®

Westway Feed Products LLC
2001 Timberloch Place, Suite 400
The Woodlands, TX 77380

Net Weight Appears on Invoice

Density: 1.25 kg/L at 20°C.

The viscosity of this product will vary inversely with temperature.

Liquid Bulk Shipment

Registration Number: 621379

1. THIS SUPPLEMENT IS FOR BEEF CATTLE ONLY.
2. THIS SUPPLEMENT MAY BE FED FREE CHOICE IN A LICK WHEEL FEEDER, MIXED AS A PART OF A COMPLETE RATION, OR TOP DRESSED.
3. THIS PRODUCT IS A SUPPLEMENT! ADEQUATE ROUGHAGE AND OR GRAINS MUST BE AVAILABLE AT ALL TIMES. INTAKE OF THIS PRODUCT SHOULD NOT EXCEED THE MAXIMUM LEVEL LISTED IN THE TABLE ABOVE.
4. IN FREE CHOICE APPLICATIONS CONSUMPTION SHOULD BE CLOSELY MONITORED TO ENSURE THAT IT IS WITHIN THE GUIDELINES PROVIDED IN THE TABLE ABOVE BASED ON THE WEIGHT OF ANIMALS BEING FED. MANAGEMENT TECHNIQUES SHOULD BE EMPLOYED IF CATTLE ARE NOT CONSUMING THE PRODUCT AT THE DESIRED LEVEL.
5. FEEDERS DESIGNED FOR LIMITING THE INTAKE ARE RECOMMENDED. INTAKE CAN BE AFFECTED BY A NUMBER OF FACTORS SUCH AS FEEDER PLACEMENT RELATIVE TO WATER, SHADE OR LOAFING AREAS AND NUMBER OF ANIMALS PER FEEDER OR LICK WHEEL. THESE FACTORS CAN BE USED TO MANAGE LEVEL OF CONSUMPTION.
6. THIS PRODUCT CONTAINS SUPPLEMENTAL COPPER AND AT NORMAL CONSUMPTION LEVELS MAY CONTRIBUTE SOME 5 TO 12 PPM COPPER TO THE DIET. TOTAL DIET COPPER FROM ALL SOURCES SHOULD NEVER EXCEED 20 PPM. CONSULT YOUR DEALER FOR PRODUCTS SPECIFICALLY DESIGNED FOR SHEEP AND GOATS.
7. THIS PRODUCT MUST BE AGITATED WHILE BEING DISPENSED.
8. INTRODUCTION OF THIS PRODUCT AFTER HORN FLY INFESTATION IS ESTABLISHED WILL REQUIRE TREATMENT OF CATTLE WITH ADULTICIDES IF ELIMINATION OF THE ADULT FLY POPULATION IS DESIRED.

WARNING

THIS PRODUCT CONTAINS SUPPLEMENTAL COPPER. DO NOT FEED THIS PRODUCT TO SHEEP AND GOATS AS CONSUMPTION MAY BE FATAL. TOTAL DIETARY COPPER CONSUMPTION SHOULD NOT EXCEED 20 PPM. CONSULT YOUR DEALER FOR PRODUCTS SPECIFICALLY DESIGNED FOR SHEEP AND GOATS THAT CONTAIN NO ADDED COPPER.

OBSERVE THE FOLLOWING PRECAUTIONS:

- PROVIDE UNLIMITED DRINKING WATER
- PROVIDE ADEQUATE ROUGHAGE IN TERMS OF BOTH QUANTITY AND QUALITY
- PROVIDE ADEQUATE MINERAL SUPPLEMENT FREE-CHOICE
- DO NOT EXCEED THE DAILY MAXIMUM INTAKE SPECIFIED ABOVE
- DO NOT OFFER FREE-CHOICE TO OVERLY HUNGRY OR DEPRIVED ANIMALS AS OVER CONSUMPTION MAY RESULT

SUGGESTION: USE OF THIS PRODUCT MAY RAISE THE MOISTURE CONTENT OF FEEDS TO WHICH IT IS APPLIED. IF ANTICIPATED FINAL FEED MOISTURE IS TO EXCEED 12% AND STORAGE FOR PERIODS LONGER THAN 2 TO 3 DAYS IS EXPECTED, SPECIAL CONSIDERATION TO THE INCLUSION OF PRESERVATIVES IS NEEDED. YOUR WESTWAY REPRESENTATIVE WILL PROVIDE ASSISTANCE IF NEEDED.

If this product is introduced after an adult fly infestation is established and immediate elimination of the adult fly population is desired, it is suggested to treat cattle with an approved adulticide.

PRECAUTIONS: PREVENT ACCESS BY UNAUTHORIZED PERSONNEL. Wash thoroughly with soap and water after handling product and before eating, drinking or smoking. Wash contaminated clothing with soap and water before reuse. Do not wear contaminated shoes. Permits may be required for discharges containing this pesticide into lakes, ponds, streams or public waters. DO NOT discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans or other waters.

FIRST AID:

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS: Toxic to aquatic organisms.

DISPOSAL

Canadian manufacturers should dispose of unwanted active ingredients and containers in accordance with municipal and provincial regulations. For additional details and clean up of spills, contact the manufacturer or the provincial regulatory agency

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

SELLER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, AS TO DESCRIPTION, UTILITY, PRODUCTIVENESS, OR ANY OTHER MATTER, EXCEPT AS TO GUARANTEED ANALYSIS.