



Kansas State University®

Division of Biology

232 Ackert Hall
Manhattan, KS 66506 -4901
785-532-6615
Fax: 785-532-6653

BACTERIOLOGY LABORATORY REPORT ON GERM PRO SURFACE DISINFECTANT

June 29, 2006

1. Germ Pro Surface Disinfectant was tested to determine the Minimum Inhibitory Concentration against *Clostridium difficile* and its spores. In our opinion the disinfectant is effective to at least a 1:7 (Germ Pro:Bacteria) dilution under the conditions of our test.
2. The data in Table 1. support this conclusion.
3. Protocol:
 - a. *Clostridium difficile* was obtained from American Type Culture Collection and grown for 48 hr at 37°C in an anaerobic chamber in Thyoglycollate Broth Containing: Tryptone Casein 15 g, Yeast extract 5g, Dextrose 5.5 g, NaCl 2.5 g, L-Cysteine 0.5 g and Sodium thioglycollate 0.5 g. The culture was judged microscopically to contain at least 10% endospores.
 - b. Culture material was mixed and added to Germ Pro Surface Disinfectant for 24 hr @ 37°C.
 - c. Following incubation, 0.10 ml aliquots were plated on agar solidified Thyoglycollate Medium and incubated anaerobically at the same temperature for 24 hr.
 - d. Table 1 indicates that no colony forming units were obtained.

Table 1.

Minimum Inhibitory Test – Germ Pro ® Surface Disinfectant					
Serial Dilution*	1.5ml:0.5ml	1ml:1ml	0.75ml:1.25	0.5ml:1.5ml	0.25ml:1.75ml
Bacteria					
<i>Clostridium difficile</i> with ~10% spores (cfu/0.10ml)	0	0	0	0	0

* an overnight culture of *Clostridium difficile* containing >1.0 x 10⁸ cfu/ml was incubated with Germ Pro Surface Disinfectant for 24 hours at 37.5 °C. Then 0.10 ml aliquots were spin plated and incubated for 24 hr @ 37°C

Sincerely yours,

Dr. George L Marchin
Associate Professor of Microbiology

and Immunology

Division of Biology

Kansas State University

Manhattan, Kansas

Voice: 785 532 6635

FAX: 785 532 6653