

Lecture Note X – Summarized Learning Objectives for Topic 2

Descriptive Chemistry and Laboratory

Feb 2022

- *Doing chemistry requires both understanding ideas and remembering key information!*
- *Hands-on intensive lab experience is the most efficient way to study this topic.*

[Learning Objectives]

1. Flame Test, the Color of Ions, the Heat of Reactions/Dissolution, etc.

1.1 Flame Test

Introduction: A flame test is an analytical procedure used in chemistry to detect the presence of certain elements, primarily _____ (metal or nonmetal) ions, based on each element's characteristic _____ (absorption or emission) spectrum. What causes the bright colors in fireworks?

Operation: In high-school chemistry courses, wooden splints are commonly used, mostly because solutions can be dried onto them, and they are inexpensive. Bunsen burners are also commonly used as their flame is light.

[USNCO Example – N2016-P1-Q12]

[USNCO Example – N2020-P1-Q7]

- | | | | | | |
|--|---|---------------|-------------|---------------|------------|
| <p>12. Which is the safest method for performing a flame test?</p> <p>(A) Dissolve the metal salt in methanol, then squirt the solution into a lit Bunsen burner from at least 1 meter away.</p> <p>(B) Dissolve the metal salt in methanol, then pour the methanol into a crystallizing dish, igniting it with the flame from a Bunsen burner.</p> <p>(C) Soak a wooden splint in an aqueous solution of the metal salt, then burn the splint in the flame from a Bunsen burner.</p> <p>(D) Soak a wooden splint in an aqueous solution of the metal salt, then heat the splint on top of a ceramic hotplate.</p> | <p>7. Which ion gives a green flame test?</p> <table border="0"><tr><td>(A) Potassium</td><td>(B) Calcium</td></tr><tr><td>(C) Strontium</td><td>(D) Barium</td></tr></table> | (A) Potassium | (B) Calcium | (C) Strontium | (D) Barium |
| (A) Potassium | (B) Calcium | | | | |
| (C) Strontium | (D) Barium | | | | |

Typical flame colors: Colored flames of methanol solutions of different compounds, burning on cotton wool.
From left to right: LiCl, SrCl₂, CaCl₂, NaCl, BaCl₂, B(CH₃)₃, CuCl₂, CsCl and KCl.



- from Wikipedia

Fill the blanks with their flame colors.


Li	Na	K	Cu/B	Ca/Sr

1.2 The Color of Aqueous Ions – commonly used for AP Chemistry experiments

name	___manganate	chromate	dichromate	/
formula	$\text{MnO}_4^- (\text{aq})$			$\text{Cu}^{2+} (\text{aq}) / \text{Cu}(\text{NH}_3)_4^{2+} (\text{aq})$
color				___/deep blue
application	strong oxidant, redox titration, no extra indicator needed	LCP demo, strong oxidant, $\text{___} + 2\text{H}^+ \rightleftharpoons \text{___} + \text{H}_2\text{O}$ reduced to ___ (color: ___) in acidic		dehydration of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ formation of complex
name	/	/	/	/
formula	$\text{Co}^{2+} (\text{aq})$ or $\text{Co}(\text{H}_2\text{O})_6^{2+}$	$\text{CoCl}_4^{2-} (\text{aq})$	$\text{Fe}^{3+} (\text{aq})$	$\text{Ag}^+ / \text{Zn}^{2+} (\text{aq})$
color				<u>colorless</u>
application	LCP demo $\text{Co}(\text{H}_2\text{O})_6^{2+} + 4\text{Cl}^- \rightleftharpoons \text{CoCl}_4^{2-} + 6\text{H}_2\text{O}$ how does the color change if adding water?		K measurement for the formation of $\text{Fe}(\text{SCN})^{2+}$ ___	___ d subshell $\rightarrow$ NO d-d transition $\rightarrow$ no color


TRANSITION METAL ION COLOURS

Transition metals form coloured compounds and complexes. These colours can vary depending on the charge on the metal ion, and the number and type of groups of atoms (called ligands) attached to the metal ion. In aqueous solutions, the ions form complexes with the colours shown to the right.



Electrons are arranged around the nucleus of the metal atom in orbitals. Transition metals, unlike other metals, have partially filled d orbitals, which can hold up to 10 electrons. When ligands are present, some d orbitals become higher in energy than before, and some become lower. Electrons can then move between these higher and lower d orbitals by absorbing a photon of light. This absorption of light affects the perceived colour of the compound or complex. The wavelength of the light absorbed is affected by the size of the energy gap between the d orbitals, which is in turn affected by the type of ligand and the charge on the metal ion.

2014 COMPOUND INTEREST WWW.COMPOUNDINTEREST.COM



Extension:

- Color of Precipitates
- Crystal Field Theory (d orbital splitting)
- *Ligand-Metal Charge Transfer

- from compounds of interest

[USNCO Example – N2021-P1-Q9/N2020-P1-Q12]

9. Addition of 10 mL of distilled water to 0.1 g of which salt produces a yellow, slightly cloudy solution?

- (A) KMnO_4 (B) FeCl_3
(C) $\text{Co}(\text{NO}_3)_2$ (D) CuSO_4

12. Mixing which 0.1 M aqueous solutions results in formation of a colored precipitate?

- (A) BaCl_2 and CH_3COOH
(B) BaCl_2 and Na_2CO_3
(C) CuCl_2 and CH_3COOH
(D) CuCl_2 and Na_2CO_3

1.3 The Heat of Reactions/Dissolution – fill the blanks with *exothermic* or *endothermic*

- The dilution of concentrated H_2SO_4 is _____, the dissolution of solid NaOH in water is _____.
- The acid-base neutralizations are _____, most redox reactions are _____.
- Reaction of baking soda (formula: _____) with vinegar (formula: _____) is _____;
- The dissolution of solid sodium bicarbonate in water is _____.
- Dissolution of most ammonium salts, such as NH_4Cl , NH_4NO_3 is _____.
- *Reaction of solid barium hydroxide (formula: _____) with solid ammonium chloride (formula: _____) is _____. What is the reaction? _____

[USNCO Example – L2020-Q11]

[USNCO Example – N2015-P1-Q7]

11. Which reaction is not exothermic?

- (A) Dilution of concentrated hydrochloric acid in water.
- (B) Dilution of concentrated sulfuric acid in water.
- (C) Dissolution of solid sodium hydroxide in water.
- (D) Dissolution of solid sodium bicarbonate in water.

7. Each of the following substances dissolves exothermically in water EXCEPT

- (A) $\text{NaOH}(s)$.
- (B) $\text{NH}_4\text{NO}_3(s)$.
- (C) $\text{CuSO}_4(s)$.
- (D) $\text{H}_2\text{SO}_4(l)$.

2. Solubility Rules

- Pretty much all nitrates, acetates, Na^+ , K^+ , NH_4^+ are soluble, a lot of precipitates are white unless _____.
- Most chlorides are soluble except _____ and _____ (dissolve in hot water).
- Most sulfates are soluble except Pb^{2+} and several group _____ cations, such as Ca^{2+} , Sr^{2+} , and Ba^{2+} .
- Most hydroxides, carbonates, oxalates, sulfides are insoluble except those listed in Rule #1.
- *Fluorides are different from other halides, AgF is _____ while other AgX ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) are insoluble, MF_2 ($\text{M} = \text{Ca}, \text{Sr}, \text{Ba}$) are _____ while other MX_2 ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) are soluble.
- Precipitates made by conjugate base of weak acids are more soluble in _____, such as hydroxides, carbonates, oxalates, and sulfides. Why? How about the precipitates of halides (excepts fluorides)?
- Some precipitates dissolves when forming complex ions: $\text{AgCl}(s)$ dissolved in concentrated _____ (formula of complex: _____), $\text{Al}(\text{OH})_3(s)$ dissolved in concentrated _____ (_____).

[USNCO Example – N2021-P1-Q8]

[USNCO Example – N2016-P1-Q8]

8. A solution that may contain either 0.1 M $\text{Ag}^+(aq)$, 0.1 M $\text{Pb}^{2+}(aq)$, or both, is treated with 1 M aqueous HCl . A white precipitate forms which does not appear to dissolve in hot water. Which conclusion about the cations present may be drawn?

- (A) Only Ag^+ is present.
- (B) Only Pb^{2+} is present.
- (C) Ag^+ is present, and Pb^{2+} may be present.
- (D) Pb^{2+} is present, and Ag^+ may be present.

8. A student has 10 mL of a solution that might contain any or all of the following cations at 0.01 M concentrations: Mn^{2+} , Ba^{2+} , Ag^+ , and Cu^{2+} . Addition of 10 mL of 1 M HCl causes a precipitate to form. After the precipitate is filtered off, 1 M H_2SO_4 is added to the filtrate and another precipitate forms. What is the second precipitate?

- (A) MnSO_4
- (B) BaSO_4
- (C) Ag_2SO_4
- (D) A mixture of BaSO_4 and Ag_2SO_4

3. *Properties of Representative Metals/Nonmetals

3.1 General Properties of Metals

- Metals conduct heat and electricity, tend to _____ electrons (lose or gain).
- _____ metals and most _____ metals are highly reactive, which *violently* react with water to produce $M^I(OH)$ or $M^{II}(OH)_2$ and _____.
 $Na + H_2O \rightarrow Na^+ + OH^- + \underline{\hspace{1cm}}$
- Metals with a standard reduction potential _____ 0 V (> or <) (**pre-H** metals) react with acid to produce the corresponding salts and _____, such as Mg, Al, and several 4th period transition metals: _____ and _____.
 $Fe + H^+ \rightarrow \underline{\hspace{1cm}} + H_2(g)$
- Some metals such as Cu, Ag, Pt, Au (metallic money) are unreactive, and _____ is commonly used as inert electrodes.
- Most metal oxides are _____ (acidic or basic), reacting with acids to form metal cations and _____.
 $Fe_2O_3 + H^+ \rightarrow Fe^{3+} + \underline{\hspace{1cm}}$
- More reactive metals can displace the less reactive metals from their cation solutions:



[USNCO Example – N2017-P1-Q8/L2020-Q9]


8. An element is a solid at room temperature but soft enough to be cut with an ordinary knife. When placed in water, the element reacts violently. What element is it?

(A) Na (B) Mg (C) Cu (D) Hg

MOST REACTIVE	
potassium	K
sodium	Na
calcium	Ca
magnesium	Mg
aluminium	Al
carbon	C
zinc	Zn
iron	Fe
tin	Sn
lead	Pb
hydrogen	H
copper	Cu
silver	Ag
gold	Au
platinum	Pt
LEAST REACTIVE	

3.2 General Properties of Nonmetals

- Electronegative nonmetals tend to _____ electrons, typical examples are halogens, dioxygen.
- Most nonmetals react with hydrogen gas to produce their covalent hydrides.
- Typical nonmetal oxides are _____, reacting with _____ to produce the corresponding acids.
- The more reactive halogens (X_2) can displace the less reactive halogens from their halide solutions:



[USNCO Example - N2015-P1-Q11/L2021-Q11]

11. Elemental silicon is oxidized by O_2 to give a compound which dissolves in molten Na_2CO_3 . When this solution is treated with aqueous hydrochloric acid, a precipitate forms. What is the precipitate?

(A) SiH_4 (B) $SiCO_3$ (C) SiO_2 (D) $SiCl_4$

9. A strip of metallic zinc is placed in a beaker containing dilute aqueous copper(II) nitrate. Which statement correctly describes what takes place?
- (A) No reaction takes place.
- (B) The mass of the metal strip decreases as the zinc is oxidized.
- (C) A white precipitate of $CuNO_3$ is formed.
- (D) Bubbles of $NO(g)$ form as the nitrate ion is reduced.

11. Chlorine gas is bubbled into a colorless aqueous solution of sodium iodide. Which is the best description of what takes place?
- (A) A precipitate of white $NaCl$ forms.
- (B) A precipitate of metallic Na forms.
- (C) The solution turns pale green as the chlorine dissolves.
- (D) The solution turns yellow-brown as iodide reacts with the chlorine.

3.3 Amphoteric Metals

Several metals close to the metal/nonmetal boundary are amphoteric, meaning that they can react with both _____ and _____. Their oxides and hydroxides are also _____. The most typical example is _____.

Reaction example: $\text{Al} + \text{OH}^-(\text{aq}) + \text{_____} \rightarrow \text{Al}(\text{OH})_4^- + \text{_____}$

[USNCO Example - N2015-P1-Q12]

[USNCO Example - N2016-P1-Q9]

- | | |
|---|---|
| <p>12. A metal dissolves in 3.0 M NaOH solution with evolution of gas to form a clear, colorless solution. Upon neutralization, the solution forms a gelatinous precipitate. What is the metal?</p> <p>(A) Al (B) Ag (C) Cu (D) Mg</p> | <p>9. When 6 M sodium hydroxide is added to an unknown white solid, the solid dissolves. What is a possible identity for this solid?</p> <p>(A) $\text{Mg}(\text{OH})_2$ (B) $\text{Al}_2(\text{SO}_4)_3$
(C) BaCO_3 (D) AgBr</p> |
|---|---|

How about if *excess* strong acid is added in the second step?

4. Strong/Weak Electrolytes, Acid-Base Property of Salt Solutions

4.1 Commonly Used Acids and Bases

- List the THREE most commonly used strong acids in a HS chemistry lab: _____, _____, _____ (first step strong, second step weak), a few more strong acids are $\text{HBr}(\text{aq})$, $\text{HI}(\text{aq})$, $\text{HClO}_4(\text{aq})$.
- $\text{HF}(\text{aq})$ is a _____ (strong or weak) acid, the only exception in hydrohalic acids.
- Most carboxylic acids (RCOOH) are _____ acids, such as acetic acid _____ (formula).
- List the THREE most used strong bases in a HS chemistry lab: _____, _____, $\text{Ba}(\text{OH})_2$.
- NH_4OH (aqueous solution of ammonia) is a typical _____ (strong or weak) base.
- Amines are weak _____, which are organic derivatives of ammonia, NH_3 .

4.2 Acid-Base Properties of Salt Solutions

- Salts made by cations of **strong** bases and conjugate bases of **strong** acids are _____ (acidic, neutral, or basic), such as $\text{NaCl}(\text{aq})$.
 - Salts made by cations of **strong** bases and conjugate bases of **weak** acids are _____ (acidic, neutral, basic), such as $\text{NaF}(\text{aq})$ and $\text{CH}_3\text{COONa}(\text{aq})$.
 - Salts made by cations of **weak** bases and conjugate bases of **strong** acids are _____ (acidic, neutral, basic), such as $\text{NH}_4\text{Cl}(\text{aq})$.
 - Salts made by highly charged cations and conjugate bases of **strong** acids are _____ (acidic, neutral, basic), such as $\text{FeCl}_3(\text{aq})$.
 - the weaker the acids are, the _____ (more or less) basic the conjugate bases are (conjugate see-saw).
- *Extension: 0.1 M $\text{Na}_2\text{CO}_3(\text{aq})$ is _____ (more or less) basic than 0.1 M $\text{NaHCO}_3(\text{aq})$.

How to explain the above statements? Taking $\text{NH}_4\text{Cl}(\text{aq})$ and $\text{NaF}(\text{aq})$ as examples.

[USNCO Example – L2018-Q9]

[USNCO Example - N2017-P1-Q10]

9. A 0.1 M solution of which salt is the most basic? 10. A 0.1 M solution of which salt is the most acidic?
- (A) NaNO_3 (B) NaClO_4 (A) $\text{Al}(\text{NO}_3)_3$ (B) MgBr_2
 (C) NaHSO_4 (D) NaHCO_3 (C) NaHCO_3 (D) NaHCO_2

5. Typical Gas Evolution and Redox Reactions

5.1 Typical Gas Evolution Reactions

gas evolved	reaction	net ionic equation
$\text{H}_2(\text{g})$	Pre-H metals reacting with _____	
$\text{CO}_2(\text{g})$	carbonates or bicarbonates reacting with _____	
$\text{SO}_2(\text{g})$	sulfites or bisulfites reacting with _____	
$\text{NH}_3(\text{aq})$	ammonium reacting with strong _____	
$\text{NO}_x(\text{g})$	metals including post-H metals reacting with _____ acids	why no $\text{H}_2(\text{g})$ produced?

[USNCO Example - N2020-P1-Q10]

[USNCO Example - L2017-Q10]

10. Which substance produces a toxic and explosive gas when added to strong acids? 10. Addition of small amounts of which solids to 4 M HCl will result in gas evolution?
- (A) NaN_3 (B) Na_2CO_3 I. Zn II. Na_2SO_3
 (C) NaClO_4 (D) Na_2SO_3 (A) I only (B) II only
 (C) Both I and II (D) Neither I nor II

5.2 Typical Redox Reactions

redox reaction	characters	net ionic equation
metals reacting with water/acid/base	H^+ or H_2O is reduced into _____	
decomposition of $\text{H}_2\text{O}_2(\text{aq})$	_____ thermic, catalyzed by a variety of catalysts, such as $\text{MnO}_2(\text{s})$, Br^- , I^- , Fe^{3+} , etc.	
standardization of MnO_4^- (aq) by $\text{C}_2\text{O}_4^{2-}$ or $\text{Fe}^{2+}(\text{aq})$	MnO_4^- is _____ (color), used as oxidant and indicator (turns <u>pink</u> when last drop of MnO_4^- is added), solution is acidified	
titration of $\text{H}_2\text{O}_2(\text{aq})$ using standardized MnO_4^- (aq)	H_2O_2 is oxidized into _____, solution is acidified, MnO_4^- is reduced to _____ (color _____)	
titration of $\text{I}_2(\text{aq})$ using standardized $\text{S}_2\text{O}_3^{2-}(\text{aq})$	starch as indicator, forms _____ (color) complex with I_2/I_3^- , $\text{S}_2\text{O}_3^{2-}$ is oxidized into _____	


[USNCO Example - N2021-P1-Q10]

10. The amount of ascorbic acid in a vitamin C tablet is determined by titration with a solution of iodine in aqueous potassium iodide. A small amount of starch is added to the vitamin C solution before the titration. What is the function of the starch?
- (A) It forms an intensely colored complex with triiodide ion.
- (B) It increases the viscosity of the analyte solution.
- (C) It catalyzes the dissociation of triiodide into iodine and iodide.
- (D) It binds to the inert ingredients of the vitamin C tablet.

[USNCO Example - N2015-P1-Q9]

9. A student standardizes a solution of $\text{Na}_2\text{S}_2\text{O}_3$ by titrating it against a solution containing a known mass of NaIO_3 that has been dissolved in an excess of a freshly prepared solution of KI in dilute HCl. Which of the following errors will lead to a value of the molarity of the thiosulfate solution that is higher than the true value?
- (A) The student overshoots the endpoint of the titration.
- (B) The NaIO_3 is contaminated with NaCl.
- (C) The KI/HCl solution is allowed to stand overnight before it is used in the titration.
- (D) The sample of sodium thiosulfate pentahydrate used to make the $\text{Na}_2\text{S}_2\text{O}_3$ solution had partially dehydrated on standing.

6. Volumetric Glassware

volumetric glassware					
name					
uncertainty	$\pm 0.0X \text{ mL}$	$\pm 0.0X \text{ mL}$	$\pm 0.0X \text{ mL}$	$\pm 0.X \text{ mL}$	$\pm 0.0X \text{ mL}$
other characters	to contain (TC)	bulb needed, *finger on top	read the volume difference	easy to use	bulb needed, *finger on top
application	preparing a solution with a certain molarity	transfer a <i>fix</i> volume of solution	titrations	transfer a certain volume of solution with a <i>lower</i> precision	transfer a <i>certain</i> volume of solution

Beakers, Erlenmeyer flasks, or disposable plastic/glass pipets can't be used for serious volume measurement as their marks are not precise and for reference only.

[USNCO Example - N2021-PI-Q7]

7. Which glassware would be most appropriate for measuring 10.00 mL of distilled water?

(A) A 10-mL test tube
(B) A 25-mL beaker
(C) A 50-mL buret
(D) A 100-mL graduated cylinder

[USNCO Example - L2018-Q11]

11. Which would be most suitable for measuring 2.7 mL of ethanol for addition to a reaction with acidified dichromate?

(A) 10-mL graduated cylinder
(B) 10-mL volumetric flask
(C) 10-mL volumetric pipet
(D) 10-mL beaker

[USNCO Example - L2019-Q12]

12. Which is the best way to dispense liquids using a volumetric pipet?

(A) The pipet is immersed in the liquid to be dispensed, then lifted with a gloved finger on the top and the liquid allowed to drain to the mark. The remaining contents are then allowed to drain into the desired container.

(B) The pipet is immersed in the liquid to be dispensed, then lifted with a gloved finger on the top and the liquid allowed to drain to the mark. The remaining contents are then allowed to drain into the desired container, with a pipet bulb used to gently blow out any residual droplets.

(C) The tip of the pipet is submerged below the surface of the liquid and suction is applied using a pipet bulb until the liquid rises above the level of the mark. A gloved finger is then applied to the top of the pipet and the liquid allowed to drain to the mark. The remaining contents are then allowed to drain into the desired container.

(D) The tip of the pipet is submerged below the surface of the liquid and suction is applied by mouth until the liquid rises to the level of the mark. The contents are then allowed to drain into the desired container.

[USNCO Example - L2020-Q8]

8. A student is using a buret for a titration. What initial buret reading should be recorded?



(A) 6.6 mL
(B) 7.4 mL
(C) 6.63 mL
(D) 7.37 mL

7. UV-Vis Spectroscopy and Beer's Law

7.1 Principle

UV/Vis spectroscopy is routinely used in AP and analytical chemistry for the *quantitative* determination of different analytes or sample with colors, based on the *Beer's Law*

$$Abs = \epsilon lc$$


where ϵ is the _____, l is _____, c is the molarity of the solution.

7.2 Procedures

- prepare the standard solution series
- measure the spectrum (Abs vs λ), “blank” is needed
- pick up a wavelength ($\lambda_{\max}$ is commonly used unless $Abs@_{\lambda_{\max}}$ is too large or **interference** existed)
- measure the $Abs@_{\lambda}$ and plot the calibration curve
- measure the Abs of the sample solution at the picked λ , plot the Abs in the calibration curve to get the c .

[USNCO Example - N2015-P1-Q6]

6. Nitrophenol is a colorless weak monoprotic acid ($pK_a = 7.2$) whose conjugate base is bright yellow. To 2.00 mL of a solution of 0.0100 M nitrophenol is added 1.00 M NaOH in 0.001 mL portions, and the absorbance of the solution at 485 nm is monitored. What does the graph of A_{485} as a function of added volume of NaOH look like?



[USNCO Example - N2021-P1-Q12]

12. A student performs an experiment to determine the concentration of a colored salt solution by measuring the absorbance of the solution at the wavelength of maximum absorbance of the salt ($\lambda_{\max}$) and using Beer's Law to calculate the concentration. Which of the following could cause the measured concentration to be higher than the actual concentration?

- (A) The cuvette is not rinsed with the salt solution after being washed.
- (B) The cuvette is not wiped off before it is inserted into the spectrophotometer.
- (C) Less than the recommended volume of salt solution is added to the cuvette.
- (D) The spectrometer is set to a wavelength different from $\lambda_{\max}$.


7.3 Color Wheel

The color of the solution is the _____ color of the light absorbed.

[USNCO Example - N2020-P1-Q8]

8. The equilibrium constant for the formation of CoCl_4^{2-} , a species that is blue in solution, is to be measured using a colorimeter. The colorimeter has wavelength settings of 470 nm (“blue”), 565 nm (“green”), and 635 nm (“red”) to use in the experiment. What is the best setting to use?

- (A) 470 nm
- (B) 565 nm
- (C) 635 nm
- (D) All settings would be equally suitable for the measurement.



8. Titrations and Error Analysis

8.1 Error Analysis

[USNCO Example - N2020-P1-Q9(calorimetry)]

9. A student is using a coffee-cup calorimeter to determine the enthalpy change of the endothermic reaction of two aqueous solutions. After both solutions are added to the cup, the student neglects to put the lid on the cup. This would cause the magnitude of the calculated ΔH° value to be:
- (A) too small, since some heat will escape out of the cup.
 - (B) too large, since some heat will escape out of the cup.
 - (C) too small, since the solution will absorb heat from the room.
 - (D) too large, since the solution will absorb heat from the room.

[USNCO Example – L2019-Q11(gravimetric analysis)]

11. The concentration of sulfate ion in a solution is measured by precipitating the sulfate as BaSO_4 , filtering the precipitate on ashless filter paper, and heating the filter paper and precipitate in a tared crucible with a Bunsen burner. Which error will result in a sulfate concentration that is higher than the actual concentration?
- (A) The empty crucible contains a few drops of water when it is tared.
 - (B) A glass fiber filter is used instead of ashless filter paper.
 - (C) Some fine precipitate is not captured by the filter.
 - (D) Some of the sulfate-containing solution spills before the BaCl_2 solution is added.

[USNCO Example - N2015-P1-Q9(iodometry)]

9. A student standardizes a solution of $\text{Na}_2\text{S}_2\text{O}_3$ by titrating it against a solution containing a known mass of NaIO_3 that has been dissolved in an excess of a freshly prepared solution of KI in dilute HCl. Which of the following errors will lead to a value of the molarity of the thiosulfate solution that is higher than the true value?
- (A) The student overshoots the endpoint of the titration.
 - (B) The NaIO_3 is contaminated with NaCl.
 - (C) The KI/HCl solution is allowed to stand overnight before it is used in the titration.
 - (D) The sample of sodium thiosulfate pentahydrate used to make the $\text{Na}_2\text{S}_2\text{O}_3$ solution had partially dehydrated on standing.

[USNCO Example - N2019-P1-Q12(acid-base titration)]

12. The ammonia concentration of a solution is determined by titrating with aqueous HCl (previously standardized against Na_2CO_3) using a pH meter. Which of the following errors will lead to a measured concentration of NH_3 that is higher than the actual concentration?
- (A) Some of the Na_2CO_3 used in the standardization is spilled before being transferred to the titration flask.
 - (B) The glass stirring rod used to stir the ammonia solution is wiped with a paper towel after each aliquot of HCl is added.
 - (C) The ammonia solution is allowed to stand in an open beaker for an hour before being titrated.
 - (D) The pH meter has been miscalibrated so that all readings are 2.00 pH units higher than the actual pH.

8.2 Important Concepts in Titrations

- titrant (typical put in the burette)
 - analyte (typically put in the titration flask)
 - indicator (phenolphthalein, pH range 8~10, change from ____ to ____)
 - **standardization** (KMnO_4 , I_2 , HCl, NaOH, etc.)
 - *primary standard (KIO_3 , Na_2CO_3 , potassium hydrogen phthalate (KHP), $\text{Na}_2\text{C}_2\text{O}_4$)
- What is the application of each?
- equivalence point vs end point
 - half equivalence point

9. Some Extra Challenging Questions

[USNCO Example – N2000-P1-Q1-6]

1. Which of these ions is expected to be colored in aqueous solution?

I Fe^{3+} II Ni^{2+} III Al^{3+}

- (A) I only (B) III only
(C) I and II only (D) I, II, and III

2. Which substance is stored in contact with water to prevent it from reacting with air?

- (A) bromine (B) lithium
(C) mercury (D) phosphorus


3. A solution of concentrated aqueous ammonia is added dropwise to 1 mL of a dilute aqueous solution of copper(II) nitrate until a total of 1 mL of the ammonia solution has been added. What observations can be made during this process?

- (A) The colorless copper(II) nitrate solution turns blue and yields a dark blue precipitate.
(B) The colorless copper(II) nitrate solution yields a white precipitate which turns dark blue upon standing.
(C) The light blue copper(II) nitrate solution yields a precipitate which redissolves to form a dark blue solution.
(D) The light blue copper(II) nitrate solution turns dark blue and yields a dark blue precipitate.

4. What gas is produced when dilute HNO_3 is added to silver metal?

- (A) NO (B) H_2 (C) NH_3 (D) N_2


5. A substance is analyzed by paper chromatography, giving the chromatogram shown.



What is the R_f value of the substance represented by the spot at 8.0 cm?

- (A) 0.80 (B) 0.75 (C) 0.67 (D) 0.60

6. The molarity of a Cu^{2+} solution is to be determined from its absorbance, measured under the same conditions as those used to prepare this calibration curve. What will be the percent uncertainty in the concentration of a 0.050 M solution if the uncertainty in the absorbance reading is ± 0.01 absorbance units?



- (A) 5% (B) 10% (C) 15% (D) 20%

[USNCO Example – N2001-P1-Q1-5&Q7/N2010-Q6]

1. Which of these compounds is amphoteric?

I. $\text{Al}(\text{OH})_3$ II. $\text{Ba}(\text{OH})_2$ III. $\text{Zn}(\text{OH})_2$

- (A) I only (B) II only
(C) I and III only (D) II and III only

2. Calcium hydride reacts with excess water to form

- (A) CaO and H_2 (B) $\text{Ca}(\text{OH})_2$ and O_2
(C) $\text{Ca}(\text{OH})_2$ only (D) $\text{Ca}(\text{OH})_2$ and H_2

3. What is the most likely boiling point of an equimolar mixture of hexane, C_6H_{14} , and heptane, C_7H_{16} ?

Boiling Point	
C_6H_{14}	69 °C
C_7H_{16}	98 °C

- (A) below 69 °C (B) between 69 and 98 °C
(C) 69 °C (D) 98 °C


4. Which element melts at the highest temperature?

- (A) aluminum (B) silicon
(C) phosphorus (D) sulfur

5. Which substance participates readily in both acid-base and oxidation-reduction reactions?

- (A) Na_2CO_3 (B) KOH
(C) KMnO_4 (D) $\text{H}_2\text{C}_2\text{O}_4$

7. What is the purpose of this apparatus?



- (A) distilling (B) filtering
(C) refluxing (D) titrating

6. Four elements were tested in the laboratory and gave the results in the table below. Which element is a metalloid?

Element	Appearance	Conductivity	Behavior with HCl
A	Slight luster	High	Bubbles slowly
B	Shiny	Low	No reaction
C	Dull	None	No reaction
D	Shiny	High	Bubbles rapidly

- (A) Element A (B) Element B
(C) Element C (D) Element D

[USNCO Example – N2011-P1-Q1-2&Q7]

- Which solid does not react with a small amount of 3 M HNO_3 ?
 (A) calcium carbonate (B) manganese(II) sulfide
 (C) potassium sulfite (D) silver chloride
- A metal dissolves in dilute HCl in the absence of air to form a pale green solution that turns yellow upon exposure to air. The metal could be
 (A) iron. (B) manganese.
 (C) nickel. (D) vanadium.
- An aqueous solution contains the ions Ag^+ , Ba^{2+} , and Ni^{2+} . Dilute aqueous solutions of NaCl , Na_2S , and Na_2SO_4 are available. In what order should these solutions be added if the goal is to precipitate each of the three cations separately?
 (A) Na_2S , Na_2SO_4 , NaCl (B) Na_2S , NaCl , Na_2SO_4
 (C) Na_2SO_4 , Na_2S , NaCl (D) NaCl , Na_2SO_4 , Na_2S

[USNCO Example – N2013-P1-Q8-9]

- Which substance is used in self-contained breathing equipment because it absorbs exhaled CO_2 and H_2O and releases O_2 gas?
 (A) KO_2 (B) Na_2O_2 (C) NaOH (D) Li_2O
- A sample of a white solid is known to be NaHCO_3 , AgNO_3 , Na_2S , or CaBr_2 . Which 0.1 M aqueous solution can be used to confirm the identity of the solid?
 (A) $\text{NH}_3(\text{aq})$ (B) $\text{HCl}(\text{aq})$
 (C) $\text{NaOH}(\text{aq})$ (D) $\text{KCl}(\text{aq})$

[USNCO Example – N2014-P1-Q11]

- All of the following can be used as primary standards in acid-base titrations EXCEPT
 (A) oxalic acid.
 (B) potassium hydrogen phthalate.
 (C) sodium carbonate.
 (D) sodium hydroxide.

[USNCO Example – N2018-P1-Q7&10]

- Which salt is diamagnetic?
 (A) $\text{K}_2[\text{NO}(\text{SO}_3)_2]$ (B) $\text{K}_4[\text{Fe}(\text{CN})_6]$
 (C) $\text{Ce}_2(\text{SO}_4)_3$ (D) $\text{Hg}[\text{Co}(\text{SCN})_4]$
- A copper-nickel alloy is analyzed by dissolving it in 8 M nitric acid, diluting the solution with water, and then adding 1 mL of this diluted solution to an excess of aqueous potassium iodide. What are the principal forms of copper and nickel in this final mixture?
 (A) $\text{Cu}^{2+}(\text{aq})$, $\text{Ni}^{3+}(\text{aq})$ (B) $\text{CuI}_2(\text{s})$, $\text{NiI}_2(\text{s})$
 (C) $\text{CuI}(\text{s})$, $\text{Ni}^{2+}(\text{aq})$ (D) $\text{Cu}^+(\text{aq})$, $\text{Ni}(\text{s})$

[USNCO Example – N2019-P1-Q10]

- A pale yellow solid is insoluble in water or concentrated ammonia solution, but dissolves in concentrated $\text{Na}_2\text{S}_2\text{O}_3$ solution. What is it?
 (A) AgF (B) AgCl
 (C) AgI (D) $\text{Ag}(\text{CH}_3\text{COO})$

[USNCO Example – N2020-P1-Q10-11]

- Which substance produces a toxic and explosive gas when added to strong acids?
 (A) NaN_3 (B) Na_2CO_3
 (C) NaClO_4 (D) Na_2SO_3
- When aqueous ethanol is treated with an acidified solution of potassium dichromate, what is observed?
 I. Color change
 II. Gas evolution
 (A) I only (B) II only
 (C) Both I and II (D) Neither I nor II

[USNCO Example – N2021-P1-Q11]

- When blue solid $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ is heated, it turns white. Which is the best explanation for this observation?
 (A) Heating causes water molecules bonded to copper to be replaced by sulfate, lowering the energy of the light absorbed.
 (B) Heating causes an expansion of the lattice, leading to a decrease in the coordination number of copper and raising the energy of the light absorbed.
 (C) Heating causes reduction of copper(II) to copper(I).
 (D) Heating causes reduction of sulfate to sulfite.

[USNCO Example – L2012-Q7]

- A dilute solution of which acid is most likely to produce a reduction product other than H_2 when it reacts with a metal?
 (A) HF (B) HCl (C) HNO_3 (D) H_2SO_4

[USNCO Example – L2018-Q7]

- A divalent metal ion dissolved in dilute hydrochloric acid forms a precipitate when H_2S is bubbled through the solution. Which ion is it?
 (A) Ca^{2+} (B) Mn^{2+} (C) Zn^{2+} (D) Cd^{2+}