

MEDICATION LIST

PRODUCT	USE	DOSAGE	METHOD
ANTIBIOTICS AND OTHER MEDICATION			
Maxitrel LA	Treatment of heartwater, tick-borne gallsickness (anaplasmosis), pneumonia, footrot, joint-ill and naval ill.	1 mℓ per 10kg	Subcutaneous, intramuscular or intravenous injection
Engemycin 10%	For the treatment of heartwater, bacterial pneumonia, footrot, mastitis, navel/joint-ill and bacterial wound infections.	1 mℓ per 10 kg	Intramuscular or intravenous injection; In sheep and goats more than 50 kg body mass, the dose should be divided and administered at 2 injection sites.
Doxymycin QA	For the treatment of heartwater, anaplasmosis (tick-borne gallsickness), pneumonia, footrot, navel-ill and joint-ill.	1 mℓ per 10 kg	Intramuscular or intravenous injection
Duplocilin	For the treatment of infections caused by bacteria sensitive to penicillin, where long-acting therapy is required.	1 mℓ / 20 kg. Injection may be repeated at 72 hour intervals depending on clinical judgment given at a different injection site.	Intramuscular injection
NuFlor	Pasteurella	1mℓ / 15kg	Intramuscular injection
Finadyne	Inflammation and Fever	1 mℓ/ 25kg	Slow intravenous/ intramuscular injection
Dexa 0,2 Phenix	Kortikosteroïde	1mℓ / 15kg	Intramuscular injection
Disulfox LA	Coccidiosis and bacterial scours in lambs	Initial Dose - 1 to 2 mℓ/10 kg Maintenance Dose - 0.5 to 1 mℓ/10 kg	Subcutaneous or intravenous injection
INTERNAL PARASITES			
Dectomax	Used to treat and control internal and external parasites	1 – 1,5 mℓ / 50 kg	Subcutaneous, intramuscular injection
Lintex - L	Worm remedy against tapeworms in sheep, goats, calves and immature conical fluke in sheep and goats. Can be used for animals of all ages and pregnant animals.	1 mℓ / 4 kg	Oral drench
Prodose Orange	Nasal bot, liver fluke, roundworm and milk tapeworm remedy for sheep and goats with residual protection against re-infestation of wireworm and hookworm.	2 mℓ / 10 kg	Oral drench
Prodose Red	Roundworm remedy	3 mℓ / 10 kg	Oral drench
Startect	A broad spectrum oral combination drench effective against adult and immature (L4) stages of sensitive strains of internal parasites, including those resistant to macrocyclic lactones, salicylanilides, levamisole, organophosphors and benzimidazole drenches. Also controls nasal bot larvae and itch mites.	1 mℓ / 5 kg	Oral drench
First Drench	Oral anthelmintic for the control of roundworms, tapeworms and nasal worm	Lambs and kids older than 6 weeks of age a dose of 2,5 mℓ / 10 kg	Oral drench
SPRAYS, OINTMENTS & POWDERS			
Supa Spray Plus	Contains Sulfadiazine with proven anti-bacterial and wound healing properties. Forms protective aluminium layer over wounds		
Stokholm	Hooves		
Supona Aerosol	Controls ticks by spot treatment. Kills blowfly maggots in wounds. Use in cuts, abrasions, after dehorning, branding, shearing, tail docking and castrations to prevent infections and maggot infestations		
Oberdne	Wound, Eye and Footrot Spray		
Speen cream	A powerful antiseptic & softening agent to keep udder soft and supple		
Activated charcoal	Activated charcoal can be used on goats to help prevent and treat digestive problems and poisoning		
F10	Disinfectant		
Zinc Sulphate	Footrot	10% Solution	Footbath
Liquid Paraffin & Iodine	Orf/Vuilbek	-	Mix in equal parts and dab on effected area
SUPPLEMENTS AND DIGESTIVE MODIFIERS			
Ovi-Min Gel	Provides essential trace minerals and fat-soluble vitamins to assist in fertility, immunity and growth	Lambs to wean: 5mℓ / 12kg, Weaning and feedlot: 10ml / 25kg, Adult: 20 - 25ml/65kg or dilute 60ml per 100 L water daily for 5 days.	Oral drench.
Kyroligo	Supplement in all animals where nutritional supplementation is required. For use in healthy animals to ensure normal development. Kyroligo can also be used therapeutically, for convalescence after illness and for general debilitation and weakness	5 - 8 mℓ	By deep intramuscular injection only.
B-Co-Bolic	Injectable tonic for Phosphor, Selenium and Vitamin B deficiencies in ruminants. To normalize and/or boost metabolism. To aid in the recovery after disease.	Mature: 3 – 5 mℓ and Lambs / Kids: 2 – 3 mℓ	Intramuscular injection
Rumix	By restoring normal function, ruminal movements are re-established, digestion is regulated, appetite improved and feed conversion promoted.	Adult Goats - A quarter sachet (25 g) twice daily for 4 days,	Oral drench / mix with feed
Calbro Forte	Indicated for Hypocalcaemia and hypomagnesaemia as well as Ketosis	10 – 20 mℓ	Subcutaneous, intramuscular or intravenous injection
Browse Plus	Browse Plus is a specialised formula designed as a drinking water or in feed / lick additive which enhances the animal's digestive process and results in more nutrient utilisation.	1 – 4 g per animal per day	Add to drinking water, licks or feed.
Electro Guard	Assist in the prevention and treatment of symptoms associated with stress related conditions.	3 - 4 mℓ / 10kg	Oral drench.
EQUIPMENT			
1 inch 20 gauge Hypodermic Needles			
3mℓ, 5mℓ & 10mℓ Syringes			
Oral Drench			
Feeding Tube			
Foot bath			
Medical Gloves			
Scalpel			