

Risks in Distance Running for Children

In recent years, there has been a surge in the participation of children in distance running. It is not unusual for an aspiring prepubescent athlete to run 10 to 15 miles daily and to participate in distance races, including marathons (26.2 miles).

Although running is a natural activity that can maintain and improve aerobic fitness, racing and particularly training for long distances have their risks. Distance running may induce musculoskeletal, endocrine, hematologic, thermoregulatory, and psychosocial damage. Most reports on such potential damage have not been evaluated with proper epidemiologic scrutiny. It is unknown whether the risk is greater for children than for adults. Nevertheless, the American Academy of Pediatrics wishes to alert the physician to the presence of such risks. Even without established guidelines and extensive documentation, physicians can give children, parents, and coaches advice that fosters healthy physical and psychosocial growth.

The most common musculoskeletal problems in the young runner are overuse injuries (ie, those that result from a mechanical stress repeated during a long period). These include epiphyseal plate injuries, stress fractures, patellofemoral syndrome, and chronic tendonitis.¹⁻⁴ The incidence of such injuries seems to be related to the total distance covered in training and competition.⁴ Such overuse injuries may lead to a chronic disability (eg, chronic arthritis and growth deformity). Therefore, early medical intervention is important.

Female distance runners often experience delayed menarche.⁵ Its etiology and relevance to health have yet to be established. In most cases, menarche will occur several months after cessation or reduction in volume of training. When a female athlete displays a delayed menarche, a medical

etiology should be considered—rather than assuming causation by exercise.⁶

Although studies have not been performed in prepubertal children, iron depletion, manifested by low serum ferritin concentration, even with a normal hemoglobin concentration, is not uncommon among adolescent runners of both sexes.⁷ An iron-poor diet is probably a main contributor to this disorder. Hemolysis is an often-cited cause of iron loss in runners and is manifested by low serum haptoglobin levels.⁸ One possible mechanism is repetitive foot strike (previously known as “march hemoglobinuria”). Hematuria and mild occult gastrointestinal bleeding have been observed and may contribute to iron depletion.

A child's ability to maintain thermal homeostasis during prolonged running is less efficient than that of an adult, particularly when the climate is very hot or very cold.⁹ This deficiency may result in heat- or cold-related disorders, including heatstroke or hypothermia. Of particular concern is the dehydration that accompanies prolonged running, even if the child is given fluid *ad libitum*.¹⁰ In addition, children take longer than adults to acclimatize to hot, humid climates, which further increases their risk for heat-related disorders.⁹

Psychologic and social problems for the child runner can result from spending long hours in training and setting unrealistic goals. This is similar to the effects of participation in other competitive sports, in which the child may be submitted to inappropriate pressures. A prepubertal child should be allowed to participate for the enjoyment of running, without fear of parental or peer rejection or pressure.

Total mileage (and number of hours) covered by the child during training rather than the distance run on the day of competition may entail the greatest risk to the child's well-being and health. Therefore, suggestions cannot be made for specific maximal racing distances for children. It is important to recognize, however, that heat-related disorders are particularly pronounced in races that exceed 30 minutes in duration.

Until further data are available concerning the

The recommendations in this statement do not indicate an exclusive course of treatment to be followed. Variations, taking into account individual circumstances, may be appropriate.

This statement has been approved by the Council on Child and Adolescent Health.

PEDIATRICS (ISSN 0031 4005). Copyright © 1990 by the American Academy of Pediatrics.

relative risk of endurance running at different ages, the American Academy of Pediatrics recommends that, if children enjoy the activity and are asymptomatic, there is no reason to preclude them from training for and participating in such events. The risks discussed in this statement should be considered by pediatricians when counseling families about the advisability of their children participating in distance running.

COMMITTEE ON SPORTS MEDICINE, 1989–1990

Michael A. Nelson, MD, Chairman

Barry Goldberg, MD

Sally S. Harris, MD

Gregory L. Landry, MD

William L. Risser, MD

Liaison Representatives

Oded Bar-Or, MD, Canadian Paediatric Society

Richard Malacrea, National Athletic Trainers Association

Roswell Merrick, National Association for Sport and Physical Education

AAP Section Liaison

David M. Orenstein, MD, Section on Diseases of the Chest

Arthur M. Pappas, MD, Section on Orthopaedics

REFERENCES

1. Caine DJ, Koenraad JL. Growth plate injury: a threat to young distance runners? *Physician Sportsmed.* 1984;12:118–124
2. Chandy TA, Grana WA. Secondary school athletic injury in boys and girls: a three-year comparison. *Physician Sportsmed.* 1985;13:106–111
3. Micheli LJ, Santopietro F, Gerbino P, Grove MD. Etiologic assessment of overuse stress fractures in athletics. *Nova Scotia Med Bull.* April/June 1980:43–47
4. Orava S, Saarela J. Exertion injuries to young athletes: a follow-up research of orthopaedic problems of young track and field athletes. *Am J Sports Med.* 1978;6:68–74
5. Malina RM. Menarche in athletes: a synthesis and hypothesis. *Ann Hum Biol.* 1983;10:1–24
6. Shangold MM. Menstruation. In: Shangold MM, Mirkin G, eds. *Women and Exercise.* Philadelphia, PA: F.A. Davies; 1988:129–144
7. Nickerson HJ, Tripp AD. Iron deficiency in adolescent cross-country runners. *Physician Sportsmed.* 1983;11:60–66
8. Rowland TW. Iron deficiency and supplementation in the young endurance athlete. In: Bar-Or O, ed. *Advances in Pediatric Sport Sciences.* Champaign, IL: Human Kinetics; 1989;3:169–190
9. Bar-Or O. *Pediatric Sports Medicine for the Practitioner.* New York, NY: Springer Verlag; 1983
10. Bar-Or O, Dotan R, Inbar O, Rotshtein A, Zonder H. Voluntary hypohydration in 10- to 12-year-old boys. *J Appl Physiol.* 1980;48:104–108

Risks in Distance Running for Children
Pediatrics 1990;86;799

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/86/5/799>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.aappublications.org/site/misc/Permissions.xhtml>

Reprints

Information about ordering reprints can be found online:
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Risks in Distance Running for Children

Pediatrics 1990;86;799

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/86/5/799>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 1990 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

