CONTINUOUS

· Belts around belly, must stay close to unit

INTERMITTENT WITH DOPPLER

- Good if you're healthy and low risk
- Listen to baby periodically throughout labor with a handheld Doppler


TELEMETRY/WIRELESS UNIT

If you need to be monitored continuously but want to move around

INTERNAL MONITORS

- FETAL Inserts into baby's scalp
- · IUPC Inserts into uterus, measures strength of contractions





BENEFITS OF FETAL MONITORING

- Provides information about how baby is tolerating labor
- Fetal heart rate patterns give providers an indication of how baby is tolerating labor
- Continuous monitoring may be helpful if there are health concerns

BENEFITS OF INTERNAL MONITORING

- Can provide the most accurate information about how well baby is tolerating labor
- Provides information about strength of contractions (IUPC)
- Can be used for Amnioinfusion (IUPC)
- Can help with finding right dose of pitocin (IUPC)

RISKS OF FETAL MONITORING Monitoring strips can sometimes

- Monitoring strips can sometimes be misinterpreted and do not diagnose issues alone
- Continuous monitoring can increase your risk of cesarean
- Wired and internal monitors may restrict movement

RISKS OF INTERNAL MONITORING

- Infection risks associated with bag of waters being broken
- Injury to baby's scalp (scalp electrode)
- Must stay close to the bed and monitors
- Can be uncomfortable





