

CLASS A: Chest Pain & Respiratory Distress

PROTOCOL(S) USED IN: Chest Pain, Hypertension, Respiratory Distress

PHARMACOLOGY AND ACTIONS:

- A. Vasodilator
- B. Decreases peripheral resistance
- C. Generalized smooth muscle relaxation
- D. Reduces venous tone

INDICATIONS:

- A. Chest, arm, neck pain thought to be related to coronary ischemia.
- B. Angina
- C. Pulmonary edema

CONTRAINDICATIONS:

- A. Hypotension
- B. Hypovolemia
- C. ICP
- D. Aortic Stenosis
- E. Severe bradycardia or tachycardia
- F. Patients who have taken Viagra® (sildenafil citrate) or Levitra® (vardenafil HCl) within 24 hours, or who have taken Cialis® (tadalafil) within 48 hours. Contact OLMC for direction.

SIDE EFFECTS AND NOTES:

- A. Common side effects are headache, flushing, dizziness or burning under the tongue.
- B. Hypotension; IV line should be established prior to administration
- C. Reflex tachycardia
- D. Syncope
- E. May be effective in relieving chest pain due to esophageal spasm
- F. Therapeutic effect is enhanced but adverse effects are increased when patient is upright.
- G. NTG loses potency easily; should be stored in a dark glass container with tight lid and not exposed to heat.

ADULT DOSING:

0.4 mg SL every 3-5 minutes or IV/IO infusion starting at **5 mcg/min** and titrating to effect as long as **systolic BP ≥ 100 mmHg**

PEDIATRIC DOSING: Not recommended in pediatric patients.