## **Suprep, Suflave or Clenpiq Afternoon Prep Instructions**

## The Week BEFORE Colonoscopy

- Pick up your prescription from your pharmacy
- 48 HOURS PRIOR TO PROCEDURE PLEASE DO NOT HAVE ANY NUTS, POPCORN, OR ANY TYPE OF SEEDS.
- STOP ALL BLOOD THINNERS 7 DAYS PRIOR TO PROCEDURE

	When	Start clear liquid diet				
	you wake	Clear liquids include all of the following that are NOT colored RED OR PUPRLE:				
The Day BEFORE Colonoscopy	ир	Gatorade Plain jell-o Popsicles Coffee (no Cream) Apple Juice Kool-Aid Soda Pop Clear broth Tea Water Crystal Light Lemonade (no Pulp) Soda				
		Remember: Do not eat any solid foods the entire day, do not drink orange juice, milk or any milk products.  • Diabetics: skip your dose of diabetic medication the day before and the morning of the procedure				
	6:00 PM	<ul> <li>Use the Split dose instructions on the box of Suprep, Suflave or Clenpiq</li> <li>Take first dose at 6PM</li> </ul>				

The Day of Colonoscopy	6:00 AM	<ul> <li>Take second dose at 6:00 AM.</li> <li>You can have clear liquids up to 6 hours prior to your procedure.</li> <li>Ok to take regular medications. Diabetics; skip your dose of diabetic medication the morning of the procedure</li> </ul>		
		You must have a responsible person to drive you home after your procedure. From check in to check out it will take approximately 3-4 hours total. Be aware that the facility WILL NOT let you leave under any circumstances		
		Arrive at the surgery center		
		Baylor Scott & White Surgicare Rockwall Center		
		825 W Yellowjacket Lane		
		Rockwall, Texas 75087		

Patient signature	Date	