Suprep, Suflave or Clenpiq Morning Prep Instructions

The Week BEFORE Colonoscopy

- Pick up your prescription from your pharmacy
- 48 HOURS PRIOR TO PROCEDURE PLEASE DO NOT HAVE ANY NUTS, POPCORN, OR ANY TYPE OF SEEDS.
- STOP ALL BLOOD THINNERS 7 DAYS PRIOR TO PROCEDURE

2 Days Before your	PLEASE DO NOT HAVE ANY NUTS, POPCORN, OR ANY TYPE OF SEEDS
Procedure	

The Day BEFORE Colonoscopy	When you wake up	 Start clear liquid diet Clear liquids include all of the following that PUPRLE: Gatorade Plain jell-o Popsicles Apple Juice Kool-Aid Soda Pop Tea Water Crystal Light 	Coffee (no Cream) Clear broth			
		Soda				
		 Remember: Do not eat any solid foods the entire day, do not drink orange juice, milk or any milk products. Diabetics: skip your dose of diabetic medication the day before and the morning of the procedure The more you drink and hydrate the better your prep will be! 				
	6:00 PM	 Use the Split dose instructions on the box of Suprep, Suflave or Clenpiq Take first dose at 6PM 				
	10:00 PM	Take second dose of prep at 10:00PM				
	12:00 PM	 NOTHING TO EAT OR DRINK AFTER MIDNIGHT UNTIL AFTER YOUR PROCEDURE It is ok to take your regular medication with a sip of water 				

	You must have a responsible person to drive you home after your procedure. From check in to check out it will take approximately 3-4 hours total. Be aware that the facility WILL NOT let you leave under any circumstances		
The Day of Your Colonoscopy	Arrive at the surgery center Baylor Scott & White Surgicare Rockwall Center 825 W Yellowjacket Lane Rockwall, Texas 75087		

Patient signature	Date	