

Informed Consent For Cataract Surgery And/Or Implantation of an Intraocular Lens

Risks of cataract surgery include, but are not limited to:

1. Complications of removing the natural lens may include hemorrhage (bleeding); rupture of the capsule that supports the IOL; perforation of the eye; clouding of the outer lens of the eye (corneal edema), which can be corrected with a corneal transplant; swelling in the central area of the retina (called cystoid macular edema), which usually improves with time; retained pieces of lens in the eye, which may need to be removed surgically; infection; detachment of the retina, which is definitely an increased risk for highly nearsighted patients, but which can usually be repaired; uncomfortable or painful eye; droopy eyelid; increased astigmatism; glaucoma; and double vision. These and other complications may occur whether or not an IOL is implanted and may result in poor vision, total loss of vision, or even loss of the eye in rare situations. **Additional surgery may be required to treat these complications.**
2. Complications associated with the IOL may include increased night glare and/or halo, double or ghost images, and dislocation of the IOL. Multifocal IOLs may increase the likelihood of these problems. In some instances, corrective lenses or surgical replacement of the IOL may be necessary for adequate visual function following cataract surgery.
3. Complications associated with local anesthesia injections around the eye include perforation of the eye, destruction of the optic nerve, interference with the circulation of the retina, droopy eyelid, respiratory depression, hypotension, cardiac problems, and in rare situations, brain damage or death.
4. If a monofocal IOL is implanted, either distance or reading glasses or contacts will be needed after cataract surgery for adequate vision.
5. Complications associated with monovision. Monovision may result in problems with impaired depth perception. Choosing the wrong eye for distance correction may result in feeling that things are the “wrong way around.” Once surgery is performed, it is not always possible to undo what is done, or to reverse the distance and near eye without some loss of visual quality.
6. Complications associated with multifocal IOLs. While a multifocal IOL can reduce dependency on glasses, it might result in less sharp vision, which may become worse in dim light or fog. It may also cause some visual side effects such as rings or circles around lights at night. It may be difficult to distinguish an object from a dark background, which will be more noticeable in areas with less light. Driving at night may be affected. If you drive a considerable amount at night, or perform delicate, detailed, “up-close” work requiring closer focus than just reading, a monofocal lens in conjunction with eyeglasses may be a better choice for you. If complications occur at the time of surgery, a monofocal IOL may need to be implanted instead of a multifocal IOL.
7. If an IOL is implanted, it is done by a surgical method. It is intended that the small plastic, silicone, or acrylic IOL will be left in the eye permanently.
8. If complications occur at the time of surgery, the doctor may decide not to implant an IOL in your eye even though you may have given prior permission to do so.
9. Other factors may affect the visual outcome of cataract surgery, including other eye diseases such as glaucoma, diabetic retinopathy, age-related macular degeneration; the power of the IOL; your individual healing ability; and, if certain IOLs are implanted, the function of the ciliary (focusing) muscles in your eyes.

10. The selection of the proper IOL, while based upon sophisticated equipment and computer formulas, is not an exact science. After your eye heals, its visual power may be different from what was predicted by preoperative testing. You may need to wear glasses or contact lenses after surgery to obtain your best vision. Additional surgeries such as IOL exchange, placement of an additional IOL, or refractive laser surgery may be needed if you are not satisfied with your vision after cataract surgery.
11. The results of surgery cannot be guaranteed. If you chose a multifocal IOL, it is possible that not all of the near (and intermediate) focusing ability of your eye will be restored. Additional treatment and/or surgery may be necessary. Regardless of the IOL chosen, you may need laser surgery to correct clouding of vision. At some future time, the IOL implanted in your eye may have to be repositioned, removed surgically, or exchanged for another IOL.
12. If your ophthalmologist has informed you that you have a high degree of hyperopia (farsightedness) and/or that the axial length of your eye is short, your risk for a complication known as nanophthalmic choroidal effusion is increased. This complication could result in difficulties completing the surgery and implanting a lens, or even loss of the eye.
13. If your ophthalmologist has informed you that you have a high degree of myopia (nearsightedness) and/or that the axial length of your eye is long, your risk for a complication called a retinal detachment is increased. Retinal detachments can usually be repaired but may lead to vision loss or blindness.
14. Since only one eye will undergo surgery at a time, you may experience a period of imbalance between the two eyes (anisometropia). This usually cannot be corrected with spectacle glasses because of the marked difference in the prescriptions, so you will either temporarily have to wear a contact lens in the non-operated eye or will function with only one clear eye for distance vision. In the absence of complications, surgery in the second eye can usually be accomplished within 3 to 4 weeks, once the first eye has stabilized.

Patient (or person authorized to sign for patient)

Date

Physician Signature

Date

Version 2/27/06