

Marijuana Facts



Marijuana Use Disorder

Some people who use marijuana will develop marijuana use disorder, meaning that they are unable to stop using marijuana even though it's causing health and social problems in their lives.

One study estimated that approximately 3 in 10 people who use marijuana have marijuana use disorder. The risk of developing marijuana use disorder is stronger in people who start using marijuana during youth and who use marijuana more frequently.

Inhaling or Eating Marijuana

- Vaping THC-containing products has been linked to <u>lung</u> <u>injury</u> and even death. Use of concentrates in vaping or dabbing devices delivers very high levels of THC to the user and may carry health risks because of the high concentration or strength of THC being used.
- Eating edibles (foods and drinks that contain marijuana)
 can take longer to kick in compared to smoking marijuana.
 The delayed effect of consuming edibles may increase the
 risk of consuming too much, which can lead to poisoning
 or serious injury. READ

For Immediate Action

- Share these Facts with your faith community.
- Forward to 10 others.

Click here to donate to Illinois Churches in Action (tax deductible)



www.ilcaaap.org 1132 W Jefferson Springfield, IL 60702

STAY CONNECTED



Illinois Church Action on Alcohol and Addiction Problems | 1132 West Jefferson, Springfield, IL 62702

Unsubscribe ilcaaap@sbcglobal.net

Update Profile |Constant Contact Data Notice

Sent byilcaaap@sbcglobal.netpowered by



Try email marketing for free today!