

Pediatric M-Finger, M-Finger/Partial M-Finger



Pediatric M-Fingers, M-Fingers

1. If using a custom silicone socket, add a small amount of water to dampen skin and push your limb into the prosthesis.
2. If a strap is used on the forearm section, secure in place.
3. Adjust BOA reel to the ideal cable tension.
4. Flex wrist to bend fingers, extend wrist to extend fingers.
5. New users should start with a wearing schedule and ease in to a full day of use.
6. To remove the prosthesis, remove the strap, loosen the BOA reel tension and remove limb from prosthesis.
7. Clean inside of silicone liner with fragrant free anti-bacterial wipes daily. Consult with your Doctor and prosthetist regarding preferred cleansing wipes.

Partial M-Fingers

1. Apply wristband to hand.
2. Apply small amount of water to dampen finger/fingers and push each finger into the prosthetic sockets.
3. Adjust cable tension to desired tension by moving the metal cable sleeve up or down. Moving up will tighten the cable tension, moving down will loosen the tension.
4. New users should start with a wearing schedule and ease in to a full day of use.
5. To remove Partial M-Fingers from limb, pull on socket to remove.
6. Remove wrist band.
7. Clean inside of silicone liner daily with fragrant free anti-bacterial wipes. Consult with your Doctor and prosthetist regarding preferred cleansing wipe.

M-Finger



Shown with custom made prosthesis

Pediatric M-Finger



M-Finger



Partial M-Finger



Shown with custom made prosthesis

Partial M-Finger



PARTIAL HAND SOLUTIONS

86 Ladyslipper Lane
Southington, CT, USA 06489
+1 (860) 538-5532

EC

REP

EMERGO EUROPE
Prinsessegracht 20
2514 AP The Hague
The Netherlands

