

2020 HIV DRUG CHART

This quick-reference chart compares antiretroviral (ARV) options for the treatment of HIV, including adult dosing and dietary restrictions. Visit poz.com/drugchart for more info.

*Generic version available in the U.S.

(Pills not shown actual size)



ATRIPLA
efavirenz + tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 600 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take on an empty stomach. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



BIKTARVY
bictegravir + tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 50 mg bictegravir + 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with or without food.



CABENUVA
cabotegravir + rilpivirine
A long-acting injectable regimen. Maintenance dosing requires injections of cabotegravir + rilpivirine every four to eight weeks. A 30-day lead-in period using oral cabotegravir + Edurant (rilpivirine) is required. Take with food. *Approval pending at press time. Photo unavailable.*



COMPLERA
rilpivirine + tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 25 mg rilpivirine + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with a meal.



DELSTRIGO
doravirine + tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 100 mg doravirine + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



DOVATO
dolutegravir + lamivudine
One tablet once a day. Each tablet contains 50 mg dolutegravir + 300 mg lamivudine. Take with or without food.



GENVOYA
elvitegravir + cobicistat + tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 150 mg elvitegravir + 150 mg cobicistat + 10 mg tenofovir alafenamide + 200 mg emtricitabine. Take with food.



JULUCA
dolutegravir + rilpivirine
One tablet once a day. Each tablet contains 50 mg dolutegravir + 25 mg rilpivirine. Take with a meal.



CIMDUO
tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



COMBIVIR *
zidovudine + lamivudine
One tablet twice a day. Each tablet contains 300 mg zidovudine + 150 mg lamivudine. Take with or without food.



DESCOVI
tenofovir alafenamide + emtricitabine
One tablet once a day. Each tablet contains 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with or without food.



EMTRIVA
emtricitabine (also known as FTC)
One 200 mg capsule once a day. Take with or without food.



EPIVIR *
lamivudine (also known as 3TC)
One 300 mg tablet once a day, or one 150 mg tablet twice a day. Take with or without food. Also approved for the treatment of hepatitis B virus but at a lower dose. People living with both viruses should use the HIV dose.



EPZICOM *
abacavir + lamivudine
One tablet once a day. Each tablet contains 600 mg abacavir + 300 mg lamivudine. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.



RETROVIR *
zidovudine (also known as AZT)
One 300 mg tablet twice a day. Take with or without food.



TEMIXYS
tenofovir disoproxil fumarate + lamivudine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take with or without food.



TRUVADA
tenofovir disoproxil fumarate + emtricitabine
One tablet once a day. Each tablet contains 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with or without food.



VIREAD *
tenofovir disoproxil fumarate
One 300 mg tablet once a day. Take with or without food.



ZIAGEN *
abacavir
One 300 mg tablet twice a day, or two 300 mg tablets once a day. Take with or without food. Should be used only by individuals who are HLA-B*5701 negative.



EDURANT
rilpivirine
One 25 mg tablet once a day. Take with food.



INTELENCE
etravirine
One 200 mg tablet twice a day. Take with food.



PIFELTRO
doravirine
One 100 mg tablet once a day. Take with or without food.



SUSTIVA *
efavirenz
One 600 mg tablet once a day, or three 200 mg capsules once a day. Take on an empty stomach or with a low-fat snack. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



VIRAMUNE *
nevirapine
One 200 mg Viramune immediate release (IR) tablet once a day for the first 14 days, then one 400 mg Viramune extended release (XR) tablet once a day. Take with or without food.



EVOTAZ
atazanavir + cobicistat
One tablet once a day. Each tablet contains 300 mg atazanavir + 150 mg cobicistat. Take with food.



KALETRA
lopinavir + ritonavir
Two tablets twice a day, or four tablets once a day, depending on HIV drug resistance. Each tablet contains 200 mg lopinavir + 50 mg ritonavir. Take with or without food.



PREZCOBIX
darunavir + cobicistat
One tablet once a day. Each tablet contains 800 mg darunavir + 150 mg cobicistat. Take with food.



PREZISTA
darunavir
One 800 mg tablet (or two 400 mg tablets) plus one 100 mg Norvir tablet, or one 600 mg tablet plus one 100 mg Norvir tablet twice a day, depending on drug resistance. Take with food.



REYATAZ *
atazanavir
Two 200 mg capsules once a day, or one 300 mg capsule plus one 100 mg Norvir tablet once a day. Take with food.



ODEFSEY
rilpivirine + tenofovir alafenamide + emtricitabine

One tablet once a day. Each tablet contains 25 mg rilpivirine + 25 mg tenofovir alafenamide + 200 mg emtricitabine. Take with a meal.



STRIBILD
elvitegravir + cobicistat + tenofovir disoproxil fumarate + emtricitabine

One tablet once a day. Each tablet contains 150 mg elvitegravir + 150 mg cobicistat + 300 mg tenofovir disoproxil fumarate + 200 mg emtricitabine. Take with food.



SYMFI AND SYMFI LO
efavirenz + tenofovir disoproxil fumarate + lamivudine

One tablet of either Symfi or Symfi Lo once a day. Each tablet of Symfi contains 600 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Each tablet of Symfi Lo (above) contains 400 mg efavirenz + 300 mg tenofovir disoproxil fumarate + 300 mg lamivudine. Take on an empty stomach. Dose should be taken at bedtime to minimize dizziness, drowsiness and impaired concentration.



SYMITUZA
darunavir + cobicistat + tenofovir alafenamide + emtricitabine

One tablet once a day. Each tablet contains 800 mg darunavir + 150 mg cobicistat + 10 mg tenofovir alafenamide + 200 mg emtricitabine. Take with food.



TRIUMEQ
dolutegravir + abacavir + lamivudine

One tablet once a day. Each tablet contains 50 mg dolutegravir + 600 mg abacavir + 300 mg lamivudine. Take with or without food. Should be used only by individuals who are HLA-B*57:01 negative.



RUKOBIA
fostemsavir

One 600 mg tablet once a day for people with HIV treatment experience. Take with or without food.



SELZENTRY
maraviroc

One 150 mg, 300 mg or 600 mg tablet twice a day, depending on other meds used, for people with HIV treatment experience. Take with or without food.



TROGARZO
ibalizumab

Administered intravenously as a single loading (or initial) dose of 2,000 mg followed by a maintenance dose of 800 mg every two weeks for people with HIV treatment experience.



NORVIR *
ritonavir

Six 100 mg tablets twice a day. The full dose of Norvir is rarely used. It is most often used at lower doses to boost the levels of other ARVs in the blood. Take with food.



TYBOST
cobicistat

One 150 mg tablet once a day in combination with ARVs that require boosting. Used only to boost other drugs. Take with food.



ISENTRESS
raltegravir

Two 600 mg Isentress HD tablets (above) once a day for those who are treatment naive or whose virus has been suppressed on an initial regimen of Isentress. One 400 mg Isentress tablet twice daily for people with HIV treatment experience. Take with or without food.



TIVICAY
dolutegravir

One 50 mg tablet once a day for those first starting ARV therapy or for those who have not used an integrase inhibitor in the past. One 50 mg tablet twice a day for treatment-experienced individuals who have HIV that is resistant to other integrase inhibitors and when taken with certain ARVs. Take with or without food.

Integrase Inhibitors

These antiretroviral medications are rarely prescribed and no longer recommended:



APTIVUS
tipranavir



CRIVIAN
indinavir



FUZEON
enfuvirtide



INVIRASE
saquinavir



LEXIVA
fosamprenavir



TRIZIVIR *
abacavir + zidovudine + lamivudine



VIRACEPT
nelfinavir

PK Boosters