Microblading Pre-care and information

Thank you for choosing Kristi PMU and Wax for your service.

As part of your microbading pre care aftercare, it is crucial that you:

- Wear usual daily make up (eyebrow) to the clinic on the day of your treatment
- Make an appointment or order your sensitivity patch test. Test has to be done 48 hours prior to the treatment
- Do not take aspirin, ibuprofen or Coumadin for two days before and two days after the treatment. However you must not discontinue any prescribed medication before consulting your doctor
- Alcohol must not be consumed in the 24 hours prior to the treatment
- Refrain from using sunbeds and avoid natural sun exposure for two weeks before and two weeks after the treatment
- Refrain from waxing 48 hours prior to the appointment
- Refrain from having electrolysis for at least 5 days prior to the treatment
- Eyebrow tinting should not be undertaken one week prior to the treatment

Sensitivity Patch testing:

- 1. Wash your hands.
- 2. Use clean cotton buds.
- 3. Place the pigment sample behind your left ear.
- 4. Use a another clean cotton buds.
- 5. Place the numbing cream behind your right ear.
- 6. After 24 hours wipe the products of with a cotton pad, if there is no visible reaction, such as redness, swelling, irritation or itching, than treatment can be proceed

Please not that, negative sensitivity patch test result does not guarantee that you will not have an allergic reaction.

Please not colour enclosed is not necessary the colour will be used for your treatment, it is for patch test purposes only.

Important treatment information:

- This treatment is a form of cosmetic tattooing.
- Permanent cosmetics may require multiple treatment sessions.
 For best results, you will be required to return for at least one retouch treatment between 4

and 12 weeks after the initial treatment.

- Immediately after the procedure, during the healing process the results can appear darker.
- The pigment colour will fade during the healing process. It is your responsibility to keep the treatment area clean and free from contaminants.
- Infections can occur.
- Allergic reactions to topical anaesthetics and pigment can occur.
- There may be swelling and redness following the procedure.
- You may experience minor bleeding.
- A retouch cannot be performed until the treatment area is completely healed.
- In rare cases, the pigment could migrate under the skin.
- Application of permanent cosmetics can be painful.
- The pigment will fade over time and a colour-boosting procedure will need to be performed approximately 12 to 18 months after the original procedure.
- A negative sensitivity patch test will not guarantee that you will not have an allergic reaction.
- There are few effective methods for pigment removal.
- Possible scarring and inconsistency of color can occur.
- After having a permanent cosmetics procedure, discuss with a doctor before having an MRI scan.
- Those receiving an eye treatment may require someone to drive them home after the appointment.
- Corneal abrasion may occur during eyeliner procedures. However, this is rare.
 If you have experienced previous problems with cold sores/herpes and wish to receive a lip treatment, you may have a reoccurrence of cold sores following the procedure.
- Anti-herpes medication is available over the counter or on prescription and has been shown to prevent or minimize such outbreaks.

- Permanent cosmetics procedures can trigger outbreaks of dormant herpes whether or not you are aware of its existence, but it cannot be the cause of herpes.
- Lip treatments will appear dry and flaky for approximately one week following the procedure.
- Camouflage procedures are experimental in nature and procedures will need to be tested for skin colour matching.
- Possible scarring, inconsistency of colour and loss of eyelashes may occur.