



# 2024 Napa Primary Care Conference

## Infectious Disease in the Active and Athletic Patient

Dennis K-Borna, MD, FACSM, FAAFP

*No disclosures to report*

# Goals

- Recognize the important role that infections play in sports
- Identify and treat cutaneous infections, including those with bacterial, viral, and fungal etiologies
- Identify and treat “non-cutaneous” or systemic infections
- Understand appropriate return to play strategies
- Understand the simple, but important ways to help prevent infectious disease in sports





# Importance


- Most infections are contagious
- The close proximity of athletes to each other is a key risk factor
- The close contact involved in many sports also puts certain athletes at risk
- Exhaustive exercise increases the risk and severity of some viral illnesses

# Importance

- Even relatively mild systemic infections can dramatically decrease an athlete's performance
- Cutaneous infections can be severely limiting, and can be disqualifying
- Both systemic and cutaneous infections have decimated entire teams




# Cutaneous Infections



- **Bacterial**

- Cellulitis/Abscesses
- Impetigo
- Folliculitis
- Otitis externa
- Conjunctivitis
- Hand infections
- Corynebacteria infections



- **Viral**


- HSV
- molluscum
- warts



- **Fungal**

- tinea


# Non-cutaneous Infections



- **Mononucleosis**
- **URI and other viral syndromes (Covid-19)**
- **GI infections**
- **Bloodborne pathogens**


# Cutaneous Infections

- **Bacterial**
  - Cellulitis/Abscesses
  - Impetigo
  - Folliculitis
  - Otitis externa
  - Conjunctivitis
  - Hand infections
    - paronychia
    - felons
  - Corynebacteria infections



# Cutaneous Infections


- **Viral**
  - Herpesvirus infections
    - Herpes Gladiatorum
    - Herpes Zoster
    - Herpetic Whitlow
  - Molluscum contagiosum
  - Warts






# Cutaneous Infections

- **Fungal**
  - Tinea capitis
  - Tinea corporis
  - Tinea cruris
  - Tinea pedis




# Bacterial Cutaneous Infections




- Cellulitis/Abscesses
  - can occur just about anywhere there is skin
  - athletes are particularly susceptible due to abrasions and physical contact
  - *Staph aureus* and *Strep pyogenes* are the most common pathogens
  - be wary of *Pseudomonas*, *Aeromonas*, and *Vibrio* in water athletes
  - MRSA is becoming more prevalent

# Cellulitis



- Treatment
  - Cephalexin 250-500mg po qid for 5-14 days
  - Dicloxacillin 250-500mg po qid for 5-14 days
  - Amox-Clavulanate 875/125 po bid for 5-14 days
  - Macrolides (erythromycin, azithromycin, clarithromycin)
  - Quinolones (moxifloxacin, levofloxacin)

# Abscesses




- Treatment
  - Incision and drainage
  - pack with packing gauze
  - oral antibiotics for mild-moderate cases
  - IV antibiotics for severe cases





# Bacterial Cutaneous Infections



- MRSA infections
  - now account for about 30% of all *Staph* infections
  - about 20% of cases in sports require IV antibiotics
  - suspect abscesses/boils and all non-healing infections initially treated with beta-lactams



# MRSA Infections

- Treatment


- Oral antibiotics

- Trimethoprim-sulfamethoxazole (Bactrim/Septra)
    - Clindamycin
    - Rifampicin
    - Tetracycline/doxycycline/minocycline


- IV antibiotics

- Vancomycin
    - Linezolid (Zyvox) -- PO or IV

- For recurrences in patients with known carrier status, treat with intranasal mupirocin (Bactroban)




# Bacterial Cutaneous Infections




- Impetigo
  - superficial skin infection, usually produced by *Staph aureus* and  $\beta$ -hemolytic *Strep*
  - bullous and nonbullous (more common)
  - vesicles on erythematous base with honey-crusted lesions; vesicles often progress to pustules
  - most common in children and young adults
  - more often in warm, humid environments

# Impetigo






# Impetigo



- Treatment
  - Mupirocin ointment tid for 7-10 days
  - oral cephalexin or dicloxacillin q6hrs for 7-10 days if more widespread
  - Prevent spread!




# Bacterial Cutaneous Infections



- Folliculitis
  - infection of hair follicles
  - usually caused by *Staph aureus*
  - “hot tub” folliculitis often caused by gram-negative bugs
  - usually occurs in areas of friction, especially in shave areas
  - can progress to abscesses




# Folliculitis



- Treatment
  - most cases with heal spontaneously in 5-10 days
  - warm compresses can help
  - oral dicloxacillin/cephalexin for 7-14 days
  - worse cases might need 3+ weeks of treatment
  - incision and drainage for fluctuant lesions
  - astringents can help with prevention
  - decrease/avoid shaving
  - use *clean* blades when you do shave

# Bacterial Cutaneous Infections




- Ear infections (Otitis externa)
  - increased risk for athletes training in hot, humid weather, and especially *water sports*
  - also occurs with use of foreign objects (Q-tips)
  - culprit often mixed flora with *Pseudomonas*
  - red, edematous ear canal with significant pain







# Otitis Externa




- Treatment
  - Cortisporin Otic drops tid-qid for ~10 days
    - use suspension if tympanic membrane is ruptured
    - otherwise, use the solution
    - may need to use ear wick if canal is extremely swollen
  - For severe/resistant cases, an oral quinolone (e.g. ciprofloxacin) may be used
  - Consider possibility of **fungal** otitis externa if not improving with treatment

# Conjunctivitis



- Majority are viral, but bacterial causes are usually Staph or Strep
- Highly transmissible, so contact sport athletes should be kept out until resolved
- Treatment options for 5-7 days
  - Erythromycin ointment
  - Trimethoprim-polymyxin B drops
  - Ciprofloxacin or ofloxacin drops (contact lens wearers)

# Bacterial Cutaneous Infections




- Hand Infections

- Paronychia

- infection around the nail folds
    - may be acute (*Staph*) or chronic (fungal)
    - treat acute cases with warm compresses, I&D if abscess is present; nail removal if subungual abscess or ingrown nail is present
    - oral antibiotics covering Staph (e.g. cephalexin or dicloxacillin) for 7-10 days if suppurative



# Bacterial Cutaneous Infections



- Hand Infections

- Felon

- close proximity to joint
    - can lead to osteomyelitis
    - x-ray can provide idea of bone involvement
    - treatment is with *good* incision and drainage
    - hand surgery consult is usual




infections  
or  
antibiotics






# Viral Cutaneous Infections

- Herpesvirus infections
  - Herpes Gladiatorum
  - Herpes Zoster
  - Herpetic Whitlow
- Molluscum contagiosum
- Warts



# Viral Cutaneous Infections



- Herpes Gladiatorum
  - caused by either HSV-1 or HSV-2
  - affects about 2-8% of high school and collegiate wrestlers
  - HSV incubation period is 5-10 days
  - skin lesions improve in 2-21 days
  - “groups of vesicles on an erythematous base”
  - stinging/burning pain of herpes
  - may have associated fever and malaise

# Viral Cutaneous Infections

- Herpes

- caus

- affect

- colle

- HSV

- skin


- “gro

- sting

- may




s base”







# Herpes Gladiatorum




- Treatment
  - oral acyclovir can shorten the time course, but must be started early
  - valacyclovir can decrease the risk of recurrence
    - 7-10 days of treatment
  - benzoyl peroxide + aggressive drying can reduce the risk of secondary bacterial infection
  - lesions must be crusted over for at least 72 hours and covered before return to play
    - At least 5 days of antiviral treatment to return






# Herpes Zoster



- Similar to herpes gladiatorum, but forms along a dermatomal pattern
- Antiviral therapy with 72 hours of symptoms
- Associated with more pain from neuritis
- Analgesics initially for neuritis pain
- Tricyclic antidepressants may be added for refractory neuropathic pain


# Viral Cutaneous Infections

- Herpetic Whitlow
  - infection of the hand (usually one or more fingers) caused by HSV-1 and HSV-2
  - often appears very similar to paronychia or felon
  - axillary lymphadenopathy symptoms and constitutional are not uncommon
  - distal pulp space is swollen but soft
  - characteristic vesicles are present
  - self-limited; oral acyclovir may be helpful if systemic symptoms are present




# Viral Cutaneous Infections

- Herpes



on

# Viral Cutaneous Infections




- Molluscum Contagiosum
  - “umbilicated skin-colored papules” 2-4mm
  - caused by a poxvirus
  - risk factors include close contact, skin abrasion, swimming pools and hot tubs
  - frequently self-limited, but liquid nitrogen or electrocautery can hasten resolution

# Viral Cutaneous Infections

- Molluscum Contagiosum


- “umbilicated” papules, 2-4mm
- caused by poxvirus
- risk factors include skin abrasion, swimming in pools
- frequently treated with cryotherapy or electrocautery






# Viral Cutaneous Infections

- Warts (verrucae)
  - caused by papillomaviruses
  - increased in sports involving calluses
  - treat with cryodestruction or topical salicyclic acid
  - surgical removal can be done for refractory cases




# Viral Cutaneous Infections

- Plantar warts
  - can be significantly limiting to runners and other athletes
  - similar treatments to other warts, but can be more stubborn
  - laser
  - intralesional immunotherapy
  - duct tape?
    - 6 days, 1 day off, repeated weekly




# Cutaneous Infections

- **Fungal**
  - Tinea capitis
  - Tinea corporis
  - Tinea cruris
  - Tinea pedis




# Fungal Cutaneous Infections



- Tinea Capitis
  - “ringworm” of the scalp
  - typically round patches of scale, usually with associated bald patches
  - 90% caused by *Trichophyton tonsurans*
  - oral agents are the treatment of choice:
    - Griseofulvin for 6-12 weeks
    - Itraconazole 3-5mg/kg/day for 6 weeks
    - Terbinafine (Lamisil) for 2-4 weeks




# Fungal Cutaneous Infections



- Tinea Corporis
  - well-marginated scaling circular plaques with central clearing
  - *Trichophyton rubrum* most common
  - KOH scrapings showing branching hyphae with septa
  - treat with topical antifungal creams bid for 2 weeks
  - resistant or extensive cases may require 2-4 weeks of oral therapy



# Fungal Cutaneous Infections



- Tinea Corporis

- well-
- clear
- *Trich*
- KOH
- septa
- treat
- resist
- oral t




entral

eks

ks of




# Fungal Cutaneous Infections




- Tinea Cruris
  - “jock itch” is characterized by crescent-shaped large, well-margined erythematous plaques in the crural folds that burn and itch
  - typically caused by *Microsporum*, *Trichophyton*, and *Epidermophyton*
  - treated with topical antifungals bid for 2-3 weeks  
educate patients about preventive measures

# Fungal Cutaneous Infections




- Tinea Pedis
  - “athlete’s foot” is characterized by itching, burning, scaling, eruptions on the foot, usually between toes
  - skin may be macerated
  - vesicles may be present
  - various *Trichophyton* species are common, resulting in variations in appearance

# Tinea Pedis




# Tinea Pedis





# Tinea Pedis




# Topical Antifungals



Antifungal	<b>Tinea Corporis</b> Effectiveness	<b>Tinea Cruris</b> Effectiveness	<b>Tinea Pedis</b> Effectiveness	Cost
<b>Terbinafine</b>	High	High	High	Moderate-High
<b>Naftifine</b>	High	High	High	Moderate-High
<b>Clotrimazole</b>	High	High	Moderate-High	Low (OTC)
<b>Miconazole</b>	Moderate-High	Moderate-High	Moderate-High	Low (OTC)
<b>Econazole</b>	Moderate-High	Moderate-High	Moderate-High	Moderate
<b>Ketoconazole</b>	Moderate-High	Moderate-High	Moderate-High	Moderate
<b>Ciclopirox</b>	Moderate-High	Moderate-High	Moderate-High	High
<b>Tolnaftate</b>	Moderate	Moderate	High	Low (OTC)

# Fungal Cutaneous Infections



- Tinea Pedis
  - treat with topical antifungal creams bid for 2-4 weeks; may mix with mild topical steroid cream initially
  - Burow's solution soaks often helpful for symptoms
  - oral agents are an option for tough cases, and for cases with extensive nail involvement
  - **nystatin** is *not* effective for dermatophyte infections

# Corynebacterium Infections

- Can often mimic athlete's foot and/or plantar warts
- characterized by foul foot odor with pitted keratolysis, and often with associated hyperhidrosis
- often will have “coral red fluorescence” under Wood's lamp
- treated with proper foot care, and sometimes with topical aluminum chloride (Drysol) and/or topical 2% erythromycin solution

# Corynebacterium Infections

- Can often

Figure: Courtesy




Figure 1. Pitted  
sent as multipl  
bearing areas c



- or planter


y of Dirk M. Elston, MD,  
ardner, MD



of figure 1 shows  
/ pits (arrow) on the  
sole. The lesions and  
odor pointed to the  
keratolysis.




# Non-cutaneous Infections



- Athletes are subject to many of the same infections as the general population
- Infection incidence and severity appear to increase with severe exertion
- Prevention is of utmost importance

# Non-cutaneous Infections



- Mononucleosis
- Upper respiratory infections and other viral syndromes
- Gastrointestinal infections
- Bloodborne pathogens

# Mononucleosis




- Signs & Symptoms
  - syndrome of malaise, headache, fatigue, anorexia, and myalgias
  - exam findings include tonsillar enlargement, cervical lymphadenopathy, soft palate petechiae, and splenomegaly
  - may also have atypical lymphocytosis and elevated liver function tests

# Mononucleosis



- Typically caused by Epstein-Barr virus
- Incubation period of 30-45 days
- Self-limited illness, but complications include splenomegaly with splenic rupture
- Return to play recommendations vary, but all athletes should be withheld for at least 3 weeks

# Non-cutaneous Infections




- Upper respiratory infections
  - usually include illnesses of infectious rhinitis, pharyngitis, sinusitis, and bronchitis
  - there is increased risk of URIs with longer running, and with high-intensity exercise
  - *moderate* exercise may decrease the risk of URIs
  - “above the neck rule”



# Non-cutaneous Infections

- Gastrointestinal infections
  - beware of “weird” organism infections in water athletes
    - *Leptospirosis*
    - *Giardiasis*
    - *Cryptosporidiosis*
  - the primary concern of treatment is hydration
    - oral fluids for mild dehydration
    - IV fluids for more pronounced dehydration or metabolic disturbances
  - oral antibiotics for nonviral pathogens



# Non-cutaneous Infections

- Bloodborne pathogens


- HIV

- the asymptomatic HIV+ athlete can still perform at a very high level, but there may be risks to extreme training and overtraining
    - (almost) no confirmed HIV transmissions in sports to date

- Hepatitis


- some patients can tolerate sports quite well
    - clinical signs and symptoms should guide return to play

# COVID-19




- There is a strong correlation with physical activity and better Covid-19 outcomes
- Myocarditis incidence estimated at 0.5-3%
- Symptoms to look for:
  - Chest pain
  - Shortness of breath out of proportion to URI
  - Palpitations
  - Syncope

# COVID-19



- **Asymptomatic or mild cases**
  - Should be improving for a minimum of 1 day prior to return to physical activity progression
  - 3 day return to play after resolution of symptoms
  - Masking for 10 days from symptom onset
  - Monitor for chest pain, shortness of breath out of proportion to URI, palpitations, or syncope

# COVID-19



- **Moderate symptoms**

- $\geq 4$  days of fever  $> 100.4$
- $\geq 1$  week of myalgia, chills, lethargy, or non-ICU hospital stay
- Physician evaluation
- EKG
- 5 day return to play, after a minimum of 1 day of symptom resolution
- Masking for 10 days from symptom onset

- **Severe symptoms**

- 3-6 months, with a Cardiology clearance



# Prevention

- Athletes should minimize contact with people who are obviously ill
  - this may include avoiding crowds, travel and young children
  - masking and distancing
- Keep intertriginous areas and feet dry as much as possible
- Shower after all events, and before events that involve close skin contact (e.g. wrestling)
- WASH YOUR HANDS!

# Prevention



59

# Prevention

- Keep clothing, equipment, and facilities sanitary
- Assure that athletes' immunizations are up-to-date
- Avoid overtraining, sleep deprivation, and improper nutrition
- Safeguard the water supply




# Prevention

- The athletes should be advised that it is their responsibility to report all wounds and injuries in a timely manner, including those recognized before the sporting activity.
- Post event consideration should include reevaluation of any wounds sustained during the sporting event.
- The care provider managing an acute blood exposure must follow the guidelines of universal precautions.



# Prevention Vaccines





# Prevention Vaccines



# Conclusions

- Infections play a major role in sports.
- Infections can be either cutaneous, or more systemic, and can have a number of different etiologies.
- While prompt recognition and treatment of infections is very important, prevention is by far the most critical step in this aspect of sports medicine.

**Thank you!**

