

High-Intensity Interval Training Increases Cardiac Output and $\dot{V}O_{2\max}$

TODD A. ASTORINO¹, ROSS M. EDMUNDS^{1,2}, AMY CLARK¹, LEESA KING^{1,3}, RACHAEL A. GALLANT¹, SAMANTHA NAMM¹, ANTHONY FISCHER¹, and KIMI M. WOOD¹

¹Department of Kinesiology, California State University, San Marcos, San Marcos, CA; ²Department of Physical Therapy, SUNY, Stony Brook, Stony Brook, NY; and ³National College of Natural Medicine, Portland, OR

ABSTRACT

ASTORINO, T. A., R. M. EDMUNDS, A. CLARK, L. KING, R. A. GALLANT, S. NAMM, A. FISCHER, and K. M. WOOD. High-Intensity Interval Training Increases Cardiac Output and $\dot{V}O_{2\max}$. *Med. Sci. Sports Exerc.*, Vol. 49, No. 2, pp. 265–273, 2017. Increases in maximal oxygen uptake ($\dot{V}O_{2\max}$) frequently occur with high-intensity interval training (HIIT), yet the specific adaptation explaining this result remains elusive. **Purpose:** This study examined changes in $\dot{V}O_{2\max}$ and cardiac output (CO) in response to periodized HIIT. **Methods:** Thirty-nine active men and women (mean age and $\dot{V}O_{2\max}$ = 22.9 ± 5.4 yr and 39.6 ± 5.6 mL·kg⁻¹·min⁻¹) performed HIIT and 32 men and women (age and $\dot{V}O_{2\max}$ = 25.7 ± 4.5 yr and 40.7 ± 5.2 mL·kg⁻¹·min⁻¹) were nonexercising controls (CON). The first 10 sessions of HIIT required eight to ten 60 s bouts of cycling at 90%–110% percent peak power output interspersed with 75 s recovery, followed by randomization to one of three regimes (sprint interval training (SIT), high-volume interval training (HIIT_{HI}), or periodized interval training (PER) for the subsequent 10 sessions. Before, midway, and at the end of training, progressive cycling to exhaustion was completed during which $\dot{V}O_{2\max}$ and maximal CO were estimated. **Results:** Compared with CON, significant ($P < 0.001$) increases in $\dot{V}O_{2\max}$ in HIIT + SIT (39.8 ± 7.3 mL·kg⁻¹·min⁻¹ to 43.6 ± 6.1 mL·kg⁻¹·min⁻¹), HIIT + HIIT_{HI} (41.1 ± 4.9 mL·kg⁻¹·min⁻¹ to 44.6 ± 7.0 mL·kg⁻¹·min⁻¹), and HIIT + PER (39.5 ± 5.6 mL·kg⁻¹·min⁻¹ to 44.1 ± 5.4 mL·kg⁻¹·min⁻¹) occurred which were mediated by significant increases in maximal CO (20.0 ± 3.1 L·min⁻¹ to 21.7 ± 3.2 L·min⁻¹, $P = 0.04$). Maximal stroke volume was increased with HIIT ($P = 0.04$), although there was no change in maximal HR ($P = 0.88$) or arteriovenous O₂ difference ($P = 0.36$). These CO data are accurate and represent the mean changes from pre- to post-HIIT across all three training groups. **Conclusions:** Increases in $\dot{V}O_{2\max}$ exhibited in response to different HIIT regimes are due to improvements in oxygen delivery. **Key Words:** INTERVAL EXERCISE, MAXIMAL OXYGEN UPTAKE, STROKE VOLUME, THORACIC IMPEDANCE, FICK EQUATION, CYCLE ERGOMETRY

In untrained individuals, significant improvements in maximal oxygen uptake ($\dot{V}O_{2\max}$) occur in response to 2–12 wk of high-intensity interval training (HIIT) (5,12,26,30,35). Data from a meta-analysis (6) show a mean increase in $\dot{V}O_{2\max}$ equal to 0.51 L·min⁻¹ in response to 6–13 wk of HIIT, with higher-volume HIIT eliciting even greater improvements (0.8–0.9 L·min⁻¹) in $\dot{V}O_{2\max}$. Compared with moderate-intensity continuous training (MICT), results from some studies (9,38) exhibit similar increases in $\dot{V}O_{2\max}$ in response to HIIT, although this is not a universal finding (30). This discrepancy in $\dot{V}O_{2\max}$ response to exercise training across studies is likely due to differences in volume and intensity of each HIIT regime as well as the influence of genetics on $\dot{V}O_{2\max}$ response to training (8). This increase

in $\dot{V}O_{2\max}$ is important because it increases exercise tolerance and long-term health status (28). For example, a 3.5-mL·kg⁻¹·min⁻¹ increase in $\dot{V}O_{2\max}$ is associated with a 13% and 15% lower risk of all-cause mortality and cardiovascular event, respectively (23).

Despite the widely seen increases in $\dot{V}O_{2\max}$ demonstrated in response to HIIT, the specific adaptations explaining this outcome remain elusive. The Fick equation states that oxygen uptake is determined by the product of cardiac output (CO) and arteriovenous oxygen difference (a-vDO₂). In untrained men and women, Daussin et al. (12) reported that maximal CO (CO_{max}) and a-vDO₂ are increased in response to 8 wk of HIIT which led to significant improvements in $\dot{V}O_{2\max}$. Similarly, Warburton et al. (38) showed significant increases in CO_{max} and $\dot{V}O_{2\max}$ in response to 12 wk of HIIT in untrained men. In contrast, MacPherson et al. (25) demonstrated a 12% increase in $\dot{V}O_{2\max}$ in response to 6 wk of treadmill sprint interval training (SIT) despite no increase in CO_{max}, which implies that peripheral adaptations mediate the improvement in $\dot{V}O_{2\max}$. In another study (14), resting and/or submaximal measures of stroke volume (SV) and CO assessed with echocardiography were increased in response to HIIT, although no maximal values were determined which does not allow identification of adaptations responsible for the training-induced improvements in $\dot{V}O_{2\max}$. Across all

Address for correspondence: Todd Anthony Astorino, Ph.D., Department of Kinesiology, California State University, San Marcos, 333. S. Twin Oaks Valley Road, UNIV 320, San Marcos, CA 92096-0001; E-mail: astorino@csusm.edu.
Submitted for publication June 2016.
Accepted for publication August 2016.

0195-9131/17/4902-0000/0
MEDICINE & SCIENCE IN SPORTS & EXERCISE®
Copyright © 2017 by the American College of Sports Medicine

DOI: 10.1249/MSS.0000000000001099

studies, the participant population, specific characteristics of HIIT, and method of CO determination varied, which may elicit some of the dissimilar findings reported. Overall, changes in $\dot{V}O_{2\max}$ in response to HIIT are poorly understood despite significant increases in $\dot{V}O_{2\max}$ seen after as little as six to 10 sessions of training (5,14).

Although previous findings show similar and in some cases superior adaptations with HIIT versus MICT, the optimum HIIT regimen is unknown, as in most studies, adaptations to interval training are compared with those from MICT. Only a few studies have directly compared changes in $\dot{V}O_{2\max}$ between unique HIIT regimes, and these data are equivocal. In untrained women (5), similar increases in $\dot{V}O_{2\max}$ were exhibited between HIIT regimes differing in absolute intensity. Findings from Matsuo et al. (26) in untrained men showed similar increases in $\dot{V}O_{2\max}$ between high-volume HIIT and low-volume SIT. In contrast, only high-volume HIIT led to significant increases in $\dot{V}O_{2\max}$ in overweight individuals compared with low-volume HIIT or MICT (7).

In each study (5,7,26), the specific duration, intensity, and recovery duration differed across HIIT regimes. However, in no study was the actual structure of HIIT modified during the course of training. In the discipline of resistance training, periodization is widely used (36) to minimize onset of plateau and attempt to sustain physiological adaptations. Because increases in $\dot{V}O_{2\max}$ seem to level off during prolonged HIIT when only intensity is adjusted (5,31), there seems to be potential for periodized training within the domain of HIIT. In addition, it could be argued that constantly modifying the structure of HIIT may improve the participants' perception of training by providing an ever-changing training stimulus. To our knowledge, no study has incorporated periodized HIIT in an attempt to augment adaptations to training. In the present study, we incorporated 10 sessions of low-volume HIIT to rapidly and significantly enhance $\dot{V}O_{2\max}$, as widely reported (5,14), and subsequently randomized participants to 1 of 3 HIIT regimes consisting of an additional 10 sessions to examine if periodization of subsequent training continues to augment $\dot{V}O_{2\max}$. Furthermore, CO was assessed throughout training to identify if improved central O_2 delivery is responsible for potential increases in $\dot{V}O_{2\max}$ seen in response to HIIT.

METHODS

Participants. In response to word-of-mouth and electronic bulletins posted between January 2014 and February 2016, 192 individuals initially contacted the primary investigator to potentially participate in the study. Of these prospective participants, 77 enrolled in the study, 43 whom initiated HIIT and 34 nonexercising controls. Physical characteristics of participants who completed all requirements of the study are shown in Table 1. Participant ethnicity included whites ($n = 41$), Hispanic ($n = 24$), African-American ($n = 2$), Asian/Filipino ($n = 7$), and Middle Eastern ($n = 3$) and reflected the ethnic composition of the University

TABLE 1. Baseline participant characteristics (mean \pm SD).

Parameter	HIIT + SIT	HIIT + HIIT _{HI}	HIIT + PER	CON
Age (yr)	22.5 \pm 5.8	21.9 \pm 1.9	23.1 \pm 5.5	25.8 \pm 5.8*
Gender ($n = M/F$)	7/7	7/6	5/7	15/17
Height (cm)	172.0 \pm 7.8	174.4 \pm 9.6	170.9 \pm 9.9	172.2 \pm 11.3
Mass (kg)	68.5 \pm 10.3	69.6 \pm 11.4	68.2 \pm 7.0	72.2 \pm 14.3
$\dot{V}O_{2\max}$ (mL·kg ⁻¹ ·min ⁻¹)	39.6 \pm 6.8	41.1 \pm 4.9	39.5 \pm 5.3	40.5 \pm 4.9
PPO (W)	226.9 \pm 36.6	240.0 \pm 34.4	229.2 \pm 51.7	243.0 \pm 40.9
PA (h·wk ⁻¹)	6.7 \pm 2.3	5.5 \pm 2.1	6.0 \pm 2.4	6.2 \pm 2.4

PPO from progressive exercise test.

M, male; F, female; PA, habitual physical activity.

* $P < 0.05$ compared to all HIIT groups.

community. All performed a minimum of 150 min·wk⁻¹ of physical activity including resistance training, noncompetitive sport, aerobic exercise, surfing, and so on, which was confirmed with a questionnaire (29). Inclusion criteria included healthy, body mass index less than 30 kg·m⁻², weight stable in the last 12 months, nonsmoker, as well as lack of knee pain and medication/supplement use which may modify study outcomes. All women enrolled in the study were eumenorrheic. Written informed consent was obtained from all participants before initiating the study, whose procedures were approved by the university institutional review board and were conducted in accordance with the Declaration of Helsinki (1964). During the study, four withdrew from the experimental group due to lack of time ($n = 2$), unrelated injury ($n = 1$), and failure to adhere to inclusion criteria ($n = 1$), and two controls did not complete the follow-up trial due to cessation of habitual exercise outside of the study.

The data from 30 of these participants are presented elsewhere (3). The previous study focused on changes in $\dot{V}O_{2\max}$ and CO in response to the initial 10 sessions of low-volume HIIT; whereas, the current study allowed us to present new findings in a larger sample concerning effects of 20 sessions of HIIT regimes differing in structure on these outcomes.

Experimental design. At baseline, $\dot{V}O_{2\max}$ and hemodynamics were measured in all individuals participating in the study. Before this trial and all subsequent assessments, participants completed an overnight fast (12 h) and abstained from medication use for 12 h and physical activity for 48 h. Participants subsequently completed 20 sessions of periodized HIIT with time of day maintained within participants. The posttraining assessment was performed 48–120 h after the final session of HIIT. Participants were advised to maintain their dietary intake and physical activity during the study.

Assessment of body composition. Initially, height and body mass were determined using a balance beam scale with stadiometer (Health-o-Meter; Creative Health Products, Ann Arbor, MI). Subsequently, subcutaneous fat was recorded using a metal skinfold caliper (Lange, Santa Cruz, CA) at the chest, abdominal, and thigh (men) and triceps, suprailiac, and thigh (women) (20,21) in rotational order following standardized procedures (19). If scores differed by more than 10%, a third score was obtained and an average value was calculated from the two scores meeting this criterion. Gender- and ethnicity-specific equations were used to calculate body density and percent body fat.

Assessment of $\dot{V}O_{2\max}$. After 5 min of resting gas exchange data were obtained to yield resting values of $\dot{V}O_2$ (3–4 mL·kg⁻¹·min⁻¹), participants initiated progressive cycling on an electrically braked cycle ergometer (Velotron DynaFit Pro, RacerMate, Seattle, WA) consisting of 7 min at 30 (women) or 40 W (men) followed by 20 W increases in power output every 3 min until mean RER was greater than 1.0 for an entire stage, after which power output was increased by 20 W·min⁻¹ until voluntary exhaustion (cadence < 50 rpm) and attainment of maximal oxygen uptake ($\dot{V}O_{2\max}$) and peak power output (PPO). Maximal oxygen uptake was determined as the mean of the two highest values attained during exercise from any 30-s period. In our laboratory, coefficient of variation and typical error for $\dot{V}O_{2\max}$ determined from this progressive protocol are equal to 3.0% and 1.1 mL·kg⁻¹·min⁻¹, respectively. Attainment of $\dot{V}O_{2\max}$ was confirmed with constant load cycling to exhaustion at 110% PPO performed 10 min after the initial bout (27). During exercise, gas exchange data ($\dot{V}O_2$, ventilation (V_E), carbon dioxide production (VCO_2), and RER) were obtained every 15 s using a metabolic cart (ParvoMedics TrueOne, Sandy, UT) which was calibrated before exercise according to the manufacturer's recommendation. Gas exchange data were visually inspected, and any values differing by more than two standard deviations from the mean value were removed from subsequent analysis. PPO from this trial was used to set intensities for resultant HIIT. $\dot{V}O_{2\max}$ assessment was repeated after 10 and 20 sessions of training, respectively.

Assessment of hemodynamic function. An impedance cardiograph device (Physioflow Enduro, Manatec, Strasbourg, France) was used to evaluate hemodynamic function. This method has been described in detail elsewhere (10,32) and is valid and reliable at rest and during exercise up to $\dot{V}O_{2\max}$. This method detects changes in transthoracic impedance during phases of the cardiac cycle to calculate SV, which is multiplied by HR to estimate CO.

Participants entered the laboratory and were required to sit quietly for approximately 5 min. An alcohol swab was used to clean the neck, right chest, trunk at V6, and spine, and then an electrode gel (NuPrep; Weaver and Company, Aurora, CO) was rubbed into these areas, and the skin was further cleaned with a paper towel. Two sets of electrodes (Skintact ECG electrodes; Leonhard Lang GmbH, Innsbruck, Austria), one electrode transmitting and the other sensing, were applied above the supraclavicular fossa at the left base of the neck and at the height of the xiphoid on the spine. Another pair of electrodes (one placed on the right chest and another at V6) was used to monitor the ECG trace. Once applied, these leads were taped to the skin to minimize movement. The participant was seated on the cycle ergometer for 2 min, blood pressure was manually recorded twice at the antecubital space with a stethoscope and sphygmomanometer (Omron Health Care Inc., Vernon Hills, IL), and they were told not to talk and remain motionless. Then, the device was calibrated after a 30-beat procedure using the baseline blood pressure value, which was averaged.

Once calibration was completed, resting values for HR, SV, and CO were obtained after which the warm-up began. During exercise, HR, SV, and CO values were determined every 15 s. Maximal values of HR, SV, and CO were identified as the highest values at any point during exercise. Maximal arteriovenous oxygen difference (a-vDO₂) was calculated as the quotient of $\dot{V}O_{2\max}$ (mL·min⁻¹)/CO_{max} (L·min⁻¹) and expressed in mL·dL⁻¹.

Incorporation of HIIT. At least 48 h after baseline testing, all participants initiated 10 sessions of progressive low-volume HIIT on the same electrically braked cycle ergometer consisting of 8–10 min of training time per day (Table 2). Participants were randomized using a Latin Squares design (13) to one of three regimes for the remaining 10 sessions. SIT (HIIT + SIT) consisted of 8–12 “all-out” sprints (4–6 min training duration per day) during which participants were required to pedal maximally. High-volume interval training (HIIT + HIIT_{HI}) required repeated 2.5-min bouts of cycling with 60 s recovery, leading to training duration equal to 12.5–17.5 min·d⁻¹. Lastly, the periodized regime (HIIT + PER) consisted of 3 sessions of high-volume HIIT, 3 sessions of SIT, and 4 sessions of low-volume HIIT during which training duration varied from 5 to 15 min per session (Table 2).

Before all sessions, a 5-min warm-up at 20% PPO was completed, and active recovery was performed at this work rate between bouts. During the initial 10 sessions, overload was incorporated after every three sessions by incorporating an additional one to two sessions and increasing work rate by 10% PPO. Participants were given additional recovery when requested. All exercise training was supervised, performed in a climate-controlled laboratory (temperature and relative humidity = 19°C–22°C and 30%–50%, respectively), and held at the same time of day (±1 h) within participants. During HIIT, HR was determined using telemetry (Polar Electro, Woodbury, NY) to identify peak (mean score of all values recorded at the end

TABLE 2. Description of 20 sessions of HIIT performed in the study.

Training Regimes	No. Bouts	Bout Duration (s)	Intensity (% PPO)	Recovery (s)	Session Duration (min)
HIIT + SIT					
1–3 ^a	8	60	90	75	23.0
4–6	9	60	100	75	25.25
7–10	10	60	110	75	27.5
11–13	8	30	130	120	25.0
14–16	10	30	140	120	30.0
17–20	12	30	150	120	35.0
HIIT + HIIT _{HI}					
1–3	8	60	90	75	23.0
4–6	9	60	100	75	25.25
7–10	10	60	110	75	27.5
11–13	5	150	70	60	22.5
14–16	6	150	75	60	26.0
17–20	7	150	80	60	29.5
HIIT + PER					
1–3	8	60	90	75	23.0
4–6	9	60	100	75	25.25
7–10	10	60	110	75	27.5
11–13	6	150	70	60	26.0
14–16	10	30	140	120	30.0
17–20	8	60	100	75	23.0

^aSpecific days of HIIT sessions. $\dot{V}O_{2\max}$ was determined after session 10 with PPO from this bout used to set intensity of sessions 11–20.

of each bout) and session HR, and strong verbal encouragement was provided. Average work rate (W) which included the HIIT bouts as well as recovery was also recorded using the Velotron software. Training was performed 3 d·wk⁻¹ and at least 24 h was provided between sessions, which were held on Monday/Wednesday/Friday, Monday/Tuesday/Thursday, and Tuesday/Thursday/Friday.

Habitual physical activity. Participants were advised to maintain their habitual physical activity during the study. They were given a training log upon study initiation and were required to denote daily physical activity in this log, which was submitted at study termination.

Data analyses. Data are expressed as mean ± SD and were analyzed using SPSS Version 22.0 (Chicago, IL). The Shapiro–Wilks test was used to assess normality. One-way ANOVA was performed to identify differences in demographic and physiological variables across all groups at baseline. Repeated three-way ANOVA (time = pre versus post, gender, and regimen = HIIT + SIT, HIIT + HIIT_{HI}, HIIT + PER, and CON) with repeated-measures was performed to identify differences in $\dot{V}O_{2\max}$ and maximal SV, CO, and $a\text{-vDO}_2$ during the study. If a significant F ratio was obtained, Tukey's *post hoc* test was used to identify differences between means. Partial eta-squared (η^2_p) and Cohen's *d* were used as an estimate of effect size. Pearson product moment correlation was used to determine the association between variables. Ninety-five percent confidence intervals (95% CI) are reported for the change in $\dot{V}O_{2\max}$ in response to HIIT. At a statistical power equal to 0.80 and expected difference in absolute $\dot{V}O_{2\max}$ from pretraining to posttraining equal to 15%, the minimum total sample size was determined to be 20 individuals (15). Using similar parameters other than a difference in $\dot{V}O_{2\max}$ response across regimens equal to 5%, a minimum of nine participants per regimen was determined which is similar to a previous study (26). Statistical significance was established as $P < 0.05$.

RESULTS

Training fidelity. Training compliance was equal to 778 of 780 (99.7% of all sessions). To verify the intensity of




FIGURE 1—Change in HR (mean ± SD) in response to 20 sessions of HIIT. * $P < 0.05$ versus sessions. HIIT is separated into phase 1 (sessions 1–3), phase 2 (sessions 4–6), phase 3 (sessions 7–10), phase 4 (sessions 11–13), phase 5 (sessions 14–16), and phase 6 (sessions 17–20).

HIIT (37), average peak HR in response to training is shown in Figure 1. HR increased during training ($P < 0.001$, $\eta^2_p = 0.25$) but there was no time–regimen interaction ($P = 0.41$). HR significantly increased ($P < 0.05$) from sessions 1–3, 4–6, and 7–10, when it peaked at $91.0\% \pm 2.6\%$ of baseline HR_{max}. HR declined ($P < 0.05$) in sessions 11–13 and was similar during the last seven sessions of training compared with values from sessions 1–10. Average work rate during HIIT increased across time ($P < 0.001$, $\eta^2_p = 0.36$) and a significant time–regimen interaction ($P < 0.001$, $\eta^2_p = 0.43$) was shown. During the first 10 sessions of HIIT, average work rate increased every three sessions of HIIT although it was similar ($P > 0.05$) across regimes (115–130 W). During sessions 11–20, participants in HIIT + HIIT_{HI} were exercising at significantly higher ($P < 0.05$) average session power outputs (127–143 W equal to 52.9%–59.6% PPO) versus HIIT + SIT (107–112 W equal to 47.3%–49.6% PPO) and HIIT + PER (109–124 W equal to 47.6%–54.6% PPO). Body mass did not change during the study (data not reported).

Differences in $\dot{V}O_{2\max}$, gas exchange variables, and PPO in response to HIIT. At baseline, there were no differences ($P > 0.05$) in any variable between groups, although differences ($P < 0.05$) between men and women occurred in $\dot{V}O_{2\max}$, $\dot{V}CO_2$, PPO, and V_E . Relative and absolute $\dot{V}O_{2\max}$ showed a significant main effect for time ($P < 0.001$, $\eta^2_p = 0.62$) as well as time–regimen interaction ($P < 0.001$, $\eta^2_p = 0.39$ and $P < 0.001$, $P < 0.001$, $\eta^2_p = 0.51$), as increases in $\dot{V}O_{2\max}$ were shown in all HIIT regimes but not in CON (Table 3). Individual data for change in absolute $\dot{V}O_{2\max}$ across all HIIT regimes are shown in Figure 2.

From baseline to posttraining, mean percent change in $\dot{V}O_{2\max}$ was equal to 10.9 ± 7.6 (95% CI, 6.5%–15.3%), $8.9\% \pm 6.1\%$ (95% CI, 5.2%–12.6%), and $12.3\% \pm 6.0\%$ (95% CI, 8.3%–16.3%) for HIIT + SIT, HIIT + HIIT_{HI}, and HIIT + PER, respectively, which were all higher ($P < 0.001$) than the change in CON ($-0.4\% \pm 6.6\%$; 95% CI, -2.3% to 2.0%). $\dot{V}O_{2\max}$ determined from the verification test was also increased ($P < 0.001$) in response to HIIT + SIT (39.3 ± 5.5 mL·kg⁻¹·min⁻¹ vs 42.3 ± 6.4 mL·kg⁻¹·min⁻¹, +7.7%), HIIT + HIIT_{HI} (40.2 ± 4.4 mL·kg⁻¹·min⁻¹ vs 43.7 ± 4.9 mL·kg⁻¹·min⁻¹, +9.0%), and HIIT + PER (39.1 ± 4.9 mL·kg⁻¹·min⁻¹ vs 42.2 ± 5.2 mL·kg⁻¹·min⁻¹, +8.0%), with no change demonstrated in CON (39.7 ± 5.0 mL·kg⁻¹·min⁻¹ vs 39.0 ± 5.4 mL·kg⁻¹·min⁻¹).

Post hoc analyses showed that $\dot{V}O_{2\max}$ in HIIT + HIIT_{HI} increased ($P < 0.05$) at 10 sessions of training (Cohen's $d = 1.68$) versus baseline after which it was unchanged (Cohen's $d = 0.50$) (Table 3). In contrast, the other regimes showed significant and sustained increases in $\dot{V}O_{2\max}$. In the first 10 sessions of HIIT + SIT, $\dot{V}O_{2\max}$ increased by 6.1% (Cohen's $d = 1.49$) and by an additional 3.5% in the last 10 sessions of training (Cohen's $d = 0.87$). Similarly, $\dot{V}O_{2\max}$ increased ($P < 0.05$) by 6.3% in the initial 10 sessions (Cohen's $d = 1.68$) and subsequently ($P < 0.05$) by 5% (Cohen's $d = 1.31$) in the latter 10 sessions of HIIT + PER.

TABLE 3. Maximal exercise responses (mean \pm SD) for power output, $\dot{V}O_{2\max}$, HR, SV, CO, and a-vDO₂ in response to 10 and 20 sessions of HIIT.

Regime	PPO (W)	$\dot{V}O_{2\max}$ (mL·kg ⁻¹ ·min ⁻¹)	$\dot{V}O_{2\max}$ (L·min ⁻¹)	HR (bpm)	SV (mL)	CO (L·min ⁻¹)	a-vDO ₂ (mL·dL ⁻¹)
HIIT + SIT							
Pre	230 \pm 35	39.8 \pm 7.3	2.7 \pm 0.6	186 \pm 9	104 \pm 16	19.2 \pm 2.6	14.3 \pm 2.1
Mid	242 \pm 36*	42.2 \pm 6.8*	2.9 \pm 0.5*	187 \pm 9	108 \pm 12	20.2 \pm 2.1*	14.5 \pm 2.0
Post	252 \pm 39***	43.6 \pm 6.1***	3.0 \pm 0.5***	189 \pm 7	113 \pm 10***	21.3 \pm 1.7***	14.1 \pm 1.9
HIIT + HIIT _{HI}							
Pre	240 \pm 34	41.1 \pm 4.9	2.9 \pm 0.6	185 \pm 9	112 \pm 19	20.6 \pm 3.0	14.1 \pm 1.6
Mid	262 \pm 43*	43.8 \pm 6.0*	3.1 \pm 0.7*	189 \pm 9	117 \pm 16*	21.8 \pm 2.9*	14.4 \pm 1.7
Post	265 \pm 47*	44.6 \pm 7.0*	3.1 \pm 0.8*	186 \pm 8	120 \pm 18*	22.2 \pm 3.1*	14.3 \pm 2.0
HIIT + PER							
Pre	229 \pm 56	39.5 \pm 5.6	2.7 \pm 0.8	189 \pm 8	107 \pm 19	20.3 \pm 3.7	13.3 \pm 2.5
Mid	245 \pm 50*	42.0 \pm 5.1*	2.8 \pm 0.8*	186 \pm 11	111 \pm 24	21.1 \pm 4.7	13.6 \pm 2.7
Post	249 \pm 53*	44.1 \pm 5.4***	3.0 \pm 0.9***	189 \pm 7	115 \pm 21*	21.8 \pm 4.4*	13.8 \pm 2.4
CON							
Pre	243 \pm 40	40.4 \pm 5.1	2.9 \pm 0.6	184 \pm 10	112 \pm 16	20.6 \pm 3.0	14.0 \pm 2.0
Post	244 \pm 41	40.2 \pm 5.5	2.9 \pm 0.6	184 \pm 9	112 \pm 15	20.5 \pm 2.9	14.0 \pm 1.9

* $P < 0.05$ from pretraining value.** $P < 0.05$ from value after session 10.

Significant effects of time ($P < 0.001$) and a time–regimen interaction ($P < 0.001$, $\eta^2_p = 0.35$) were shown for V_E (data not reported). Similar differences across time and a time–regimen interaction were found for $\dot{V}CO_2$ ($P < 0.001$, data not reported). There was no change in RER in response to HIIT ($P = 0.79$). PPO increased ($P < 0.001$) in all HIIT regimes by 20–25 W with no change seen in CON (Table 3).

Relationship between baseline $\dot{V}O_{2\max}$ and change in $\dot{V}O_{2\max}$ in response to HIIT. There was a significant inverse association ($r = -0.36$, $P = 0.01$) between baseline $\dot{V}O_{2\max}$ and the training-mediated change in $\dot{V}O_{2\max}$, suggesting that a low baseline $\dot{V}O_{2\max}$ is related to a greater change in $\dot{V}O_{2\max}$ with training.

Differences in hemodynamic variables in response to HIIT. At baseline, there were no differences in any variable between groups, although men showed higher ($P < 0.05$) SV, CO, and a-vDO₂ versus women. Data were combined across men and women because there was no time–gender interaction. HR did not change in response to training ($P = 0.88$). Cardiac output increased across time ($P < 0.001$, $\eta^2_p = 0.36$) and a significant time–regimen interaction occurred ($P = 0.002$, $\eta^2_p = 0.25$) in that it increased in response to training and did not change in CON (Table 3). *Post hoc* analyses showed that CO in HIIT + SIT increased ($P < 0.05$) by 5.3% (Cohen's $d = 0.91$) and then an additional 5.4% (Cohen's $d = 1.0$) after 10 and 20 sessions of training. Nevertheless, in HIIT + HIIT_{HI}, CO increased by 6% from 10 sessions of training (Cohen's $d = 1.1$) after which it was maintained (Cohen's $d = 0.37$). In HIIT + PER, it was similar after 10 sessions (Cohen's $d = 0.73$) and was higher ($P < 0.05$) after 20 sessions of training (7.5%, Cohen's $d = 1.37$) (Table 3). Compared with baseline, SV increased ($P < 0.05$ between session 20 and preceding values) in HIIT + SIT and increased ($P < 0.05$) after 10 and 20 sessions of training for HIIT + HIIT_{HI} and after 20 sessions for HIIT + PER, respectively. SV did not change in CON. Individual changes in SV and CO are shown in Figure 3. Results showed no difference in a-vDO₂ in response to training ($P = 0.36$) (Table 3).

DISCUSSION

The present study is the first to examine the effects of 6 wk of HIIT characterized by a modification of intensity, duration, and overall structure *during* training on changes in cardiac output and $\dot{V}O_{2\max}$ in active men and women. This approach models what exercisers frequently do on a day-to-day basis yet is rarely used in research. In all participants, the initial 10 sessions of HIIT led to similar and significant increases in $\dot{V}O_{2\max}$, but the subsequent 10 sessions varying in intensity and volume did not lead to continued increases in $\dot{V}O_{2\max}$ in all regimes. Our findings refute previous studies using both SIT and HIIT (22,25) as we show that significant increases in SV and CO, but not a-vDO₂, occur in response to HIIT which are consequent with increases in $\dot{V}O_{2\max}$. The overall magnitude of increase in $\dot{V}O_{2\max}$ was similar across regimes, although the time course of change differed across groups, and the increase in $\dot{V}O_{2\max}$ was inversely related to baseline cardiorespiratory fitness.

Our data showing similar increases in $\dot{V}O_{2\max}$ across different HIIT regimes varying in intensity and structure support data from recent studies in habitually active individuals. Similar increases in $\dot{V}O_{2\max}$ were demonstrated in men performing 4 wk of repeated 15 or 30 s bouts of SIT (39) as well as in men and women performing six sessions of repeated 30 or 10 s Wingate tests (18). However, in sedentary individuals (age and $\dot{V}O_{2\max} = 41$ yr and 33.9 mL·kg⁻¹·min⁻¹), 6 wk of high-volume HIIT consisting of repeated 4 min bouts led to significantly greater increases in $\dot{V}O_{2\max}$ compared with low-volume HIIT (ten 1 min bouts) or MICT (7). These data, however, are opposed by similar increases in $\dot{V}O_{2\max}$ found in sedentary, young, nonobese women who completed 12 wk of moderate or more intense HIIT (5) as well as in sedentary men completing HIIT (repeated 3 min efforts at 85%–90% $\dot{V}O_{2\max}$) and SIT (repeated 30 s efforts at 120% $\dot{V}O_{2\max}$) (26). Discrepancies between studies include the markedly higher training volume used in the Baekkerud et al. (7) study, which has been identified (6) to elicit more substantial increases in $\dot{V}O_{2\max}$ in response to HIIT, as well as heterogeneity in $\dot{V}O_{2\max}$




FIGURE 2—Individual changes in $\dot{V}O_{2\max}$ ($L \cdot \min^{-1}$) in response to 20 sessions of (A) HIIT + SIT, (B) HIIT + HIIT_{HI}, and (C) HIIT + PER. 0 = pre-training value, 10 = value recorded after initial 10 training sessions, and 20 = posttraining value. Thin lines are individual participants and thick line is the mean change in $\dot{V}O_{2\max}$ across participants within group.

response to HIIT (4). This individual response to HIIT (Fig. 2) is also exhibited by 43% of participants showing greater than the mean change in $\dot{V}O_{2\max}$ (10.7%) in response to HIIT, 46% showing less than this change, and 10% showing no change (less than the typical error for $\dot{V}O_{2\max}$). Increases in $\dot{V}O_{2\max}$ are inversely associated with baseline $\dot{V}O_{2\max}$ after HIIT (4) or SIT (17), although this relationship typically does not occur after prolonged MICT in untrained participants (8,34). Our data also showed that in 20 participants with $\dot{V}O_{2\max}$ below $40 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$, the mean value of our sample, 55% exhibited greater than average increases in $\dot{V}O_{2\max}$ in response to HIIT, which suggests that a portion of

the training-induced increase in cardiorespiratory fitness is associated with baseline values.

An intriguing finding of the present study is that an additional 10 sessions of HIIT_{HI} did not further augment $\dot{V}O_{2\max}$ compared with SIT or PER (Table 3). This is unexpected considering that this regime is characterized by higher training duration (12.5–17.5 min per session) and mean power output (see Results) which seem to elicit greater increases in $\dot{V}O_{2\max}$ (6,7) compared with SIT (4–6 min per session) or PER (5–15 min per session). In contrast, Hazell et al. (18) denoted that it is the peak intensity achieved during SIT that mediates the resultant change in $\dot{V}O_{2\max}$ rather than training volume. Similar findings from highly trained cyclists show that completion of repeated 30-s intervals allowing greater time above 90% $\dot{V}O_{2\max}$ led to greater increases in $\dot{V}O_{2\max}$ compared with longer intervals (5 min) at lower intensity (33). It is plausible that after the initial 10 sessions of training, the novel use of SIT and PER, which dramatically modifies the HIIT regime by including brief efforts at supramaximal intensities, promotes continued improvements in $\dot{V}O_{2\max}$ which do not happen when primarily training volume but not intensity was enhanced in HIIT_{HI}.

Our findings show a significant increase in maximal CO in response to all HIIT regimes ranging from 1.5 to 2.1 $L \cdot \min^{-1}$ (8%–11%) (Table 3). However, the time course of increase in CO varied as it steadily increased in HIIT + SIT, plateaued after 10 sessions of training in HIIT + HIIT_{HI}, and did not initially change in HIIT + PER after which it increased. Although speculative, the slightly different SV response in each group may have led to this result. In untrained adults (12) (age and $\dot{V}O_{2\max} = 47$ yr and $26 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$), CO_{\max} was increased by 12% in response to 24 sessions of HIIT. However, in this study, HIIT consisted of repeated 5-min bouts including 4 min at work rate coincident with lactate threshold and 1 min at 90% PPO, so this regimen was higher volume and lower intensity compared with our HIIT protocols. In untrained men (age and $\dot{V}O_{2\max} = 30$ yr and $38 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) (38), 36 sessions of HIIT (repeated 2 min efforts at 90% $\dot{V}O_{2\max}$) led to significant increases in $\dot{V}O_{2\max}$ which was associated with increased CO_{\max} but not a-v DO_2 . In these studies, an increase ($P < 0.05$) in maximal SV led to the increase in CO, supporting our results (Table 3), although maximal HR was also increased in the Daussin et al. (12) study. One likely mechanism explaining this increase in SV is a significant increase in plasma volume (PV) via HIIT, which is seen after only three and six sessions of training (38) and elicits an increase in blood volume (BV) and thus SV. Only six sessions of HIIT at 95%–100% $\dot{V}O_{2\max}$ also promoted significant increases in PV (10.8%) leading to an 11% increase in $\dot{V}O_{2\max}$ in untrained men (age and $\dot{V}O_{2\max} = 25$ yr and $39 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) (14). In young active adults ($\dot{V}O_{2\max} = 46.8 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$), completing 6 wk of maximal treadmill-based SIT, there was no change in maximal CO and the increase in $\dot{V}O_{2\max}$ was due to a greater a-v DO_2 (25). Similarly, six sessions of HIIT at 100% PPO did not change maximal PV, BV, or CO despite a significant




FIGURE 3—Individual changes in SV (mL) in response to 20 sessions of (A) HIIT + SIT, (B) HIIT + HIIT_{HI}, and (C) HIIT + PER. 0 = pre-training value, 10 = value recorded after initial 10 training sessions, and 20 = posttraining value. Thin lines are individual participants and thick line is the mean change in SV across participants within group. Individual changes in cardiac output (L·min⁻¹) in response to 20 sessions of (D) HIIT + SIT, (E) HIIT + HIIT_{HI}, and (F) HIIT + PER. 0 = pretraining value, 10 = value recorded after initial 10 training sessions, and 20 = posttraining value. Individual participants (*thin lines*). Mean change in cardiac output across participants within group (*thick line*).

8% increase in $\dot{V}O_{2max}$ (22), which led the authors to conclude that peripheral adaptations increase $\dot{V}O_{2max}$ after short-term HIIT. One characteristic of both studies is that participants' $\dot{V}O_{2max}$ was 15%–20% higher compared with studies in which CO was enhanced with HIIT (12,38). It may be that persons with lower baseline $\dot{V}O_{2max}$ are more apt to experience hypervolemia soon after initiation of HIIT leading to rapid increases in BV, SV, and thus CO. In addition, it is plausible that participants with higher $\dot{V}O_{2max}$ such as used in previous studies (22,25) require higher training volumes or duration of HIIT/SIT to significantly increase SV and CO. This discrepancy requires additional study to ascertain if there is a specific $\dot{V}O_{2max}$ value or training dose for which central rather than peripheral adaptations mediate the increased $\dot{V}O_{2max}$ demonstrated in response to HIIT.

In the current study, we used thoracic impedance to estimate SV and CO during progressive exercise to $\dot{V}O_{2max}$.

This technique was also used in another study (12), showing significant increases in SV and CO in response to HIIT. Previous data show an intraclass correlation coefficient equal to 0.95 for repeated determinations of maximal CO obtained in habitually active men and women (2), and data from our control participants showed low typical error equal to 0.5 L·min⁻¹ for two determinations of CO conducted 6 wk apart. In a previous study (24), this technique was able to detect small differences in CO across different protocols leading to $\dot{V}O_{2max}$.

Our study is strengthened by precise allocation of intensity for all training sessions as well as high compliance to training. In addition, the fidelity of HIIT was demonstrated (Fig. 1) by showing that all regimes elicited intensities approaching 90% HR_{max}. Results from the verification test and maximal HR/RER data suggest that participants did exhibit “true” $\dot{V}O_{2max}$ and that our reported increases in $\dot{V}O_{2max}$ are repeatable and

not due to random error. We also recruited both men and women to apply our data to a broader population of active adults. Nevertheless, there is some evidence (16) that women may respond differently to HIIT versus men, although this is not a universal finding (1). In addition, accelerometry was not used to monitor habitual physical activity completed outside the study, although data from exercise training logs showed no difference in hours per week of physical activity (data not reported). We did not include a group of individuals who completed endurance training, so we cannot state if any of the HIIT regimes elicit superior increases in $\dot{V}O_{2\max}$ compared with traditional MICT. Although control participants had similar physical activity, body fat, and $\dot{V}O_{2\max}$ versus those who completed HIIT, they were recruited by convenience rather than through random assignment. Assessments of $\dot{V}O_{2\max}$ and CO were made 2–5 d after the previous HIIT session, so slight reductions in PV could have occurred (11), leading to small underestimations in CO and consequently $\dot{V}O_{2\max}$. As an equal number of participants in each regimen completed their assessments 4–5 d after their final HIIT session, the effect of this discrepancy on our results is likely small.

REFERENCES

- Astorino TA, Allen RP, Roberson DW, et al. Adaptations to high-intensity training are independent of gender. *Eur J Appl Physiol*. 2011;111(7):1279–86.
- Astorino TA, Bovee C, DeBoe A. Estimating hemodynamic responses to the Wingate test using thoracic impedance. *J Sports Sci Med*. 2015;14(4):834–40.
- Astorino TA, Edmunds RM, Clark A, et al. Increased cardiac output and maximal oxygen uptake in response to 10 sessions of high intensity interval training. *J Sports Med Phys Fit*. 2016 (in press).
- Astorino TA, Schubert MM. Individual responses to completion of short-term and chronic interval training: a retrospective study. *PLoS One*. 2014;9(5):e97638.
- Astorino TA, Schubert MM, Palumbo E, et al. Magnitude and time course of changes in maximal oxygen uptake in response to distinct regimens of chronic interval training in sedentary women. *Eur J Appl Physiol*. 2013;113(9):2361–9.
- Bacon AP, Carter RE, Ogle EA, Joyner MJ. $\dot{V}O_{2\max}$ trainability and high intensity interval training in humans: A meta-analysis. *PLoS One*. 2013;8(9):e73182.
- Bækkerud FH, Solberg F, Leinan IM, et al. Comparison of three popular exercise modalities on $\dot{V}O_{2\max}$ in overweight and obese. *Med Sci Sports Exerc*. 2015;48(3):491–8.
- Bouchard C, An P, Rice T, et al. Familial aggregation of $\dot{V}O_{2\max}$ response to exercise training: results from the HERITAGE Family Study. *J Appl Physiol (1985)*. 1999;87:1003–8.
- Burgomaster KA, Howarth KR, Phillips SM, et al. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *J Physiol*. 2008;586(1):151–60.
- Charloux A, Lonsdorfer-Wolf E, Richard R, et al. A new impedance cardiographic device for the non-invasive evaluation of cardiac output at rest and during exercise: a comparison with the “direct” Fick method. *Eur J Appl Physiol*. 2000;82:313–20.
- Coyle EF, Hemmert MK, Coggan AR. Effects of detraining on cardiovascular responses to exercise: role of blood volume. *J Appl Physiol (1985)*. 1986;60(1):95–9.
- Daussin FN, Ponsot E, Dufour SP, et al. Improvement of $\dot{V}O_{2\max}$ by cardiac output and oxygen extraction adaptation during intermittent versus continuous endurance training. *Eur J Appl Physiol*. 2007;101:377–83.
- Devillers A, Hall JI. Rank 3 latin square designs. *J Combinatorial Theory Series*. 2006;113(5):894–902.
- Esfandiari S, Sasson Z, Goodman JM. Short-term high-intensity interval and continuous moderate-intensity training improve maximal aerobic power and diastolic filling during exercise. *Eur J Appl Physiol*. 2014;114(2):331–43.
- Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavioral Res Meth*. 2007;39(2):175–91.
- Gillen JB, Percival ME, Skelly LE, et al. Three minutes of all-out intermittent exercise per week increases skeletal muscle oxidative capacity and improves cardiometabolic health. *PLoS One*. 2014; 9(11):e111489.
- Gurd BJ, Giles MD, Bonafiglia JT, et al. Incidence of nonresponse and individual patterns of response following sprint interval training. *Appl Physiol Nutr Metab*. 2016;41(3):229–34.
- Hazell TJ, MacPherson RE, Gravelle BM, Lemon PW. 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol*. 2010;110:153–60.
- Heyward VH, Gibson AL. *Advanced Fitness Assessment and Exercise Prescription*. 7th ed. Champaign (IL): Human Kinetics; 2014. pp. 219–64.
- Jackson AS, Pollock ML. Generalized equations for predicting body density of men. *Br J Nutr*. 1978;40:497–504.
- Jackson AS, Pollock ML, Ward A. Generalized equations for predicting body density of women. *Med Sci Sports Exerc*. 1980; 12(3):175–81.
- Jacobs RA, Flück D, Bonne TC, et al. Improvements in exercise performance with high-intensity interval training coincide with an increase in skeletal muscle mitochondrial content and function. *J Appl Physiol (1985)*. 2013;115:785–93.
- Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*. 2009;301(9):2024–35.

CONCLUSIONS

Overall, in young individuals with average cardiorespiratory fitness, 20 sessions of periodized HIIT led to significant increases in $\dot{V}O_{2\max}$ which were attendant with increases in maximal SV and CO. Data suggest that increases in $\dot{V}O_{2\max}$ as a result of HIIT are mediated by improvements in central O_2 delivery rather than peripheral adaptations as frequently reported (9,22,25).

The authors thank the participants for their dedication to the study. In addition, we extend gratitude to Kayla Snow, Christian Ridings, Jason Bandong, Brendyn Heath, Gina Ordille, Jordan Ridings, Matt Montell, and Angelina Mata-Villa for assistance with data collection. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation.

The authors declare no conflict of interest in completion of this study, and no funding was acquired to pursue this study. The results of this study do not constitute endorsement by ACSM.

No funding was obtained to support this work, and the authors declare that they have no conflicts of interest in executing this study.

24. Lepretre PM, Koralsztein JP, Billat VL. Effect of exercise intensity on relationship between $\dot{V}O_{2\max}$ and cardiac output. *Med Sci Sports Exerc.* 2004;36(8):1357–63.
25. MacPherson REK, Hazell TJ, Olver TD, Paterson DH, Lemon PWR. Run sprint interval training improves aerobic performance but not maximal cardiac output. *Med Sci Sports Exerc.* 2011;43(1):115–22.
26. Matsuo T, Saotome K, Seino S, et al. Effects of a low-volume aerobic-type interval exercise on $\dot{V}O_{2\max}$ and cardiac mass. *Med Sci Sports Exerc.* 2014;46(1):42–50.
27. Midgley AW, McNaughton LR, Carroll S. Verification phase as a useful tool in the determination of the maximal oxygen uptake of distance runners. *Appl Physiol Nutr Metab.* 2006;31:541–8.
28. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med.* 2002;346(11):793–801.
29. National Cancer Institute, *Past Year Total Physical Activity Questionnaire*. Bethesda, MD: National Institutes of Health; 2013, pp. 1–2.
30. Nybo L, Sundstrup E, Jakobsen MD, et al. High-intensity training versus traditional exercise interventions for promoting health. *Med Sci Sports Exerc.* 2010;42(10):1951–8.
31. Poole DC, Gaesser GA. Response of ventilatory and lactate thresholds to continuous and interval training. *J Appl Physiol (1985).* 1985;58(4):1115–21.
32. Richard R, Lonsdorfer-Wolf E, Charloux A, et al. Non-invasive cardiac output evaluation during a maximal progressive exercise test, using a new impedance cardiograph device. *Eur J Appl Physiol.* 2001;85:202–7.
33. Rønnestad BR, Hansen J, Vegge G, Tønnessen O, Slettalokken G. Short intervals induce superior training adaptations compared with long intervals in cyclists - an effort-matched approach. *Scand J Med Sci Sports.* 2015;25:143–51.
34. Scharhag-Rosenberger F, Walitzek S, Kindermann W, Meyer T. Differences in adaptations to 1 year of aerobic endurance training: individual patterns of nonresponse. *Scand J Med Sci Sports.* 2012;22:113–8.
35. Sijie T, Hainai Y, Fengying Y, JianXiong W. High intensity interval exercise training in overweight young women. *J Sports Med Phys Fitness.* 2012;52:255–62.
36. Strohacker K, Fazzino D, Breslin WL, Xu X. The use of periodization in exercise prescriptions for inactive adults: a systematic review. *Prev Med Rep.* 2015;2:385–96.
37. Taylor KL, Weston M, Batterham AM. Evaluating intervention fidelity: an example from a high-intensity interval training study. *PLoS One.* 2015;10(4):e0125166.
38. Warburton DER, Haykowsky MJ, Quinney HA, et al. Blood volume expansion and cardiorespiratory function: effects of training modality. *Med Sci Sports Exerc.* 2004;36(6):991–1000.
39. Zelt JG, Hankinson PB, Foster WS, et al. Reducing the volume of sprint interval training does not diminish maximal and submaximal performance gains in healthy men. *Eur J Appl Physiol.* 2014;114(11):2427–36.