

5-ASA or Mesalamine (Rectal)

Scan for more!



MEZERA® foam or suppositories, PENTASA® enema or suppositories, SALOFALK® enema or suppositories



What this medication is used for

Widely used for children with Crohn's disease or ulcerative colitis.



Health Canada also approved for Crohn's disease and ulcerative colitis in adults.



How long to use this medication

This medication will be used to get your IBD under control and keep it under control long-term.



How this medication works

5-ASA work locally to reduce inflammation in the intestine.

There are a number of 5-ASA medications which differ based on route of administration and medication coating.



How to use this medication

Use this medication at night before going to bed. Rectal doses should be held for at least 30 minutes, but it could take time and practice before be able to do so. If possible, hold the dose all night.

-Although some individuals receive benefit quite quickly, it can take a little longer in others. Keep taking your medication.



Monitoring during treatment

Regular blood tests, stool tests, endoscopy and/or other imaging will be done to monitor for response and side-effects to medication.



Side Effects

Abdominal pain, diarrhea, nausea or headaches can occur.

-Inflammation of the pancreas, lungs, liver, kidneys and heart may very rarely occur.



When I should call my doctor

Please call if:

-You want to or have already stopped the medicine

-If you are concerned about any side effects

-Worsening IBD symptoms such as blood in the stool, diarrhea and/or abdominal pain

-If you become pregnant



Where treatment is administered

Fill your prescription at your pharmacy.



Vaccines

✓ Live vaccines
- Safe

✓ Inactivated vaccines
- Safe