



Review article

Critical review on the socio-economic impact of tendinopathy

Chelsea Hopkins^{a,b}, Sai-Chuen Fu^{a,b}, Eldrich Chua^{a,b}, Xiaorui Hu^{a,b}, Christer Rolf^{b,c},
Ville M. Mattila^{c,d}, Ling Qin^{a,e,f}, Patrick Shu-Hang Yung^{a,b}, Kai-Ming Chan^{a,b,*}

^a Department of Orthopaedics and Traumatology, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong SAR, China

^b Lui Che Woo Institute of Innovative Medicine, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong SAR, China

^c Division of Orthopaedics and Biotechnology, Department of Clinical Science, Intervention and Technology, Karolinska Institute, Stockholm, Sweden

^d Department of Orthopaedics, Tampere University Hospital, Tampere, Finland

^e Musculoskeletal Research Laboratory, Department of Orthopaedics and Traumatology, The Chinese University of Hong Kong, Hong Kong SAR, China

^f Translational Medicine Research and Development Centre, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China

Received 30 October 2015; revised 21 January 2016; accepted 21 January 2016

Available online 22 April 2016

Abstract

There are currently no studies that determine the total burden that tendinopathy places on patients and society. A systematic search was conducted to understand the impact of tendinopathy. It demonstrated that the current prevalence is underestimated, particularly in active populations, such as athletes and workers. Search results demonstrate that due to the high prevalence, impact on patients' daily lives and the economic impact due to work-loss, treatments are significantly higher than currently observed. A well-accepted definition by medical professionals and the public will improve documentation and increase awareness, in order to better tackle the disease burden.

Copyright © 2016, Asia Pacific Knee, Arthroscopy and Sports Medicine Society. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: incidence; prevalence; quality-of-life; socio-economic; tendinopathy

Introduction

Clinicians obtain insight into the burden of tendinopathy from their patients, yet their ability to alleviate this burden remains limited. From their observations, it is to be believed that tendinopathy has a significant socio-economic impact, but there is no direct evidence to support this claim. This review aims to determine the socio-economic burden of tendinopathy and how this burden may be alleviated. The definition and classification of tendinopathy currently adopted by medical subject headings are displayed in [Figure 1](#). Tendinopathy is a

blanket term for “tendinitis”, “tendinosis”, and “tenosynovitis”. “Tendinitis” was the original term to define pain and inflammation within the tendon, and “tendinosis” was the preferential term to describe the degenerative changes observed. Strictly speaking, “tenosynovitis” refers to inflammation of the synovial sheath surrounding the tendon, thus it should not be regarded as tendinopathy in which degenerative changes are mainly observed in the tendon itself. By contrast, spontaneous tendon rupture, which occurs without prior symptoms, is attributed to mechanical weakness of tendons due to tendinopathic changes.¹ In summary, tendinopathy is characterised by chronic tendon degeneration, resulting in pain and rupture, which are the basic criteria used when searching for relevant information.

The disease burden of tendinopathy can be primarily reflected by the number of patients, the effect on the patients' quality-of-life, cost effectiveness of treatments, and the

* Corresponding author. The Chinese University of Hong Kong, Department of Orthopaedics and Traumatology, Room 74029, 5F, Lui Che Woo Clinical Sciences Building, Prince of Wales Hospital, Shatin, New Territories, Hong Kong SAR, China.

E-mail address: kaimingchan@cuhk.edu.hk (K.-M. Chan).




Figure 1. Nomenclature, definitions, and types of tendon disorders.

economic implications of work disability. Therefore, we performed a systematic search of prevalence and incidence data of tendinopathy, and gathered information about quality-of-life, work disability, and treatments specific to tendinopathy.

Prevalence and incidence of tendinopathy

A literature search was performed in PubMed in October 2015 using the search strategy: (Tendinopathy OR tendinitis OR tendonitis OR tendinosis OR tendon rupture OR tendon tear OR jumper's knee OR Sinding-Larsen-Johansson OR epicondylitis OR tennis elbow) AND (prevalence OR incidence OR epidemiology). Studies are included if prevalence or incidence of tendinopathy was reported. Studies on tenosynovitis and traumatic injuries were excluded. Non-English studies, reviews, animal, and cadaveric studies were also excluded. The search returned 1819 articles, of which 132 were included based on the selection criteria. The search results were tabulated according to the nature of the cohort (athletes, workers, general population, and patients with comorbidities), sample size, age group, type of tendinopathy involved, and the reported prevalence and incidence data.

Of the cohorts identified, athletes formed the major cohort with 42 studies, followed by workers (36 studies), individuals in the general population (35 studies), and individuals with comorbidities (19 studies). Achilles' tendinopathy, patellar tendinopathy, epicondylitis, and rotator cuff tendinopathy are identified as four major types of tendinopathy according to numbers of studies and the reported prevalence. The results are shown in Table 1.

Results

Athletes

The high intensity and frequency of physical activities in athletes exposes this group to overuse injuries due to the high stress exerted on the tendons. Records of medical attendance in the 2004 Olympics² and 2007 Pan-American Games³ show that tendinopathy was within the top three most treated conditions in athletes. This record represents the significance of tendinopathy as a widespread condition in this group.

Studies on the prevalence of upper extremity tendinopathy in athletes have observed small cohorts, yet data from studies with the largest sample sizes place the prevalence for rotator cuff tendinopathy at 23.7% in volleyball players, and epicondylitis at 13.1% in climbers.^{4,5} Older age may also play a role as evidenced in a study on elderly athletes where prevalence was seen to be as high as 48.2%.⁶ There is no study on upper extremity tendinopathy in adolescents to our knowledge. A study on patellar tendinopathy reported a prevalence of approximately 17% in adults and 5.6% in adolescents.^{7–10} Similarly, Achilles' tendinopathy was reported to be 12.5% in adults and 7.8% in adolescents.^{11,12} Adolescents are seemingly less affected by tendinopathy based on these values alone. There is however no clear evidence that age influences tendinopathy.¹³ In agreement with previous studies,^{13,14} no clear trend is observed when comparing the prevalence or incidence between male and female athletes.

Lower extremity tendinopathy, particularly that of the patellar tendon, is the most frequently studied and arguably the most commonly affected. However, sports-related tendinopathy is challenging to generalise due to the difference in anatomical sites affected and the degree of exposure. For instance, dancers present with higher prevalence of Achilles' tendinopathy, while rowers would more frequently present with rotator cuff tendinopathy or epicondylitis. In addition, the degree of sport participation would differ widely between recreational athletes and professional athletes, but professional or elite athletes may suffer greater economic losses from injury as compared to recreational athletes. Studies on the degree of participation, the associated risk of tendinopathy development, and the associated impact would be valuable further studies.

Workers

Occupational exposure is of particular relevance because of the high economic impact procured by productivity-loss and compensation for disease. Highly repetitive movements are commonly observed in daily work tasks, and coupled with poor workplace ergonomics, workers are placed at an increased risk of developing tendinopathy. A distinction can be made between workers and athletes in that occupational exposure typically consists of relatively low demand and highly repetitive movements over a longer period of time compared to athletic exposure. Worker cohorts have generally been larger than the athlete cohort. Many of these cohorts have been merged from different workplaces and may possibly be highly heterogeneous even within the same study. Tendinopathy in workers is almost exclusively observed in the upper extremity. The most common and arguably most prevalent of which is lateral epicondylitis. A prevalence of 2–3% have been observed, but rates as high as 18% and 41% have also been reported in spine surgeons and coal miners, respectively.^{15–17} Similar to athletic exposures, it is evident that the type of work influences the prevalence of tendinopathy.¹⁷ Relative risk in occupational exposure with regards to frequency of repetitive motion, length of exposure, and ergonomic factors may be worthwhile studies.

Table 1
Prevalence and incidence of tendinopathy in different cohorts.

1 st Author, year, Ref	Group	Cohort	N	Age	Type of tendinopathy	Prevalence	Incidence
Zapata, 2006 ⁷¹	General	Students	791	Adolescent	Tendonitis	2	n/a
Salaffi, 2005 ²³	General	Italian general population	2155	Adults	LE	0.7	n/a
Miranda, 2005 ¹⁸	General	General population	8028	Adults	RC tendinitis	2	n/a
Rechardt, 2010 ⁷²	General	General population	6237	Adults	RC tendinitis	2.8	n/a
Tajika, 2014 ²⁴	General	Japanese mountain village community	422	Adults	LE	3.8	n/a
Joseph, 2012 ⁷³	General	Asymptomatic active university student body	52	Adults	AT (US)	3.8	n/a
Koplas, 2011 ²⁷	General	Elbow MRI examinations	801	Adults	Triceps tendon tear	3.8	n/a
Waldecker, 2012 ⁷⁴	General	Non-athletes in orthopaedic clinic	697	Adults	AT tendinopathy	5.6	n/a
Schibany, 2004 ⁷⁵	General	Asymptomatic patients	212	Adults	Supraspinatus rupture (US)	6	n/a
Zwerver, 2011 ³⁹	General	Nonelite athletes	891	Adults	Jumper's knee	8.5	n/a
Fairley, 2014 ³⁵	General	Community with no history of knee pain or injury	297	Adults	PT (MRI)	28.3	n/a
Walker-Bone, 2012 ⁴⁴	General	General population	6038	Adults	LE	0.7	n/a
Shiri, 2006 ²¹	General	General population	4783	Adults	ME	0.6	n/a
Alvarez-Nemegyei, 2011 ²⁸	General	General population	12,686	Adults	LE	1.3	n/a
Walker-Bone, 2004 ²²	General	General population	6038	Adults	ME	0.4	n/a
Shiri, 2007 ¹⁹	General	General population	6254	Adults	RC tendinopathy	2.4	n/a
Girish, 2011 ³⁶	General	Asymptomatic shoulders	51	Adults	Bicipital tendinopathy	0.3	n/a
Safran, 2002 ⁷⁶	General	General population	279,500	Adults	AT tendinopathy	0.1	n/a
Witvrouw, 2001 ⁵²	General	Students without knee conditions	138	Adults	RC tendinitis	3.3	n/a
Huttunen, 2014 ⁶⁹	General	Nationwide Sweden	27,702	Adults	Bicipital tendinitis	0.4	n/a
Ostor, 2005 ⁷⁷	General	General population	17,000	Adults	LE	0.7	n/a
Cretnik, 2010 ⁷⁸	General	General population	572,929	Adults	ME	0.6	n/a
Darmawan, 1995 ²⁵	General	Indonesian population	1118	All	RC tendinitis	3.8	n/a
Moller, 1996 ⁷⁹	General	Malmo population	n/a	All	Bicipital tendinitis	0.5	n/a
Clayton, 2008 ⁸⁰	General	General population	535,000	All	LE	1.1	n/a
Levi, 1997 ⁸¹	General	Copenhagen population	n/a	All	ME	0.3	n/a
Leppilahti, 1996 ⁸²	General	Oulu population	n/a	All	Supraspinatus (US)	39	n/a
Houshian, 1998 ⁸³	General	Danish county	220,000	All	Subscapularis (US)	25	n/a
Maffulli, 1999 ⁸⁴	General	General population	n/a	All	Supraspinatus tear (US)	22	n/a
van der Linden, 2001 ⁷⁰	General	General population	n/a	All	Biceps tendon rupture	n/a	1.2/100,000 PY
Suchak, 2005 ⁸⁵	General	Canada general population	967,200	All	PT tendinitis	n/a	13.8% (2 y CI)
Chard, 1987 ²⁰	General	Geriatric unit not admitted for shoulder complaints	100	Elderly	AT rupture	n/a	29.5/100,000 PY
Horowitz, 2013 ²⁶	General	General population	n/a	n/a	RC tendinopathy	n/a	8.1/1000 PY
de Jonge, 2011 ⁸⁶	General	General population	57,725	n/a	AT rupture	n/a	7.6/100,000 PY
					Elderly		1.3/100,000 PY
					Epicondylitis	6.6	n/a
					AT rupture	n/a	0.06 (4 y CI)
					AT rupture	n/a	11.3/100,000 PY
					AT rupture	n/a	13.4/100,000 PY
					AT rupture	n/a	18/100,000 PY
					AT rupture	n/a	37.3/100,000 PY
					AT rupture	n/a	6/100,000 PY
					Tendon rupture	n/a	6.32/100,000 PY
					AT rupture	n/a	8.3/100,000 PY
					RC tendinitis	5	n/a
					RC rupture	7	n/a
					Retropharyngeal calcific tendinitis	n/a	0.5/100,000 PY
					AT tendinopathy	n/a	1.8/1000 PY

(continued on next page)

Table 1 (continued)

1 st Author, year, Ref	Group	Cohort	N	Age	Type of tendinopathy	Prevalence	Incidence
Nyyssonen, 2008 ⁸⁷	General	Finnish population	5.2m	n/a	AT rupture	n/a	11.5/100,000 PY
McCormack, 1990 ⁸⁸	Worker	Textile workers	2047	Adults	Epicondylitis	2	n/a
Roquelaure, 2006 ¹⁵	Worker	Workers	2685	Adults	LE	2.4	n/a
Almeida, 2012 ⁸⁹	Worker	Workers	951	Adults	Tendinitis	3.2	n/a
Frost, 2002 ⁹⁰	Worker	Workers	782	Adults	Shoulder tendinitis	3.2	n/a
Descatha, 2003 ⁹¹	Worker	Workers	1757	Adults	ME	5.2	1.5% (annual CI)
Fan, 2009 ⁹²	Worker	Workers	733	Adults	LE	5.2	n/a
Rosenbaum 2013 ⁹³	Worker	Latino poultry workers	516	Adults	Epicondylitis	5.8	n/a
Kryger, 2007 ⁹⁴	Worker	Computer workers with neck or arm pain	1369	Adults	LE	5.8	n/a
Kaergaard, 2000 ⁹⁵	Worker	Sewing machine operators	243	Adults	RC tendinitis	5.8	n/a
Dimberg, 1987 ⁹⁶	Worker	Workers	540	Adults	LE	7.4	n/a
Roto, 1984 ⁹⁷	Worker	Male meat cutters	90	Adults	Epicondylitis	8.9	n/a
Ono, 1998 ⁹⁸	Worker	Nursery school cooks	209	Adults	Epicondylitis	11.5	n/a
Leclerc, 2001 ⁹⁹	Worker	Workers	598	Adults	LE	12.2	12.2% (3 y CI)
Capone, 2010 ¹⁰⁰	Worker	Plastic surgeons	339	Adults	Epicondylitis	13.5	n/a
Ritz, 1995 ¹⁰¹	Worker	Workers	290	Adults	Epicondylitis	14.1	n/a
Chiang, 1993 ¹⁰²	Worker	Workers in fish-processing	207	Adults	Epicondylitis	15	n/a
Barrero, 2012 ¹⁰³	Worker	Workers flower industry	158	Adults	Epicondylitis	15.2	n/a
Auerbach, 2011 ¹⁶	Worker	Spine surgeons	561	Adults	LE	18	n/a
Forde, 2005 ¹⁰⁴	Worker	Ironworkers	981	Adults	Tendonitis	19	n/a
Sansone, 2015 ³⁷	Worker	Female cashier	199	Adults	RC calcific tendinopathy (US)	22.6	n/a
Cunha-Miranda, 2010 ³⁴	Worker	Workers	410,496	Adults	Shoulder tendonitis	0.6	n/a
					Elbow tendonitis	0.3	
					Lower limb tendonitis	0.1	
Werner, 2002 ¹⁰⁵	Worker	Dental hygienists	305	Adults	Shoulder tendinitis	13	n/a
					Elbow tendinitis	6	
Gold, 2009 ¹⁰⁶	Worker	Automobile manufacturing workers	1214	Adults	LE	3.3	n/a
					ME	2.2	
					RC tendonitis	12	
Pullopdisakul, 2013 ¹⁰⁷	Worker	Workers	591	Adults	LE	3.4	n/a
					ME	1.7	
Nordander, 2009 ¹⁰⁸	Worker	Workers	2677	Adults	Supraspinatus tendonitis	4.4	n/a
					Infraspinatus tendonitis	3	
					Bicipital tendonitis	3.8	
					LE	2.3	
					ME	1.2	
Silverstein, 2006 ¹⁰⁹	Worker	Workers	436	Adults	RC tendinitis	4.4–7.6	2.9–5.5/100 PY
Ozdolap, 2013 ¹⁷	Worker	Coal miners	80	Adults	LE	41.2	n/a
					ME	12.5	
Werner, 2005 ³⁸	Worker	Dental hygiene students	343	Adults	Upper extremity tendinitis	5	n/a
		Clerical workers	164			12	
Fan, 2014 ¹¹⁰	Worker	Workers	607	Adults	Epicondylitis	6	7.9/100 PY
					LE	5	5.1/100 PY
					ME	2	2.4/100 PT
Garg, 2014 ¹¹¹	Worker	Workers	536	Adults	LE	7.3	3.67/100 PY
Alexandre, 2011 ¹¹²	Worker	Dentist	173,094	Adults	Tendinitis	8.7	n/a
		Physicians				5.6	
		Lawyers				5.5	
		General population				3.2	
Herquelot, 2013 ¹¹³	Worker	Workers	3710	Adults	LE	n/a	1.0/100 PY
Werner, 2005 ¹¹⁴	Worker	Workers	501	Adults	Upper extremity tendonitis	n/a	4.5% (annual CI)
Fan, 2014 Feb ¹¹⁵	Worker	Workers	611	Adults	LE	n/a	4.9/100 PY
Descatha, 2013 ¹¹⁶	Worker	Workers	699	Adults	Epicondylitis	n/a	6.9% (36 mo CI)
					LE		4.9% (36 mo CI)
					ME		4.3% (36 mo CI)

Table 1 (continued)

1 st Author, year, Ref	Group	Cohort	N	Age	Type of tendinopathy	Prevalence	Incidence
McGaughey, 2003 ¹¹⁷	Worker	Expeditioners	292.3 PY	Adults	AT tendonitis	n/a	9.2/100 PY
Barber Foss, 2012 ¹¹⁸	Athletes	Female basketball players	419	Adolescent	SLJ	5	n/a
Tenforde, 2011 ¹²	Athletes	High school athletes	748	Adolescent	AT tendonitis	7.8	n/a
Emerson, 2010 ¹¹⁹	Athletes	Elite gymnasts	40	Adolescent	AT tendinopathy	15	n/a
Steinberg, 2011 ¹²⁰	Athletes	Nonprofessional female dancers	1336	Adolescent	Ankle & foot tendonitis	18.8	n/a
Cassel, 2015 ¹⁰	Athletes	Adolescent athletes	760	Adolescent	AT tendinopathy PT tendinopathy AT rupture	1.8 5.6 0.1	n/a
Gisslen, 2005 ¹²¹	Athletes	Swedish elite junior volleyball players	57	Adolescent	Jumper's knee PT (US)	21 28.9	n/a
Le Gall, 2007 ¹²²	Athletes	Early maturing athletes	233	Adolescent	Tendinopathy	n/a	0.06/1000 AE 0.02/1000 AE
Barber Foss, 2014 ¹²³	Athletes	Late maturing athletes	268	Adolescent	SLJ	n/a	0.3/1000 AE
Beachy, 2014 ¹²⁴	Athletes	Female middle school athletes	14,038	Adolescent	Tendinitis	n/a	0.7/1000 AE
Leanderson, 2011 ¹²⁵	Athletes	Middle school athletes	476	Adolescent	Foot tendinosis Jumper's knee Tendonitis genu Tendinosis groin PT tendinitis	n/a	11.8% (7 y CI) 6.5% (7 y CI) 5.2% (7 y CI) 8.6% (7 y CI) 30.6% (6 y CI)
Hickey, 1997 ¹²⁶	Athletes	Ballet dancers	49	Adolescent	Jumper's knee AT tendinitis PT tendinopathy	n/a	8.1 (5 y CI) 2.1 (5 y CI) 0.1/1000 h
Dubravcic-Simunjak, 2003 ¹²⁷	Athletes	Elite female basketball players	469	Adolescent		(season prevalence)	
Hagglund, 2011 ¹²⁸	Athletes	Junior figure skaters	2229	Adults		2.4	
Buda, 2013 ¹¹	Athletes	Elite male soccer players	144	Adults	AT tendinitis	12.5	n/a
Pieber, 2012 ⁵	Athletes	Climbers	193	Adults	Epicondylitis	13.1	n/a
Durcan, 2014 ¹²⁹	Athletes	Climbers	83	Adults	PT tendinopathy	13.3	n/a
Lian, 2005 ⁸	Athletes	Elite rugby academies	613	Adults	Jumper's knee	14.2	n/a
McCarthy, 2013 ⁷	Athletes	Elite athletes	496	Adults	PT tendinitis	17	n/a
van der Worp, 2011 ⁹	Athletes	Women's basketball	1505	Adults	Patellar tendinopathy	17.8	n/a
Cook, 1998 ¹³⁰	Athletes	Basketball & volleyball players	160	Adults			
Lopes, 2009 ³	Athletes	Elite athletes	434	Adults	PT (US) Tendinopathy	22 22.4	n/a n/a
Wang, 2001 ⁴	Athletes	Athletes referred to PT	59	Adults	RC tendinitis	23.7	n/a
Monteleone, 2014 ³³	Athletes	Elite volleyball athletes	53	Adults	RC (US)	30	n/a
Longo, 2011 ¹³	Athletes	Elite beach volleyball players	174	Adults Elderly	PT tendinopathy	46.6	n/a
Rooks, 1995 ¹³¹	Athletes	Veteran track & field athlete	39	Adults	Upper extremity tendinitis	50	n/a
Walls, 2010 ¹³²	Athletes	Rock climbers	18	Adults	AT tendinopathy (MRI)	78	n/a
Hagemann, 2004 ¹³³	Athletes	Professional dancers	52	Adults	Supraspinatus (MRI) Supraspinatus tear (MRI) Subscapularis (MRI) Subscapularis tear (MRI)	11.5 7.7 1.9 1.9	n/a
Reuter, 2008 ¹³⁴	Athletes	Marathon kayakers	23	Adults	RC partial tear Shoulder tendinopathy	22 43	n/a
Hadala, 2009 ¹³⁵	Athletes	Ironman triathletes	30	Adults	Epicondylitis Biceps brachii tendinitis Shoulder tendinopathy	30 3.3 13.3	n/a

(continued on next page)

Table 1 (continued)

1 st Author, year, Ref	Group	Cohort	N	Age	Type of tendinopathy	Prevalence	Incidence
Comin, 2013 ⁴⁰	Athletes	Ballet dancers	79	Adults	AT (US) PT (US)	8.9 8.9	n/a
Marshall, 2007 ¹³⁶	Athletes	Collegiate women's softball athletes	9389	Adults	Shoulder tendinitis Elbow tendinitis	n/a	0.12/1000 AE 0.04/1000 AE
Krupnick, 1998 ¹³⁷	Athletes	White water paddlers	54	Adults	Tendonitis	n/a	0.19/100 AE
Kelly, 2004 ¹³⁸	Athletes	Elite football quarterbacks	1534	Adults	Biceps tendinitis	n/a	0.5/100 AE
Parekh, 2009 ¹³⁹	Athletes	National Football League	n/a	Adults	AT rupture	n/a	0.9% (per game CI)
Heir, 1996 ¹⁴⁰	Athletes	Military conscripts	6488	Adults	AT tendinitis Shoulder tendinitis	n/a	13.5/1000 conscript-mo 2.1/1000 conscript-mo
Wolf, 2010 ¹⁴¹	Athletes	US military population	n/a	Adults	LE ME	n/a	2.3/1000 PY 0.8/1000 PY
McFarland, 1998 ¹⁴²	Athletes	Collegiate baseball players	12,828 AE	Adults	RC tendinitis	n/a	3.4/1000 AE
White, 2007 ¹⁴³	Athletes	US army soldiers	93,224 AE	Adults	Tendon rupture	n/a	5.6/100 AE
Milgrom, 2003 ¹⁴⁴	Athletes	Male infantry recruits	1405	Adults	AT tendinopathy	n/a	6.8% (4 period each 14 wk CI)
McMahon, 2014 ⁶	Athletes	Elite athletes	141	Elderly	RC tendinosis RC partial tear RC complete rupture	16.3 48.2 21.3	n/a
Kettunen, 2011 ¹⁴⁵	Athletes	Former elite male athletes	785	Elderly	Shoulder tendinopathy Shoulder tendon rupture	n/a	33% (lifetime CI) 19% (lifetime CI)
Kujala, 2005 ⁴¹	Athletes	Former elite male athletes	785	Elderly	AT rupture AT tendinopathy	n/a	8.3% (lifetime CI) 23.9% (lifetime CI)
Njobvu, 2006 ²⁹	Patients	HIV positive patients	65	Adults	Tendinitis	3.1	n/a
Cannon, 2007 ¹⁴⁶	Patients	Cervical radiculopathy with upper limb symptoms	191	Adults	LE	4.7	n/a
Hautmann, 2014 ¹⁴⁷	Patients	Patients with painful heel	101	Adults	AT tendinitis	11.9	n/a
Frey, 2007 ³¹	Patients	Overweight or obese	738	Adults	Ankle & foot tendinitis	16.7	n/a
Finley, 2004 ³²	Patients	Manual wheelchair users	52	Adults	Biceps tendonitis	30.1	n/a
Baumann, 2008 ¹⁴⁸	Patients	Diagnostic shoulder arthroscopies	1007	Adults	Shoulder tendinitis Shoulder partial tear	1.5 0.8	n/a
Chhajed, 2002 ¹⁴⁹	Patients	Lung transplant recipients treated with ciprofloxacin	101	Adults	AT tendonitis AT rupture	15.8 5.9	n/a
Ramirez, 2014 ¹⁵⁰	Patients	Patients with greater trochanteric pain	107	Adults	Gluteus medius tendinosis Gluteus minimus tendinosis	36.4 67.3	n/a
Taunton, 2002 ¹⁵¹	Patients	Patients with running related injury	2002	Adults	PT tendinopathy AT tendinopathy	4.2 4.8	n/a
Bird, 2001 ¹⁵²	Patients	Patients with greater trochanteric pain	24	Adults	Gluteus medius tear Gluteus medius tendinitis	45.8 37.5	n/a
Shah, 2008 ³⁰	Patients	Stroke patients with painful shoulder	89	Adults	RC tendinopathy RC tear	53 35	n/a
Pong, 2012 ¹⁵³	Patients	Stroke patients with hemiplegic shoulders	76	Adults	Shoulder tendinopathy	68.4 Acute 80.3 Chronic	n/a
Kingzett-Taylor, 1999 ¹⁵⁴	Patients	Patients with buttock, lateral hip, or groin pain	250	Adults	Gluteal tear Gluteal tendinosis	8.8 5.2	n/a
Chung, 2013 ¹⁵⁵	Patients	Nurses with musculoskeletal disorder	3914	Adults	ME LE	n/a	0.25% (1 y CI) 0.58% (1 y CI)
Barge-Caballero, 2008 ¹⁵⁶	Patients	Heart transplant patients under quinolones	149	Adults	AT tendinopathy AT rupture	n/a	9.4% (11 y CI) 2% (11 y CI)
Ramos, 2009 ¹⁵⁷	Patients	Patients with knee pain	318	All	PT tendinopathy	32.3	n/a

Table 1 (continued)

1 st Author, year, Ref	Group	Cohort	N	Age	Type of tendinopathy	Prevalence	Incidence
Helliwell, 2003 ¹⁵⁸	Patients	Patients with soft tissue disorders	1382	All	Shoulder tendinitis	11.3	n/a
					LE	6.3	
Sode, 2007 ¹⁵⁹	Patients	First time fluoroquinolone users	28262	All	AT rupture	n/a	0.02 (90 d CI)
Zakaria, 2014 ¹⁶⁰	Patients	Diabetes patients	1296	Elderly	Tendon rupture	n/a	5.21/1000 PY

AE = athlete exposure; AT = Achilles tendon; CI = cumulative incidence; h = hours; LE = lateral epicondylitis; ME = medial epicondylitis; MRI = magnetic resonance imaging diagnosed; N = sample/cohort size; n/a = not available; PT = patellar tendon; PY = person-years; RC = rotator cuff; SLJ = Sinding-Larsen-Johansson; US = ultrasound diagnosed; y = years.

General population

In the general population, the prevalence of clinically diagnosed rotator cuff tendinopathy in adults was reported to range from 2% to 3.8%,^{18,19} with a marginally higher prevalence observed in the elderly population at 5–7%.²⁰ Although it may not be accurate to compare values from different studies, it is worth noting that the value in the general population approaches that of the worker cohorts. In the elbow joint, prevalence of lateral epicondylitis and medial epicondylitis in European cohorts were reported at 0.7–1.3% and 0.3–0.6%, respectively.^{19,21–23} Two outliers were reported on lateral epicondylitis in a Japanese mountain village cohort at 3.8% and on epicondylitis in the Indonesian general population at 6.6%.^{24,25} The divergence in values in these groups suggest that societal aspects also come into play and may be influenced by environmental, cultural, or economic differences among societies. Age and gender do not seem to influence tendinopathy within this cohort. Although upper extremity tendinopathy has been more frequently studied in the general population, less common conditions have also been observed, such as retropharyngeal tendinitis (0.5/100,000 person-years) triceps tendon tears (3.8%)^{26,27} and bicipital tendinitis (0.3%–0.5%).^{19,28}

Comorbidity cohorts

Nineteen studies reported tendinopathy in cohorts with other associated conditions such as HIV positive patients,²⁹ stroke survivors,³⁰ obese cohorts,³¹ wheelchair users,³² etc. (Table 1). Some studies do not investigate the association of tendinopathy with other disease conditions, but instead report tendinopathy as part of a group of patients with musculoskeletal complaints in general. A general trend of increased prevalence is seen when compared with the general population; however due to the variety of conditions, and how they impact tendinopathy, the data cannot be utilised to assess the prevalence of tendinopathy within this cohort. These studies provide evidence that there are intrinsic risk factors for tendinopathy and research into the relationship between them would be worthwhile in understanding the aetiology of tendinopathy.

Summary of systematic review

There is currently a gap in the available evidence on incidence rates as most studies carried out on the general

population are on tendinopathic ruptures only. The actual prevalence of tendinopathy may be higher due to diagnosis. Clinical diagnosis is the main diagnostic technique, with radiological imaging, such as magnetic resonance imaging (MRI) and ultrasound being used to support the diagnosis. Studies that defined tendinopathy using radiological imaging revealed a higher incidence rate compared with studies that used clinical evaluation only.^{33–39} This discrepancy is caused by the inclusion of patients who did not present with symptoms at the time of examination, suggesting that asymptomatic patients are left unnoticed in tendinopathy diagnosis.⁴⁰ This is important since the lifetime cumulative incidence of retired elderly athletes is approximately 25%, suggesting that symptoms may develop later than when the injury was sustained.⁴¹

In summary, specific types of tendinopathy are more prevalent in the different groups. Epicondylitis and rotator cuff tendinopathy were preferentially investigated in workers and the general population, with workers having a higher prevalence and incidence of tendinopathy. Patellar tendinopathy was more frequently investigated in the athlete population, revealing a higher prevalence and incidence when compared with the other groups. Finally, age and sex does not seem to play a factor in tendinopathy.

Effects of tendinopathy on quality-of-life and cost-effectiveness of treatment

A community-based survey compared the socio-economic impacts of four musculoskeletal conditions including tendinitis (tendinopathy), rheumatoid arthritis, osteoarthritis, and lower back pain. Although tendinopathy was found to be less influential to work loss,⁴² shoulder tendinopathy took approximately 10 months to heal and workers take greater amounts of sick leave to recover,^{43–45} report being less productive at work,⁹ and require workers' compensation for disease.^{46,47} Finally, even though patients may return to work within 6 weeks following operative repair, recovery may take a few months.⁴⁸ The burden placed on daily activities cannot be ignored, with one study claiming that about a quarter of patients with tennis elbow (epicondylitis) reported difficulty in activities such as dressing, carrying objects, driving, and sleeping.⁴⁹ The indirect costs can reach great amounts in terms of productivity loss and worker's compensation. Up to 5% of patients with lateral epicondylitis have claimed sickness absence with an average duration of 29 days in a year.⁴⁴ Thus, absenteeism (in the working population aged 25–64) due to

lateral epicondylitis in the United Kingdom alone is estimated to cost £27 million using 2012 global population statistics and median wage.^{44,50,51} Productivity-loss and disease compensation associated with tendinopathy are remarkable, and the high prevalence of tendinopathy, as revealed by the search results on tendinopathy prevalence, suggests that the disease burden may be greater than currently understood.

The goals of tendinopathy treatment are pain reduction, recurrence prevention, and return to sports or preinjury functionality. Treatment aims to remain conservative with oral nonsteroidal anti-inflammatory drugs, corticosteroid injections, and physical therapy as the mainstay in tendinopathy management. Other therapies include injections of platelet-rich plasma or autologous blood. Finally, failure of conservative treatments leads to surgical intervention to excise the tendinopathic tissue and repair the ruptured tendon.^{52,53} Yet the cost analysis on various tendinopathy treatments is inadequate. Direct outpatient medical costs were reported as ranging from €430/patient for corticosteroid injection to €921/patient for physical therapy, for lateral epicondylitis (currency in 2004).⁵⁴ Repeated medical visits are also a concern as lateral epicondylitis is recurrent, and almost half of those affected have seen their general practitioner within the past 12 months.⁴⁴ Cost/quality-adjusted-life-years for physical therapy and corticosteroid injection were £18,962 and £20,518, respectively, values which fall within the benchmark of £20,000 to £30,000 (currency in 2015) and are comparable to the common drug treatments for osteoarthritis and osteoporosis (currency in 2005 and 2004, respectively).^{55–57} Economic evaluations on other tendinopathic conditions are lacking and research on this aspect would be valuable.

Documentation and awareness of tendinopathy

Although tendinopathy is well-recognised in the academic field as listed in the medical subject headings, only tendinitis and spontaneous tendon ruptures are stated within the current version of International Classification of Diseases (ICD) by the World Health Organization (WHO), but tendinosis and tendinopathy are absent.⁵⁸ Tendinitis and tendinosis, continue to be mainstay diagnostic terms, but as tendinopathy has become the accepted term within the medical field, it should be similarly recognised by the public. Healthcare organisations, such as WHO and the Centre for Disease Control (CDC), and orthopaedic organisations, such as the Bone Joint Decade (BJD) and the Fracture Fragility Network (FFN), do not have definitions or information for the term “tendinopathy”. Evidence from our search has demonstrated that despite clinical diagnosis being the mainstay diagnostic technique, MRI and ultrasound are favourable, particularly for asymptomatic patients. Implementation of a standardised, radiological technique, would allow for the inclusion of symptomatic patients, asymptomatic patients, and patients with ruptures to be recognised under the definition of tendinopathy.

Failure to recognise and report an incident, and failure to seek medical attention, amongst other factors may lead to two thirds of tendinopathy cases going unreported, thus the

proportion of individuals with tendinopathy may be higher than reported.⁵⁹ Tendinopathy appears to be particularly prevalent in productive populations that actively contribute to societal development, such as athletes and workers. Despite these indications that tendinopathy may be highly prevalent in society, it remains an under-recognised disease.

Osteoarthritis and osteoporosis are well recognised and studied by researchers, WHO, CDC, and are also key topics in BJD and FFN. These diseases are particularly prevalent in the elderly, thus, their impact on productive demographics may be lower, when compared with tendinopathy.^{60,61} The National Coalition for Osteoporosis and Related Bone Diseases (with support from WHO) and The Arthritis Foundation (with support from CDC) have published action plans to address insufficiencies in tackling osteoporosis and osteoarthritis, respectively.^{62,63} The Australian government published their own action plan (similar to the aforementioned plans) to tackle osteoporosis, osteoarthritis, and rheumatoid arthritis.⁶⁴ These action plans provide a framework to implement research, prevention, treatment, and education within the public. Internet resources are playing a greater role in how the public recognises diseases. Both CDC and WHO have published data on recognising symptoms, prevention, and treating both osteoarthritis and osteoporosis, targeted toward the general public for easy access to information. Our search results have demonstrated that tendinopathy is not well-documented in relation to other diseases; however, the awareness of risk-factors of osteoporosis and osteoarthritis are well-recognised, such as obesity and diabetes, allowing for better disease prevention. The International Osteoporosis Foundation and National Osteoporosis Foundation conducted a study to determine the global prevalence of osteoporosis, and the North Staffordshire Osteoarthritis Project conducted a census to determine the prevalence and impact of osteoarthritis.^{65,66} These studies demonstrate that greater awareness, leads to support by well-established organisations, and aid in determining disease prevalence and impact. Current healthcare registries are a useful tool in tracking and studying diseases, and this has helped study the prevalence of osteoporosis through hip fracture and osteoarthritis through total knee replacements.⁶⁷ Thus, this implores the question as to why tendinopathy does not receive similar awareness and action, when the prevalence may be similar to osteoarthritis and osteoporosis.

Determining the true prevalence of tendinopathy is the first step in studying the impact that tendinopathy has on society, and for this, national health registries are a useful tool, with Sweden, Finland, and The Netherlands using their own registries to study the prevalence of tendon rupture.^{68–70} However, in order for national health registries to run effectively, they require standardised nomenclature and diagnostics. For example, the Swedish Hospital Discharge Registry utilises the ICD in their system, yet without the recognition of tendinopathy in the database, incidences are not recorded. There is currently no organisation specialising in raising awareness for tendinopathy. Through such an organisation of specialists in this field, we may present a greater front in establishing these

standardised definitions of tendinopathy, and having established this, it is necessary to approach international organisations such as WHO, CDC, BJD, and FFN, to gain recognition of the disease in professional fields, as well as have the term properly recognised by the ICD. Establishing these foundational aspects, tendinopathy may be better recognised by the public, patients may be encouraged to seek earlier medical attention, resources will be appropriately allocated to alleviate the burden of tendinopathy, and conclusive studies on the prevalence and socio-economic impact of tendinopathy can be implemented.

Conclusion

The definition of tendinopathy is variable, making proper documentation difficult. Tendinopathy should be defined using widely accepted criteria used by professionals, to include symptomatic, asymptomatic, and rupture patients. Our search results demonstrate that tendinopathy is prevalent in a variety of demographics, particularly in younger generations that are most active in society, yet the public awareness is low. By encouraging awareness in both the professional and public fields, we will enhance our understanding and make appropriate changes in how to tackle the disease.

These proposed changes will be slow, and require persistent effort from experts in the field of tendinopathy. Furthermore, the capacity to make such changes varies widely around the globe, in which some societies may not be able to implement the same systems or interventions as others. However, through these actions we may be able to enhance global awareness of the disease and relieve the burden tendinopathy currently places on society.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

Funding/support

No financial or material support of any kind was received for the work described in this article.

References

- Kader D, Mosconi M, Benazzo F, et al. Achilles tendon rupture. In: Maffulli N, Renstrom P, Leadbetter WB, eds. *Tendon Injuries*. London: Springer London; 2005:187–200.
- Athanasopoulos S, Kapreli E, Tsakoniti A, et al. The 2004 Olympic games: physiotherapy services in the Olympic Village polyclinic. *Br J Sports Med*. 2007;41(9):603–609.
- Lopes AD, Barreto HJ, Aguiar RC, et al. Brazilian physiotherapy services in the 2007 Pan-American Games: injuries, their anatomical location and physiotherapeutic procedures. *Phys Ther Sport*. 2009;10:67–70.
- Wang HK, Cochrane T. A descriptive epidemiological study of shoulder injury in top level English male volleyball players. *Int J Sports Med*. 2001;22:159–163.
- Pieber K, Angelmaier L, Csapo R, et al. Acute injuries and overuse syndromes in sport climbing and bouldering in Austria: a descriptive epidemiological study. *Wien Klin Wochenschr*. 2012;124:357–362.
- McMahon PJ, Prasad A, Francis KA. What is the prevalence of senior-athlete rotator cuff injuries and are they associated with pain and dysfunction? *Clin Orthop Relat Res*. 2014;472:2427–2432.
- McCarthy MM, Voos JE, Nguyen JT, et al. Injury profile in elite female basketball athletes at the Women's National Basketball Association combine. *Am J Sports Med*. 2013;41:645–651.
- Lian OB, Engebretsen L, Bahr R. Prevalence of jumper's knee among elite athletes from different sports: a cross-sectional study. *Am J Sports Med*. 2005;33:561–567.
- van der Worp H, van Ark M, Roerink S, et al. Risk factors for patellar tendinopathy: a systematic review of the literature. *Br J Sports Med*. 2011;45:446–452.
- Cassel M, Baur H, Hirschi Müller A, et al. Prevalence of Achilles and patellar tendinopathy and their association to intratendinous changes in adolescent athletes. *Scand J Med Sci Sports*. 2015;25:e310–e318.
- Buda R, Di Caprio F, Bedetti L, et al. Foot overuse diseases in rock climbing: an epidemiologic study. *J Am Podiatr Med Assoc*. 2013;103:113–120.
- Tenforde AS, Sayres LC, McCurdy ML, et al. Overuse injuries in high school runners: lifetime prevalence and prevention strategies. *PM R*. 2011;3:125–131. quiz 131.
- Longo UG, Rittweger J, Garau G, et al. Patellar tendinopathy in master track and field athletes: influence of impact profile, weight, height, age and gender. *Knee Surg Sports Traumatol Arthrosc*. 2011;19:508–512.
- Longo UG, Rittweger J, Garau G, et al. No influence of age, gender, weight, height, and impact profile in achilles tendinopathy in masters track and field athletes. *Am J Sports Med*. 2009;37:1400–1405.
- Roquelaure Y, Ha C, Leclerc A, et al. Epidemiologic surveillance of upper-extremity musculoskeletal disorders in the working population. *Arthritis Rheum*. 2006;55:765–778.
- Auerbach JD, Weidner ZD, Milby AH, et al. Musculoskeletal disorders among spine surgeons: results of a survey of the Scoliosis Research Society membership. *Spine (Phila Pa 1976)*. 2011;36:E1715–E1721.
- Ozdolap S, Emre U, Karamercan A, et al. Upper limb tendinitis and entrapment neuropathy in coal miners. *Am J Ind Med*. 2013;56:569–575.
- Miranda H, Viikari-Juntura E, Heistaro S, et al. A population study on differences in the determinants of a specific shoulder disorder versus nonspecific shoulder pain without clinical findings. *Am J Epidemiol*. 2005;161:847–855.
- Shiri R, Varonen H, Heliövaara M, et al. Hand dominance in upper extremity musculoskeletal disorders. *J Rheumatol*. 2007;34:1076–1082.
- Chard MD, Hazleman BL. Shoulder disorders in the elderly (a hospital study). *Ann Rheum Dis*. 1987;46:684–687.
- Shiri R, Viikari-Juntura E, Varonen H, et al. Prevalence and determinants of lateral and medial epicondylitis: a population study. *Am J Epidemiol*. 2006;164:1065–1074.
- Walker-Bone K, Palmer KT, Reading I, et al. Prevalence and impact of musculoskeletal disorders of the upper limb in the general population. *Arthritis Rheum*. 2004;51:642–651.
- Salaffi F, De Angelis R, Grassi W. Prevalence of musculoskeletal conditions in an Italian population sample: results of a regional community-based study. I. The MAPPING study. *Clin Exp Rheumatol*. 2005;23:819–828.
- Tajika T, Kobayashi T, Yamamoto A, et al. Prevalence and risk factors of lateral epicondylitis in a mountain village in Japan. *J Ortho Surg*. 2014;22:240–243.
- Darmawan L, Valkenburg HA, Muirden KD, et al. The prevalence of soft tissue rheumatism. *Rheumatol Int*. 1995;15:121–124.
- Horowitz G, Ben-Ari O, Brenner A, et al. Incidence of retropharyngeal calcific tendinitis (longus colli tendinitis) in the general population. *Otolaryngol Head Neck Surg*. 2013;148:955–958.
- Koplas MC, Schneider E, Sundaram M. Prevalence of triceps tendon tears on MRI of the elbow and clinical correlation. *Skeletal Radiol*. 2011;40:587–594.
- Alvarez-Nemegyei J, Peláez-Ballestas I, Rodríguez-Amado J, et al. Prevalence of rheumatic regional pain syndromes in adults from Mexico:

- a community survey using COPCORD for screening and syndrome-specific diagnostic criteria. *J Rheumatol.* 2011;86:15–20.
29. Njobvu P, McGill P. Soft tissue rheumatic lesions and HIV infection in Zambians. *J Rheumatol.* 2006;33:2493–2497.
 30. Shah RR, Haghighpanah S, Elovic EP, et al. MRI findings in painful post-stroke shoulder. *Stroke.* 2008;39:1808–1813.
 31. Frey C, Zamora J. The effects of obesity on orthopaedic foot and ankle pathology. *Foot Ankle Int.* 2007;28:996–999.
 32. Finley MA, Rodgers MM. Prevalence and identification of shoulder pathology in athletic and nonathletic wheelchair users with shoulder pain: a pilot study. *J Rehabil Res Dev.* 2004;41:395–402.
 33. Monteleone G, Tramontana A, McDonald K, et al. Ultrasonographic evaluation of the shoulder in elite Italian beach volleyball players. *J Sports Med Physical Fitness.* 2014.
 34. Cunha-Miranda L, Carnide F, Lopes MF. Prevalence of rheumatic occupational diseases - PROUD study. *Acta Reumatol Port.* 2010;35:215–226.
 35. Fairley J, Toppi J, Cicuttini FM, et al. Association between obesity and magnetic resonance imaging defined patellar tendinopathy in community-based adults: a cross-sectional study. *BMC Musculoskelet Disord.* 2014;15:266–272.
 36. Girish G, Lobo LG, Jacobson JA, et al. Ultrasound of the shoulder: asymptomatic findings in men. *AJR Am J Roentgenol.* 2011;197:W713–W719.
 37. Sansone VC, Meroni R, Boria P, et al. Are occupational repetitive movements of the upper arm associated with rotator cuff calcific tendinopathies? *Rheumatol Int.* 2015;35:273–280.
 38. Werner RA, Franzblau A, Gell N, et al. Prevalence of upper extremity symptoms and disorders among dental and dental hygiene students. *J Calif Dent Assoc.* 2005;33:123–131.
 39. Zwerver J, Bredeweg SW, van den Akker-Scheek I. Prevalence of Jumper's knee among nonelite athletes from different sports: a cross-sectional survey. *Am J Sports Med.* 2011;39:1984–1988.
 40. Comin J, Cook JL, Malliaras P, et al. The prevalence and clinical significance of sonographic tendon abnormalities in asymptomatic ballet dancers: a 24-month longitudinal study. *Br J Sports Med.* 2013;47:89–92.
 41. Kujala UM, Sarna S, Kaprio J. Cumulative incidence of achilles tendon rupture and tendinopathy in male former elite athletes. *Clin J Sport Med.* 2005;15:133–135.
 42. Kramer JS, Yelin EH, Epstein WV. Social and economic impacts of four musculoskeletal conditions. A study using national community-based data. *Arthritis Rheum.* 1983;26:901–907.
 43. Bonde JP, Mikkelsen S, Anderson JH, et al. Prognosis of shoulder tendonitis in repetitive work: a follow up study in a cohort of Danish industrial and service workers. *Occ Environ Med.* 2003;60:E8.
 44. Walker-Bone K, Palmer KT, Reading I, et al. Occupation and epicondylitis: a population-based study. *Rheumatology (Oxford).* 2012;51:305–310.
 45. Serazin C, Ha C, Bodin J, et al. Employment and occupational outcomes of workers with musculoskeletal pain in a French region. *Occ Environ Med.* 2013;70:143–148.
 46. Grewal R, MacDermid JC, Shah P, et al. Functional outcome of arthroscopic extensor carpi radialis brevis tendon release in chronic lateral epicondylitis. *J Hand Surg-Am Vol.* 2009;34A:849–857.
 47. Palmer KT, Harris EC, Coggon D. Compensating occupationally related tenosynovitis and epicondylitis: a literature review. *Occ Med-Oxford.* 2007;57(1):67–74.
 48. Seil R, Litztenburger H, Kohn D, et al. Arthroscopic treatment of chronically painful calcifying tendinitis of the supraspinatus tendon. *Arthroscopy-J Arthroscopic Related Surg.* 2006;22:521–527.
 49. Alizadehkhayat O, Fisher AC, Kemp GJ, et al. Pain, functional disability, and psychologic status in tennis elbow. *Clin J Pain.* 2007;23:482–489.
 50. Distribution of Median and Mean Income and Tax by Age Range and Gender. Personal Income by Tax Year 2015. Available from: <https://www.gov.uk/government/statistics/distribution-of-median-and-mean-income-and-tax-by-age-range-and-gender-2010-to-2011>. [accessed 12.02.16].
 51. Population, total. 2011–2015. Available from: <http://data.worldbank.org/indicator/SP.POP.TOTL> [accessed 12.02.16].
 52. Witvrouw E, Bellemans J, Lysens R, et al. Intrinsic risk factors for the development of patellar tendinitis in an athletic population – a two-year prospective study. *Am J Sports Med.* 2001;29:190–195.
 53. Dragoo JL, Wasterlain AS, Braun HJ, et al. Platelet-rich plasma as a treatment for patellar tendinopathy: a double-blind, randomized controlled trial. *Am J Sports Med.* 2014;42:610–618.
 54. Korthals-de Bos IBC, Smidt N, van Tulder MW, et al. Cost effectiveness of interventions for lateral epicondylitis – results from a randomised controlled trial in primary care. *Pharmacoeconomics.* 2004;22:185–195.
 55. Coombes BK, Connelly L, Bisset L, et al. Economic evaluation favours physiotherapy but not corticosteroid injection as a first-line intervention for chronic lateral epicondylalgia: evidence from a randomised clinical trial. *Br J Sports Med.* 2015.
 56. Stevenson M, Jones ML, De Nigris E, et al. A systematic review and economic evaluation of alendronate, etidronate, risedronate, raloxifene and teriparatide for the prevention and treatment of postmenopausal osteoporosis. *Health Technol Ass.* 2005;9:1–6.
 57. Yen ZS, Lai MS, Wang CT, et al. Cost-effectiveness of treatment strategies for osteoarthritis of the knee in Taiwan. *J Rheumatol.* 2004;31:1797–1803.
 58. International Classification of Diseases (ICD). Classifications 2015. Available from: <http://www.who.int/classifications/icd/en/>. [accessed 12.02.16].
 59. Geneva International Labour Office. *Safe Work, Estimating the Economic Costs of Occupational Injuries and Illnesses in Developing Countries: Essential Information for Decision Makers.* Geneva, Switzerland: Int Labour Office; 2012.
 60. Osteoarthritis. *Arthritis*; 2015. Available from: <http://www.cdc.gov/arthritis/basics/osteoarthritis.htm> [accessed 12.02.16].
 61. Looker AC, Frenk SM. *Percentage of Adults Aged 65 and over with Osteoporosis or Low Bone Mass at the Femur Neck or Lumbar Spine: United States, 2005–2010.* NCHS Health E-Stat; 2015. Available from: http://www.cdc.gov/nchs/data/hestat/osteoporosis/osteoporosis2005_2010.htm [accessed 12.02.16].
 62. National Coalition for Osteoporosis and Related Bone Diseases. *National Action Plan for Bone Health: Recommendations from the Summit for a National Action Plan for Bone Health.* Washington D.C., North America: National Coalition for Osteoporosis and Related Bone Diseases; 2009.
 63. Arthritis Foundation. *A National Public Health Agenda for Osteoarthritis 2010.* USA: Arthritis Foundation; 2010.
 64. National Arthritis and Musculoskeletal Conditions Advisory Group. *Evidence to Support the National Action Plan for Osteoarthritis, Rheumatoid Arthritis and Osteoporosis: Opportunities to Improve Health-related Quality of Life and Reduce the Burden of Disease and Disability.* Australian Government Department of Health and Ageing. Canberra, Australia: Editor. 2004.
 65. Johnell O, Kanis JA. An estimate of the worldwide prevalence, mortality and disability associated with hip fracture. *Osteoporos Int.* 2004;15:897–902.
 66. Thomas E, Peat G, Harris L, et al. The prevalence of pain and pain interference in a general population of older adults: cross-sectional findings from the North Staffordshire Osteoarthritis Project (NorStOP). *Pain.* 2004;110:361–368.
 67. Delaunay C. Registries in orthopaedics. *Orthop Traumatol Surg Res.* 2015;101(1 Suppl.):S69–S75.
 68. Mattila VM, Huttunen TT, Haapasalo H, et al. Declining incidence of surgery for Achilles tendon rupture follows publication of major RCTs: evidence-influenced change evident using the Finnish registry study. *Br J Sports Med.* 2015;49(16):1084–1086.
 69. Huttunen TT, Kannus P, Rolf C, et al. Acute achilles tendon ruptures incidence of injury and surgery in Sweden between 2001 and 2012. *Am J Sports Med.* 2014;42:2419–2423.
 70. van der Linden PD, Nab HW, Simonian S, et al. Fluoroquinolone use and the change in incidence of tendon ruptures in the Netherlands. *Pharm World Sci.* 2001;23:89–92.
 71. Zapata AL, Moraes AJ, Leone C, et al. Pain and musculoskeletal pain syndromes related to computer and video game use in adolescents. *Eur J Pediatr.* 2006;165:408–414.

72. Rechardt M, Shiri R, Karppinen J, et al. Lifestyle and metabolic factors in relation to shoulder pain and rotator cuff tendinitis: a population-based study. *BMC Musculoskelet Disord*. 2010;11:165.
73. Joseph MF, Trojian TH, Anderson JM, et al. Incidence of morphologic changes in asymptomatic achilles tendons in an active young adult population. *J Sport Rehabil*. 2013;22:249–253.
74. Waldecker U, Hofmann G, Drewitz S. Epidemiologic investigation of 1394 feet: coincidence of hindfoot malalignment and Achilles tendon disorders. *Foot Ankle Surg*. 2012;18:119–123.
75. Schibany N, Zehetgruber H, Kainberger F, et al. Rotator cuff tears in asymptomatic individuals: a clinical and ultrasonographic screening study. *Eur J Radiol*. 2004;51:263–268.
76. Safran MR, Graham SM. Distal biceps tendon ruptures: incidence, demographics, and the effect of smoking. *Clin Orthop Relat Res*. 2002;404:275–283.
77. Ostor AJ, Richards CA, Prevost AT, et al. Diagnosis and relation to general health of shoulder disorders presenting to primary care. *Rheumatology (Oxford)*. 2005;44:800–805.
78. Cretnik A, Kosir R, Kosanovic M. Incidence and outcome of operatively treated achilles tendon rupture in the elderly. *Foot Ankle Int*. 2010;31:14–18.
79. Moller A, Astron M, Westlin N. Increasing incidence of Achilles tendon rupture. *Acta Orthop Scand*. 1996;67:479–481.
80. Clayton RA, Court-Brown CM. The epidemiology of musculoskeletal tendinous and ligamentous injuries. *Injury*. 2008;39:1338–1344.
81. Levi N. The incidence of Achilles tendon rupture in Copenhagen. *Injury*. 1997;28:311–313.
82. Leppilahti J, Puranen J, Orava S. Incidence of Achilles tendon rupture. *Acta Orthop Scand*. 1996;67:277–279.
83. Houshian S, Tscherning T, Riegels-Nielsen P. The epidemiology of Achilles tendon rupture in a Danish county. *Injury*. 1998;29:651–654.
84. Maffulli N, Waterston SW, Squair J, et al. Changing incidence of Achilles tendon rupture in Scotland: a 15-year study. *Clin J Sport Med*. 1999;9:157–160.
85. Suchak AA, Bostick G, Reid D, et al. The incidence of Achilles tendon ruptures in Edmonton, Canada. *Foot Ankle Int*. 2005;26:932–936.
86. de Jonge S, van den Berg C, de Vos RJ, et al. Incidence of midportion Achilles tendinopathy in the general population. *Br J Sports Med*. 2011;45:1026–1028.
87. Nyyssonen T, Luthje, Kroger H. The increasing incidence and difference in sex distribution of Achilles tendon rupture in Finland in 1987–1999. *Scand J Surg*. 2008;97:272–275.
88. McCormack Jr RR, Inman RD, Wells A, et al. Prevalence of tendinitis and related disorders of the upper extremity in a manufacturing workforce. *J Rheumatol*. 1990;17:958–964.
89. Almeida MC, Cezar-Vaz MR, Soares JF, et al. The prevalence of musculoskeletal diseases among casual dock workers. *Rev Lat Am Enfermagem*. 2012;20:243–250.
90. Frost P, Bonde JP, Mikkelsen S, et al. Risk of shoulder tendinitis in relation to shoulder loads in monotonous repetitive work. *Am J Ind Med*. 2002;41:11–18.
91. Descatha A, Leclerc A, Chastang JF, et al. Medial epicondylitis in occupational settings: prevalence, incidence and associated risk factors. *J Occup Environ Med*. 2003;45:993–1001.
92. Fan ZJ, Silverstein BA, Bao S, et al. Quantitative exposure-response relations between physical workload and prevalence of lateral epicondylitis in a working population. *Am J Ind Med*. 2009;52:479–490.
93. Rosenbaum DA, Grzywacz JG, Chen H, et al. Prevalence of epicondylitis, rotator cuff syndrome, and low back pain in Latino poultry workers and manual laborers. *Am J Ind Med*. 2013;56:226–234.
94. Kryger AI, Lassen CF, Andersen JH. The role of physical examinations in studies of musculoskeletal disorders of the elbow. *Occup Environ Med*. 2007;64:776–781.
95. Kaergaard A, Andersen JH. Musculoskeletal disorders of the neck and shoulders in female sewing machine operators: prevalence, incidence, and prognosis. *Occup Environ Med*. 2000;57:528–534.
96. Dimberg L. The prevalence and causation of tennis elbow (lateral humeral epicondylitis) in a population of workers in an engineering industry. *Ergonomics*. 1987;30:573–579.
97. Roto P, Kivi P. Prevalence of epicondylitis and tenosynovitis among meatcutters. *Scand J Work, Environ Health*. 1984;10:203–205.
98. Ono Y, Nakamura R, Shimaoka M, et al. Epicondylitis among cooks in nursery schools. *Occup Environ Med*. 1998;55:172–179.
99. Leclerc A, Landre MF, Chastang JF, et al. Upper-limb disorders in repetitive work. *Scand J Work Environ Health*. 2001;27:268–278.
100. Capone AC, Parikh PM, Gatti ME, et al. Occupational injury in plastic surgeons. *Plast Reconstr Surg*. 2010;125:1555–1561.
101. Ritz BR. Humeral epicondylitis among gas- and waterworks employees. *Scand J Work Environ Health*. 1995;21:478–486.
102. Chiang H, Ko YC, Chen SS, et al. Prevalence of shoulder and upper-limb disorders among workers in the fish-processing industry. *Scand J Work Environ Health*. 1993;19:126–131.
103. Barrero LH, Pulido JA, Berrio S, et al. Physical workloads of the upper extremity among workers of the Colombian flower industry. *Am J Ind Med*. 2012;55:926–939.
104. Forde MS, Punnett L, Wegman DH. Prevalence of musculoskeletal disorders in union ironworkers. *J Occup Environ Hyg*. 2005;2:203–212.
105. Werner RA, Hamann C, Franzblau A, et al. Prevalence of carpal tunnel syndrome and upper extremity tendinitis among dental hygienists. *J Dent Hyg*. 2002;76:126–132.
106. Gold JE, d'Errico A, Katz JN, et al. Specific and non-specific upper extremity musculoskeletal disorder syndromes in automobile manufacturing workers. *Am J Ind Med*. 2009;52:124–132.
107. Pulpodissakul S, Ekpanyaskul C, Taptagaporn S, et al. Upper extremities musculoskeletal disorders: prevalence and associated ergonomic factors in an electronic assembly factory. *Int J Occup Med Environ Health*. 2013;26:751–761.
108. Nordander C, Ohlsson K, Akesson I, et al. Risk of musculoskeletal disorders among females and males in repetitive/constrained work. *Ergonomics*. 2009;52:1226–1239.
109. Silverstein BA, Viikari-Juntura E, Fan ZJ, et al. Natural course of non-traumatic rotator cuff tendinitis and shoulder symptoms in a working population. *Scand J Work Environ Health*. 2006;32:99–108.
110. Fan ZJ, Bao S, Silverstein BA, et al. Predicting work-related incidence of lateral and medial epicondylitis using the strain index. *Am J Ind Med*. 2014;57:1319–1330.
111. Garg A, Kapellusch JM, Hegmann KT, et al. The strain index and TLV for HAL: risk of lateral epicondylitis in a prospective cohort. *Am J Ind Med*. 2014;57:286–302.
112. Alexandre PC, da Silva IC, de Souza LM, et al. Musculoskeletal disorders among Brazilian dentists. *Arch Environ Occup Health*. 2011;66:231–235.
113. Herquelot E, Bodin J, Roquelaure Y, et al. Work-related risk factors for lateral epicondylitis and other cause of elbow pain in the working population. *Am J Ind Med*. 2013;56:400–409.
114. Werner RA, Franzblau A, Gell N, et al. A longitudinal study of industrial and clerical workers: predictors of upper extremity tendonitis. *J Occ Rehabil*. 2005;15:37–46.
115. Fan ZJ, Silverstein BA, Bao S, et al. The association between combination of hand force and forearm posture and incidence of lateral epicondylitis in a working population. *Hum Factors*. 2014;56:151–165.
116. Descatha A, Dale AM, Jaegers L, et al. Self-reported physical exposure association with medial and lateral epicondylitis incidence in a large longitudinal study. *Occup Environ Med*. 2013;70:670–673.
117. McGaughey I, Sullivan P. The epidemiology of knee and ankle injuries on Macquarie Island. *Injury*. 2003;34:842–846.
118. Barber Foss KD, Myer GD, Chen SS, et al. Expected prevalence from the differential diagnosis of anterior knee pain in adolescent female athletes during preparticipation screening. *J Athl Train*. 2012;47:519–524.
119. Emerson C, Morrissey D, Perry M, et al. Ultrasonographically detected changes in Achilles tendons and self reported symptoms in elite gymnasts compared with controls—an observational study. *Man Ther*. 2010;15:37–42.
120. Steinberg N, Siev-Ner I, Peleg S, et al. Injury patterns in young, non-professional dancers. *J Sports Sci*. 2011;29:47–54.
121. Gisslen K, Gyulai C, Soderman K, et al. High prevalence of jumper's knee and sonographic changes in Swedish elite junior volleyball players compared to matched controls. *Br J Sports Med*. 2005;39:298–301.

122. Le Gall F, Carling C, Reilly T. Biological maturity and injury in elite youth football. *Scand J Med Sci Sports*. 2007;17:564–572.
123. Barber Foss KD, Myer GD, Hewett TE. Epidemiology of basketball, soccer, and volleyball injuries in middle-school female athletes. *Phys Sportsmed*. 2014;42:146–153.
124. Beachy G, Rauh M. Middle school injuries: a 20-year (1988–2008) multisport evaluation. *J Athl Train*. 2014;49:493–506.
125. Leanderson C, Leanderson J, Wykman A, et al. Musculoskeletal injuries in young ballet dancers. *Knee Surg Sports Traumatol Arthrosc*. 2011;19:1531–1535.
126. Hickey GJ, Fricker PA, McDonald WA. Injuries of young elite female basketball players over a six-year period. *Clin J Sport Med*. 1997;7:252–256.
127. Dubravac-Simunjak S, Pecina M, Kuipers H, et al. The incidence of injuries in elite junior figure skaters. *Am J Sports Med*. 2003;31:511–517.
128. Hagglund M, Zwerver J, Ekstrand J. Epidemiology of patellar tendinopathy in elite male soccer players. *Am J Sports Med*. 2011;39:1906–1911.
129. Durcan L, Coole A, McCarthy E, et al. The prevalence of patellar tendinopathy in elite academy rugby: a clinical and imaging study. *J Sci Med Sport*. 2014;17:173–176.
130. Cook JL, Khan KM, Harcourt PR, et al. Patellar tendon ultrasonography in asymptomatic active athletes reveals hypoechoic regions: a study of 320 tendons. Victorian Institute of Sport Tendon Study Group. *Clin J Sport Med*. 1998;8:73–77.
131. Rooks MD, Johnston RB, Ensor CD, et al. Injury patterns in recreational rock climbers. *Am J Sports Med*. 1995;23:683–685.
132. Walls RJ, Brennan SA, Hodnett P, et al. Overuse ankle injuries in professional Irish dancers. *Foot Ankle Surg*. 2010;16:45–49.
133. Hagemann G, Rijke AM, Mars M. Shoulder pathoanatomy in marathon kayakers. *Br J Sports Med*. 2004;38:413–417.
134. Reuter RM, Hiller WD, Ainge GR, et al. Ironman triathletes: MRI assessment of the shoulder. *Skeletal Radiol*. 2008;37:737–741.
135. Hadala M, Barrios C. Sports injuries in an America's Cup yachting crew: a 4-year epidemiological study covering the 2007 challenge. *J Sports Sci*. 2009;27:711–717.
136. Marshall SW, Hamstra-Wright KL, Dick R, et al. Descriptive epidemiology of collegiate women's softball injuries: national Collegiate Athletic Association Injury Surveillance System, 1988–1989 through 2003–2004. *J Athl Train*. 2007;42:286–294.
137. Krupnick JE, Cox RD, Summers RL. Injuries sustained during competitive white-water paddling: a survey of athletes in the 1996 Olympic trials. *Wilderness Environ Med*. 1998;9:14–18.
138. Kelly BT, Barnes RP, Powell JW, et al. Shoulder injuries to quarterbacks in the national football league. *Am J Sports Med*. 2004;32:328–331.
139. Parekh SG, Wray 3rd WH, Brimmo O, et al. Epidemiology and outcomes of Achilles tendon ruptures in the National Football League. *Foot Ankle Spec*. 2009;2:283–286.
140. Heir T, Glomsaker. Epidemiology of musculoskeletal injuries among Norwegian conscripts undergoing basic military training. *Scand J Med Sci Sports*. 1996;6:186–191.
141. Wolf JM, Mountcastle S, Burks R, et al. Epidemiology of lateral and medial epicondylitis in a military population. *Mil Med*. 2010;175:336–339.
142. McFarland EG, Wasik M. Epidemiology of collegiate baseball injuries. *Clin J Sport Med*. 1998;8:10–13.
143. White DW, Wenke JC, Mosely DS, et al. Incidence of major tendon ruptures and anterior cruciate ligament tears in US Army soldiers. *Am J Sports Med*. 2007;35:1308–1314.
144. Milgrom C, Finestone A, Zin D, et al. Cold weather training: a risk factor for achilles paratendinitis among recruits. *Foot & Ankle Int*. 2003;24:398–401.
145. Kettunen JA, Kujala U, Sarna S, et al. Cumulative incidence of shoulder region tendon injuries in male former elite athletes. *Int J Sports Med*. 2011;32:451–454.
146. Cannon DE, Dillingham TR, Miao H, et al. Musculoskeletal disorders in referrals for suspected cervical radiculopathy. *Arch Phys Med Rehabil*. 2007;88:1256–1259.
147. Hautmann MG, Neumaier U, Kolbl O. Re-irradiation for painful heel spur syndrome. Retrospective analysis of 101 heels. *Strahlenther Onkol*. 2014;190:298–303.
148. Baumann B, Genning K, Bohm D, et al. Arthroscopic prevalence of pulley lesions in 1007 consecutive patients. *J Shoulder Elbow Surg*. 2008;17:14–20.
149. Chhajed PN, Plit ML, Hopkins PM, et al. Achilles tendon disease in lung transplant recipients: association with ciprofloxacin. *Eur Respir J*. 2002;19:469–471.
150. Ramirez J, Pomes I, Sobrino-Guijarro B, et al. Ultrasound evaluation of greater trochanter pain syndrome in patients with spondyloarthritis: are there any specific features? *Rheumatol Int*. 2014;34:947–952.
151. Taunton JE, Ryan MB, Clement DB, et al. A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med*. 2002;36:95–101.
152. Bird PA, Oakley SP, Shnier R, et al. Prospective evaluation of magnetic resonance imaging and physical examination findings in patients with greater trochanteric pain syndrome. *Arthritis Rheum*. 2001;44:2138–2145.
153. Pong Y, Wang LY, Huang YC, et al. Sonography and physical findings in stroke patients with hemiplegic shoulders: a longitudinal study. *J Rehabil Med*. 2012;44:553–557.
154. Kingzett-Taylor A, Tirman PFJ, Feller J, et al. Tendinosis and tears of gluteus medius and minimus muscles as a cause of hip pain: MR imaging findings. *AJR Am J Roentgenol*. 1999;173:1123–1126.
155. Chung YC, Hung CT, Li SF, et al. Risk of musculoskeletal disorder among Taiwanese nurses cohort: a nationwide population-based study. *BMC Musculoskelet Disord*. 2013;14:144.
156. Barge-Caballero E, Crespo-Leiro MG, Paniagua-Martin MJ, et al. Quinolone-related Achilles tendinopathy in heart transplant patients: incidence and risk factors. *J Heart Lung Transplant*. 2008;27:46–51.
157. Ramos LA, Carvalho RT, Garms E, et al. Prevalence of pain on palpation of the inferior pole of the patella among patients with complaints of knee pain. *Clinics*. 2009;64:199–202.
158. Helliwell PS. Towards epidemiological criteria for soft-tissue disorders of the arm. *Occ Med*. 2003;53:313–319.
159. Sode J, Obel N, Hallas J, et al. Use of fluroquinolone and risk of Achilles tendon rupture: a population-based cohort study. *Eur J Clin Pharmacol*. 2007;63:499–503.
160. Zakaria MH, Davis WA, Davis TM. Incidence and predictors of hospitalization for tendon rupture in type 2 diabetes: the Fremantle diabetes study. *Diabet Med*. 2014;31:425–430.