

## **Annual Standing Orders**

- I<u>nitial</u> next to <u>each</u> OTC medication this patient may take Indicate the NTE (Not To Exceed) as necessary
- Sign and date the bottom of the form

Patient: DOB: Today's Date:				
Initial each below to order	Medication	Instructions	Route	Indication(s)
	Tylenol 325mg	1-2 tablets every 4-6 hours as needed NTE:	PO (By Mouth)	Pain, Fever
	Ibuprofen 200mg	1-2 tablets every 4-6 hours as needed NTE:	PO	Pain
	Pepto-Bismol (Kaopectate)	30mL or 2tabs every 4 hours as needed NTE:	РО	Heartburn, Nausea, Loose Stool
	Imodium 2mg (Loperamide)	1-2 tablets every 8 hours. NTE:	РО	Loose Stool
	Robitussin DM (Guaifenesin)	5-10mL every 4 hours as needed NTE:	РО	Cough
	Milk of Magnesia	30mL once daily as needed NTE:	РО	Constipation
	Menthol Lozenges	1 lozenge every hour as needed NTE:	РО	Sore Throat
	Benadryl 25mg	1-2 tablets every 4-6 hours as needed  Not to be used as a sleep aid.  NTE:	РО	Allergies, Itching, Hives
	TUMS/Maalox 500mg (Calcium Carbonate)	1-2 tablets every 2-4 hours as needed after meal or bedtime NTE:	РО	Pyrosis (Heartburn), Indigestion
	Hydrocortisone Cream 1%	Apply thin layer to affected area as directed:	Topical (Skin)	Rash
	Bacitracin (Antibiotic	Apply thin layer to affected area as	Topical	Minor cuts, scrapes, or

(Skin)

**Edited: October 2024** 

wounds

Provider signature	Date	

directed:

Ointment)